

Spinal Anaesthesia vs General Anaesthesia for Laparoscopic Cholecystectomy: A study in Pabna Medical College Hospital, Pabna, Bangladesh

Dr. Md. Monirul Islam¹, Dr. Md. Abdus Salam^{2*}, Dr. Md. Aatur Rahman³, Dr. Ahmed Arif⁴, Dr. Md. Sirajul Islam⁵, Dr. Md. Faisal Zinnat⁶

^{1,2}Assistant professor (Anesthesiology), OSD, DGHS, Mohakhali, Dhaka Attachment with Pabna Medical College Hospital, Pabna, Bangladesh

³Senior Consultant, Department of Anaesthesiology, 250 Bedded General Hospitals, Pabna, Bangladesh

^{4,5}Junior Consultant, Department of Anaesthesiology, 250 Bedded General Hospitals, Pabna, Bangladesh

⁶Anaesthetist, Department of Anaesthesiology, 250 Bedded General Hospitals, Pabna, Bangladesh

*Corresponding author: Dr. Md. Abdus Salam

| Received: 20.05.2019 | Accepted: 26.05.2019 | Published: 30.05.2019

DOI: [10.36347/sjams.2019.v07i05.049](https://doi.org/10.36347/sjams.2019.v07i05.049)

Abstract

Case Report

Laparoscopic cholecystectomy is considered the "gold standard" for the surgical treatment of gallstone disease. A comparative clinical study was conducted in the department of Anesthesia, Pabna Medical College Hospital, Pabna, Bangladesh during the period from January 2018 to April 2019. The aim of our study was to compare between Spinal Anaesthesia and General Anaesthesia for laparoscopic cholecystectomy in healthy patients. A total of 80 patients aged between 18-65 years of both sex with ASA Grade status I and II undergoing elective laparoscopic Cholecystectomy were selected. After taking written informed consent from the study participants, we were elected 80 study subjects and divided into two groups ; Group I (n=40) received General Anaesthesia. And Group II (n=40) received spinal Anaesthesia. Male and female ratio was 1: 1.96. Study adopted an exclusion criteria which were the patients with ASA grade III and IV high risk patients, all emergency procedures, bleeding disorders, acute cholecystitis, pancreatitis and acute cholangitis, previous open surgery in upper abdomen, contraindications for pneumoperitoneum, cardiovascular disorders, respiratory disorders, renal disease and liver disease, circulatory instability, and patients with known sensitivity to local anesthetics. Pain was significantly less at 4 hours, 8 hours, 12 hours, and 24 hours after the procedure for the Spinal Anesthesia group, compared with those who received General Anesthesia. There was no statistical difference between the two groups regarding complications, hospital stay, recovery, or degree of satisfaction at follow-up. In the conclusion, we can say, Spinal Anaesthesia provides better safety and adequacy in healthy patients and provides better post-operative pain control without limiting the recovery than General Anaesthesia. Post-operative complications like nausea, vomiting, dizziness and pneumonia are less in Spinal Anaesthesia.

Keywords: Spinal Anesthesia, Surgical Treatment, General Anesthesia, Laparoscopic Cholecystectomy.

Copyright © 2019: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

INTRODUCTION

It is surprising that, regional anesthesia has been successfully used for laparoscopic cholecystectomy in patients unfit to have the procedure under General Anesthesia but has not been tested in healthy patients in whom any presumed risk would be theoretically much lower. Hamad and Ibrahim El-Khattary used spinal anesthesia for laparoscopic cholecystectomy for the first time in a small series of healthy patients but they had used nitrous oxide as a pneumoperitoneum instead of standard carbon dioxide. Recently, it has been shown that, laparoscopic cholecystectomy can be done successfully using carbon dioxide pneumoperitoneum under Spinal Anesthesia in healthy patients with symptomatic gallstone disease[1]. Laparoscopic cholecystectomy has become very popular after it was first described in 1987 by Philippe

Mouret in France. Laparoscopic surgical techniques have been rapidly accepted by surgeons worldwide with published reports describing the benefit of less postoperative pain, decreased hospital stay and earlier return to work[2]. Minimally invasive therapy is done with the general aim to minimize the trauma of interventional process whilst still achieving satisfactory result [3]. Johnson [4] noted that, "all laparoscopic procedures are merely a change in access and still require General Anesthetic; hence the difference from conventional surgery is likely to be small." This statement is predominantly based on the assumption that laparoscopy necessitates endotracheal intubation to prevent aspiration and respiratory embarrassment secondary to the induction of carbon dioxide pneumoperitoneum, which is not well tolerated in a patient who is awake during the procedure[5,6]. The

incidence of postoperative morbidity like nausea, vomiting, dizziness, respiratory complication, thromboembolism and pneumonia was much less as compared to General Anesthesia[7] Also, the total cost of Spinal Anesthesia with respect to hospital stay, induction and recovery, the need for postoperative antiemetics and analgesia and the incidence of other complication was much lower when compared to General Anesthesia[8]. This study was planned to assess comparative superiority of Spinal Anaesthesia with General Anaesthesia for elective laparoscopic Cholecystectomy in healthy patients.

Objectives

General Objective

To compare between Spinal Anaesthesia and General Anaesthesia for laparoscopic cholecystectomy in healthy patients in a tertiary hospital, Bangladesh

Specific Objective

To know more about the risk and benefit of laparoscopic cholecystectomy procedure in Bangladesh

MATERIALS AND METHODS

A comparative clinical study was conducted in the department of Anesthesia, Pabna Medical College Hospital, Pabna, Bangladesh during the period from January 2018 to April 2019. The aim of our study was to compare between Spinal Anaesthesia and General Anaesthesia for laparoscopic cholecystectomy in healthy patients. A total of 80 patients aged between 18-65 years of both sex with ASA Grade status I and II undergoing elective laparoscopic Cholecystectomy were selected for the study. After taking written informed consent from the study participants, we were elected 80 study subjects and divided into two groups; Group I (n=40) received General Anaesthesia. And Group II (n=40) received Spinal Anaesthesia. After taking approval, elective laparoscopic Cholecystectomy procedure was taken. After taking the patients to the operation theater, an intravenous line was secured in the right upper limb and infusion of 500 ml Hartmann's solution started. The initial pulse, blood pressure (BP), respiratory rate.

Oxygen saturation (SPO₂) were noted. All the patients were premeditated with Inj. Nalbuphine HCL 0.2mg per kg, Midazolam 0.02 mg/kg and Inj. Ondansetron 0.08 mg/kg intravenously (i.v.). In patients randomized for Spinal Anesthesia, the patient was first made to lie in supine position and all the monitors were attached. Oxygen was then administered through Face mask at 3 l/minute. Then the patient was made to lie in right lateral decubitus position. A 25 G Quincke spinal needle was introduced in subarachnoid space at L1-L2 interspace under all aseptic and antiseptic precautions. After confirming free flow of cerebrospinal fluid, 0.3 mg/kg of hyperbaric Bupivacaine 0.5% was injected intrathecally in cephalad direction. Then, the patients is

kept in 1^o Trendelenburg position for 5 minutes. Approximately 10 minutes after intrathecal injection, the level of analgesia was checked. A segmental sensory (pin-prick) block, extending between T4 and L5 dermatomes, was obtained without any respiratory distress. Laparoscopic cholecystectomy was performed using the same techniques in both the groups with standard for trocar insertion. Pneumoperitoneum was established by using the open (Hasson) technique with carbon dioxide at maximum intra-abdominal pressure of 12 mm Hg. Intraoperatively, the patients randomly allocated to general anesthesia group received fentanyl citrate 2 µg/kg i.v. as an adjuvant while those allocated to spinal anesthesia group were given 25 µg i.v. as bolus and when required. All the patients were monitored continuously both for clinical observation and noninvasive hemodynamic monitoring like pulse, blood pressure, respiratory rate, pulse oximetry which were recorded at 5 minute interval. Operative times as well as any intraoperative events such as shoulder pain, headache, nausea, and discomfort were recorded. Postoperative pain was assessed at 4, 8, 12 and 24 hours by using the Visual Analogue Scale (VAS) after completion of procedure. Other postoperative events either related to surgical or especially to anesthetic procedure, such as discomfort, nausea and vomiting, shoulder pain, urinary retention, pruritus, headache and other neurological sequel, were recorded.

RESULTS

All the procedures were completed within the allocated method of anesthesia. Intra-operatively, there was no bradycardia in either group. In Group-II, hypotension (i.e. >30% fall in BP) was noted in 12 (30%) cases, out of which Ephedrine HCL 2ml was given in 26 cases and the rest were managed with i.v. fluids, while in Group-I, hypotension was not noted and all of them were managed with i.v. fluids. Pain/discomfort in right shoulder was noted in 9 (22.5%) cases but it was severe enough in only 4 (10%) cases which received i.v. fentanyl 25 µg bolus once. Rests were managed with massage over right shoulder. The remaining patients did not require any additional medication or other intervention, and procedures were completed uneventfully in all cases. Intraoperative comparison of mean pulse rate in Group-I and Group-II showed less tachycardia. Figure shows that mean systolic and diastolic pressure, respectively, in both the groups, which were found to be higher in Group-I compared to Group-II. This study shows that there was no pain or respiratory distress in Group-II. Mean discharge from the hospital in Group-I was 48.33 hours and in Group-II it was 36.53 hours. There was no mortality or morbidity in either group. Regarding the postoperative complications, nausea was present in 12 (30%) cases in Group-I while none had it in Group-II. Dizziness was there in 8 (20%) cases in Group-I while none had it in Group-II. Pruritus was there in 5 (12.5%) cases in Group-I and 3 (7.5%) cases in Group-II. Pain at local site was noted in 26 (65%) cases in Group-I and in

only 5 (12.5%) cases in Group-II. There was no headache, backache, urinary retention or any other major complication. At the time of discharge, all patients were asked about the satisfaction regarding the general as well as spinal anesthesia and the patients were more satisfied with Spinal Anesthesia than General Anesthesia. In the Table II we have shown Mean Pulse Rate, Mean Systolic and Diastolic BP readings of several periods of treatment duration at a glance. From that table we found after in sufflation

pulse rate of patients of group I jumped up to 114 which did not so in patients of group II. After starting treatment the systolic BP raised suddenly up to 135 in group I and stayed for a long time which did not so in group II. About diastolic BP in case of both the anaesthesia in both groups the DBP found steady but in group II it was some lower. Considering those it can be claimed that Spinal Anaesthesia has some superiority over General Anaesthesia for Laparoscopic Cholecystectomy.

Table-1: Demographic characteristics of the study participants (n=80)

Characteristics of patients	Group- I	Group-II
Male	28	25
Female	12	15
Mean Age in Years	36.67	34.58
Mean Operative Time(minutes)	66.03	66.63
Average Hospital Stay(hours)	48.33	36.53




Fig-I: Perioperative comparison of mean pulse rate in Group-I and Group-II




Fig-II: Perioperative comparison of mean systolic blood pressure in Group-I and Group-II




Fig-IV: Perioperative comparison of mean diastolic blood pressure in Group-I and Group-II

Table-1: At a glance Mean Pulse Rate (MPR), Mean Systolic BP(MSBP), Mean Diastolic BP MDBP & EtCO2 (n=80)

Sub.	Subject	Pre-operative n	Before Insufflation n	After Insufflation n	Aft. 30 min.	Aft. 45 min.	Aft. 60 min.	Post Operative	4 Hrs	8 Hrs	12 Hrs	24 Hrs
MPR	Group 1	86	98	114	96	93	94	95	90	86	82	82
	Group-2	86	88	89	90	91	91	92	88	84	80	80
MSBP	Group 1	120	133	135	132	128	130	132	128	130	129	128
	Group 2	123	122	122	120	121	117	117	116	120	116	116
MDBP	Group 1	75	81	88	87	87	88	86	87	86	87	87
	Group-2	80	80	80	78	79	78	78	76	75	74	73
EtCO2	Group 1	35	32	37	32	32	31	35	36	36	37	36
	Group-2	34	32	38	33	32	33	32	33	31	32	32

DISCUSSION

The present study has not only confirmed the feasibility of safely performing laparoscopic cholecystectomy under Spinal Anesthesia as the sole anesthetic procedure but also shown superiority of Spinal Anesthesia in terms of better postoperative pain control as compared to general anesthesia. Pain assessed throughout any time in the postoperative period during the patients' hospital stay was significantly lesser in spinal Group-Is compared to General Anaesthesia group, which is due to residual analgesic effect of local anesthetic in subarachnoid space and decrease in discomfort due to avoidance of general anesthesia [2, 10]. Pain relief, an important component for rapid and smooth recovery, was seen in spinal anesthesia group. Intraoperatively, two things were noted - hypotension and pain/discomfort in right shoulder in the spinal group. Hypotension is due to sympathetic blockade and mechanical effect of pneumoperitoneum, while pain and discomfort over right shoulder can be attributed to diaphragmatic

irritation from pneumoperitoneum with carbon dioxide. Most of this was managed without drugs, i.e., reassurance to the patient, massage of the right shoulder, keeping the intra-abdominal pressure to 12 mm Hg, avoiding excessive tilting of table and thereby minimizing diaphragmatic irritation. The use of low pressure pneumoperitoneum was adequate, especially with spinal group, as Spinal Anesthesia causes high level of motor, sensory and sympathetic blockade and thereby good abdominal muscle relaxation as compared to General Anesthesia. In Group-I, the initial increase in pulse rate and BP after peritoneal insufflations are due to both mechanical and neurohumoral effects [11,12]. The return of pulse rate and BP to normal baseline was gradual. In Group-II, there was little variation in pulse and BP after peritoneal insufflation as spinal anesthesia tends to decrease the pulse and BP, while the neurohumoral and mechanical effects of pneumoperitoneum tend to increase them. After several minutes, the neurohumoral and mechanical effects are compensated so that there is slight decrease in the pulse rate and BP. The decrease in pulse rate and BP in

Group-II as compared to Group-I can be explained as due to decrease in pain caused by residual analgesic effect of local anesthetic in subarachnoid space. Nausea and vomiting are particularly troublesome after laparoscopic surgery; over 50% of patients required antiemetics, so prophylactic antiemetics had been given routinely. Regarding the postoperative complications, nausea, vomiting and dizziness were more common with general anesthesia due to intubation of trachea and intravenous drugs. As spinal anesthesia is a regional block, there is less procedure-related cost and hospital stay because of less postoperative pain and complications.

Limitations of the study

This was a clinical study in a single centre with small a sample size. So, the study results may not reflect the scenarios of the whole community.

CONCLUSION AND RECOMMENDATIONS

In our study, we observed that in comparison to General Anaesthesia, Spinal Anaesthesia provides better safety and adequacy in healthy patients and hence provides better post-operative pain control without limiting the recovery. Post-operative complications like nausea, vomiting, dizziness and pneumonia are less in Spinal Anaesthesia. Authors are recommending conducting multi-centre study with a large number of sample sizes.

REFERENCES

1. Hamad MA, Ibrahim El-Khattary OA. Laparoscopic cholecystectomy under spinal anesthesia with nitrous oxide pneumoperitoneum: a feasibility study. *Surg Endosc.* 2003; 17(9):1426-1428.
2. Tzovaras G, Fafoulakis F, Pratsas K, Georgopoulou S, Stamatiou G, Hatzitheofilou C. Spinal vs general anesthesia for laparoscopic cholecystectomy: interim analysis of a controlled randomized trial. *Archives of Surgery.* 2008 May 1;143(5):497-501.
3. Soper NJ, Barteau JA, Clayman RV, Ashley SW, Dunnegan DL. Comparison of early postoperative results for laparoscopic versus standard open cholecystectomy. *Surgery, gynecology & obstetrics.* 1992 Feb;174(2):114-8.
4. Wickham JE. Minimally invasive surgery: future developments. *BMJ.* 1994 Jan 15;308(6922):193-5.
5. Johnson A. Laparoscopic surgery. *Lancet.* 1997; 349(9052):631-635.
6. Crabtree JH, Fishman A, Huen IT. Videolaparoscopic peritoneal dialysis catheter implants and rescue procedures under local anesthesia with nitrous oxide pneumoperitoneum. *Advances in Peritoneal Dialysis.* 1998;14:83-6.
7. Sharp JR, Pierson WP, Brady CE. Comparison of CO₂ and N₂O-induced discomfort during peritoneoscopy under local anesthesia. *Gastroenterology.* 1982; 82(3):453-456.
8. Rodgers A, Walker N, Schug S, McKee A, Kehlet H, Van Zundert A, Sage D, Futter M, Saville G, Clark T, MacMahon S. Reduction of postoperative mortality and morbidity with epidural or spinal anaesthesia: results from overview of randomised trials. *Bmj.* 2000 Dec 16;321(7275):1493.
9. Lennox PH, Chilvers C, Vaghadia H. Selective spinal anesthesia versus desflurane anesthesia in short duration outpatient gynecological laparoscopy: a pharmacoeconomic comparison. *Anesthesia & Analgesia.* 2002 Mar 1;94(3):565-8.
10. Chilvers CR, Goodwin A, Vaghadia H, Mitchell GW. Selective spinal anesthesia for outpatient laparoscopy. V: Pharmacoeconomic comparison vs general anesthesia. *Canadian Journal of Anesthesia.* 2001 Mar 1;48(3):279.
11. Aono H, Takeda A, Tarver SD, Goto H. Stress responses in three different anesthetic techniques for carbon dioxide laparoscopic cholecystectomy. *Journal of clinical anesthesia.* 1998 Nov 1;10(7):546-50.
12. Wahba RW, Beique F, Kleiman SJ. Cardiopulmonary function and laparoscopic cholecystectomy. *Can J Anaesth.* 1995; 42:51-63