

Original Research Article

Herpes Simplex Virus Genotyping in Neurological abnormalities-Clinical Relevance for Disease Monitoring

Rekha Gorkela¹, Sachin Kumar², Anchal Chaurasia³, Binish Jawed⁴, Mukta Sehgal⁴, Meenakshi Bahuguna¹, Vatsal Prashar², Indu Bushan⁵, Manish Dev Sharma¹, Narotam Sharma^{1*}

¹Central Molecular Research Laboratory, Department of Biochemistry, SGRRIM&HS, Patel Nagar, Dehradun-UK & SGRRTS, Patel Nagar, Dehradun-UK, India

²Gurukul Kangari Vishwavidyalaya, Haridwar, Uttarakhand, India

³Kanya Gurukul Campus, Gurukul Kangri, University, Haridwar, Uttarakhand, India

⁴Biotechnology Department, Maharaj Vinayak Global University, Jaipur, Rajasthan, India,

⁵PDM Dental College, Bahadurgarh, Haryana, India

*Corresponding author

Dr. Narotam Sharma

Email: sharmanaroam5@gmail.com

Abstract: Herpes Simplex Virus (HSV) is an important etiological agent for Neurological disorders like meningitis, seizures etc. Current work includes the molecular revealing of HSV 1 and 2 in CSF specimens. 13 cases of CSF were processed for the typing of HSV by Real Time PCR and out of which 03 cases came positive. All the positive cases were HSV genotype-1. Advanced biotechnological tools promises fast, accurate and timely diagnosis of such viral diseases for further prognosis.

Keywords: Neurological abnormalities, Real Time PCR, Herpes Simplex viruses, Genotypes, Viral Meningitis.

INTRODUCTION:

Herpes simplex viruses (HSV) are categorized into two types, HSV-1, or oral herpes and HSV-2, or genital herpes. HSV-1 is mainly transmitted by oral to oral contact, whereas HSV-2 is a sexually transmitted infection (STI) that can cause painful genital ulcer [1]. HSV-1 is a highly contagious infection, which is common and endemic throughout the world. The virus causes lifelong infection, and there is no cure, although treatment can reduce symptoms. After the first (primary) infection, HSV, like other herpes viruses, remains dormant (latent) in the body and can periodically reactivate and cause symptoms [2, 3]. HSV infection causes recurring episodes of small, painful, fluid-filled blisters on the skin, mouth, lips (cold sores), eyes, or genitals [4, 5]. In the current study, characterization of HSV genotypes 1 and 2 in cerebrospinal fluid (CSF) in the patients with Viral Meningitis and other Neurological Disorders cases were studied.

MATERIALS AND METHODS:

Cerebrospinal fluid was collected from various Departments of Shri Mahant Indires Hospital, Patel Nagar, Dehradun, and Uttarakhand. 13 cerebrospinal fluids (CSF) were collected from the patients with Neurological disorders which mainly include seizures, epilepsy, meningitis, hydrocephalus, altered sensorium etc. Further the CSF was preprocessed for DNA isolation and DNA was extracted by silica column method. The amplification of glycoprotein gene in HSV was done by Qiagen Rotor-gene Q Real Time PCR. Pre PCR mix was prepared as per the manufacturer's protocol (artus HSV-1/2 PCR Kit CE-QIAGEN). All the 13 cases were processed for the molecular genotyping of HSV-1 and HSV-2 along with exogenous internal control which is used to check for any artifacts in processing and in PCR protocol. Direct detection of the fluorescence can be detected as specific amplification plots which can be visualized in channel cycling Green (470nm) and cycling Orange (detector 610nm) and Internal Control fluorescence channel cycling Yellow (530nm- 555nm) for HSV-1, HSV-2 and internal control respectively (Figure 1).




Fig 1: Amplification plots for HSV-1 and 2 in Real time PCR machine.

RESULTS:

Out of 13 CSF specimens processed, 03 cases came positive with the occurrence of HSV type 1 in all the positive cases. As the HSV-1 is prevalent worldwide it was also studied w.r.t to gender and age wise. 10 males and 03 females were considered for the study and HSV-1 positive cases were mostly seen in males. All the positive cases were with meningitis & Neurological disorders.

DISCUSSION AND CONCLUSION

HSV-1 is mainly transmitted by oral contact whereas HSV-2 causes sexually transmitted infections. Viral meningitis can cause inflammation of the brain (encephalitis). HSV meningitis is a serious infection but diagnosis previously required brain biopsy in certain cases due to low sensitivity of CSF culture and serology [6, 7]. PCR allows the detection of HSV DNA from CSF with 95% sensitivity thus avoiding invasive brain biopsy. Viral meningitis, commonly caused by either enteroviruses or HSV is more reliably detected by PCR when compared to culture and in a shorter time (one verses up to 5 days). HSV 1 classically presents as herpes gingivostomatitis an infection of the oral mucosa. It can also cause conjunctivitis, keratitis, and herpetic whitlow. HSV 2 is most common cause of genital ulcer. It has been documented that as many as one third of herpes infections are due to HSV 1, particularly in adolescent and young adult [8]. The proper management of such HSV infections required customized therapy with specific regimens. Advanced Molecular tools for specific typing of HSV have now enabled the clinicians to treat the disease effectively.

REFERENCES:

1. Jain KK. Molecular Diagnostics for Neurological Disorders. In Applications of Biotechnology in Neurology 2013 (pp. 155-210). Humana Press.
2. Itzhaki RF, Lin WR, Shang D, Wilcock GK, Faragher B, Jamieson GA. Herpes simplex virus

type 1 in brain and risk of Alzheimer's disease. The Lancet. 1997 Jan 25; 349(9047):241-4.

3. Kennedy PG. Potential use of herpes simplex virus (HSV) vectors for gene therapy of neurological disorders. Brain. 1997 Jul 1; 120(7):1245-59.
4. Whitley R. Neonatal herpes simplex virus infection. Current opinion in infectious diseases. 2004 Jun 1; 17(3):243-6.
5. Tyler KL. Herpes simplex virus infections of the central nervous system: encephalitis and meningitis, including Mollaret's. HERPES-CAMBRIDGE-. 2004 Jun; 11:57A-64A.
6. Whitley RJ, Lakeman F. Herpes simplex virus infections of the central nervous system: therapeutic and diagnostic considerations. Clinical Infectious Diseases. 1995 Feb 1; 20(2):414-20.
7. Kimberlin D. Herpes simplex virus, meningitis and encephalitis in neonates. Herpes: the journal of the IHMF. 2004 Jun; 11:65A-76A.
8. Langenberg AG, Corey L, Ashley RL, Leong WP, Straus SE. A prospective study of new infections with herpes simplex virus type 1 and type 2. New England Journal of Medicine. 1999 Nov 4; 341(19):1432-8.