

## **Congenital Diaphragmatic Hernia in an Adult: A Case Report**

**Pradhan S, Pattanaik MR, Mohanty T, Dash MR, Mallick A**

S. C. B. Medical College & Hospital, Cuttack, Odisha, Pin: 753007

### **\*Corresponding author**

Pradhan S

Email: [spswetapadmapradhan@gmail.com](mailto:spswetapadmapradhan@gmail.com)

---

**Abstract:** The incidence of congenital diaphragmatic hernia varies from 1: 2000 to 1: 4000 live births. However, the clinical manifestation of symptoms and its diagnosis in adults is very rare. Very few cases of adult patients with Bochdalek hernia have been reported. Here we report a case of an 18 year Muslim girl with eight months history of shortness of breath, left sided chest pain, heart burn and hiccup. After her medical work up she was diagnosed as a case of Bochdalek hernia. The Bochdalek hernia is commonly misdiagnosed in adult. Unlike infants who present with respiratory distress early, the most frequent symptom in an adult is mild discomfort and 25% of adult patients are asymptomatic. Our patient was misdiagnosed earlier and was treated with anti tubercular therapy for six months until she was admitted to our hospital. Congenital diaphragmatic hernia in adults is rarely suspected in the differential diagnosis of left sided chest pain with breathlessness. Correct diagnosis and early treatment is important to avoid the occurrence of serious complication.

**Keywords:** Congenital diaphragmatic hernia, Bochdalek hernia, diaphragm

---

### **INTRODUCTION**


Congenital diaphragmatic hernia (CDH) is characterized by displacement of abdominal organs into the thorax through a defect in diaphragm [1]. They are classified as Bochdalek, Morgagni and hiatus hernias. They generally present in the first few hours of life. On the other hand, traumatic hernias can present in any age group. Diaphragmatic hernias seen in adults are commonly due to trauma [2].

Congenital diaphragmatic hernias presenting late in adolescence and adult life is a very rare entity [3]. Amongst them, Bochdalek's hernia is a posterolateral congenital defect of the diaphragm, localized on left side in 70-90% of the cases [4]. It was first described in 1848 by the Czechoslovakian anatomist, Vincent Alexander Bochdalek that carries his name till today. This condition is rare in adults, and there are 130 reported cases in the medical literature [4].


### **CASE REPORT**

An 18 year old Muslim girl presented with eight months history of intermittent breathlessness, left sided chest pain aggravated after taking food. Chest pain was dull aching, diffuse all over the left hemithorax with no radiation to any other site. It was associated with frequent episodes of heart burn and hiccup. There was no history of thoracic or abdominal trauma. She was diagnosed as a case of pulmonary tuberculosis radiologically 6 months back and given ATT (antitubercular treatment) by a general practitioner. There was no radiological or clinical

improvement even after completion of 6 months of ATT. On physical examination, her blood pressure was 110 / 70 mm of Hg with a heart rate of 90 beats per minute and respiratory rate 16 per minute respectively. Her body temperature was 98.0F and SpO2 98% in room air. There was no pallor, icterus, cyanosis, clubbing, lymphadenopathy or oedema. Respiratory system examination show reduced chest expansion and breath sounds on left side. Bowel sounds were audible on left side. Chest x-ray posterior anterior view showed a large cavity with multiple air fluid levels and a normal right lung (fig - 1). Chest x-ray posterior anterior view after insertion of feeding tube shows the position of stomach (fig - 2). Barium follow through test show herniation of stomach and small bowel through left hemithorax with mesentrioaxial volvulus of stomach. There was also delayed emptying of contrast (fig 3a, 3b). Computed tomography was performed which confirmed a left sided Bochdalek hernia with herniation of stomach, small and large bowels, spleen and abdominal blood vessels into left hemithorax and shifting of mediastinum to contralateral side with compression of Left lung (fig 4a, 4b, 4c, 4d). Then the patient was referred to a cardiothoracic surgeon for surgical correction.



**Fig 1:** Chest x-ray (PA view) showing large cavity with multiple air fluid level in left mid and lower zone.




**Fig 2:** Chest X-ray with Ryle's tube shows the position of stomach




**Fig 3a:** 5 minutes after barium follow through




**Fig 3b:** 15 minutes after barium follow through




**Fig 4a:** CT scan: Lung window shows stomach containing air and food material and large intestine with haustration.



**Fig 4b:** CT scan showing presence of stomach & large intestine in thorax.



**Fig 4c: CT scan showing presence of stomach, spleen, gut in thorax**



**Fig 4d: CT scan showing presence of mesentery, large gut, spleen, abdominal blood vessels in thorax**

## DISCUSSION

The diaphragm separates thoracic and abdominal cavity. The herniation of abdominal contents into thoracic cavity through a defect in the diaphragm is called diaphragmatic hernia. The diaphragmatic defect may be congenital or acquired. The most frequent cause of diaphragmatic hernia in adult seems to be trauma (blunt or penetrating) followed by iatrogenic (oesophagus-gastric surgery for oesophagus cancer or gastric cancer), whereas in babies or newborns it is most often attributable to congenital absence or defective fusion of the septum transversum or the pleuroperitoneal membrane [5].

The diaphragm is developed in the embryo during the third week of gestation and is composed of four components: transverse septum, two pleuroperitoneal folds, cervical myotomes and dorsal mesentery. Its development is completed between the 8th and 10th week of gestation. The congenital defect that determines congenital diaphragmatic hernias (CDH) is lack of development of the pleuroperitoneal folds or the inappropriate migration or absence of the diaphragmatic musculature or weakness in the embryological points of fusion. The most frequent types of diaphragmatic hernia are the left posterolateral (Bochdalek hernia) and the sternocostal (Morgagni hernia) types.

A Bochdalek hernia is caused by a lack of closing of the pleura peritoneal cavity by incomplete diaphragmatic development before the intestine returns to the abdomen from the yolk sac between weeks 8 and 10 of gestation. The majority present during neonatal life. Being associated with congenital pulmonary abnormalities they have a poor prognosis [6, 7]. The major clinical problem is pulmonary hypoplasia, a result of the lung having failed to develop in utero as the thoracic cavity is filled by abdominal contents [7]. If hernia formation precedes lung development, pulmonary hypoplasia may occur with severe respiratory compromise at birth. But in adults, this defect is uncommon, the lung in most cases develops normally and therefore symptoms are rare [8]. Bochdalek hernia in adult can present in two ways. Either an incidental finding during x-ray examination performed for symptoms not related to hernia [9, 10] or when symptoms develop as a result of incarceration, strangulation or rupture of viscera. Symptoms may be due to digestive system involvement, which include abdominal pain, nausea, vomiting, constipation or due to respiratory system involvement that presents with chest pain, breathlessness, wheezing [9,11]. In our case the patient complained of intermittent breathlessness and chest pain. The left-sided BH occurs in approximately 70 - 90% of cases. Left-sided hernias allow herniation of both the small and large bowel and intra-abdominal solid organs into the thoracic cavity. The most frequently displaced organ is the stomach followed by the colon, spleen, small intestine and ureter [12]. Gastric volvulus is one of the rare but recognized complications of Bochdalek hernia [13]. In our case there was mesentrio-axial volvulus of stomach.

Direct chest and abdominal X-rays, barium enema examinations, ultrasonography, computerized tomography, magnetic resonances, laparoscopy and laparotomy can be used for diagnosis. The radiological features on X-ray films are not always easy to detect. Typically, radiological images show intra-thoracic gas-filled loops of the bowel with a contralateral shift of the mediastinum mimicking pleura-pulmonary pathology like pneumothorax, consolidation [14, 15]. A computed tomography (CT) scan is the gold standard radiological investigation that allows the highest accuracy for a correct diagnosis, as in our case. Treatment of Bochdalek hernia is always surgical depending on the presence of visceral complication. In an elective setting most authors recommend the thoracic approach; on the other hand, when there are septic complications, the abdominal approach is preferred. The current trend is to use minimal invasive surgical techniques such as laparoscopy, and specially thoracoscopy, which has been satisfactorily performed in adults [16, 17, 18].

## CONCLUSION

The occurrence of CDH (Bochdalek hernia) in adults is rare and misleading even to experienced

clinicians, as the patients present with symptoms mimicking other diseases. Hence high index of clinical suspicion is required for prompt diagnosis and treatment in order to avoid complications such as strangulation or bowel perforation.

## REFERENCES

1. Mullins ME, Stein J, Saini SS, Mueller PR; Prevalence of incidental Bochdalek's hernia in a large adult population. *AJR Am J Roentgenol.* 2001; 177(2):363-366.
2. Snyder WH Jr., Greaney EM Jr; Congenital Diaphragmatic Hernia; 77 Consecutive Cases. *Surgery.* 1965; 57: 576-588.
3. Anantha ramakrishnan R, Senthil KK, Karunanithi R; Congenital Diaphragmatic Hernia (Bochdalek Hernia) in an Adult. *Chettinad Health City Medical Journal.* (Internet) [Last accessed on 14 January 2015]. Available from [www.chcmj.ac.in/journal/pdf/vol1\\_no2/congenitaldiaphragmatic.pdf](http://www.chcmj.ac.in/journal/pdf/vol1_no2/congenitaldiaphragmatic.pdf)
4. Perch P, Houck W, de Anda A; Symptomatic Bochdalek hernia in an octogenarian. *Ann Thorac Surg* 2002; 73:1288-9.
5. Bianchi E, Mancini P, Vito SD, Pompili E, Taurone S, Guerrisi I, *et al.*; Congenital asymptomatic diaphragmatic hernias in adults: a case series. *Journal of Medical Case Reports.* 2013; 7:125. (Internet) [Last accessed on 13 January 2015]. Available from <http://www.jmedicalcasereports.com/content/7/1/125>
6. Gedik E, Tuncer MC, Onat S, Avcı A, Tacyıldız I, Bac B; A review of Morgagni and Bochdalek hernias in adults. *Folia Morphol* 2011; 70(1):5-12. (Internet)[Last accessed on 13 January 2015]. Available from [http:// www.czasopisma.viamedica.pl/fm/article/download/19322/15215](http://www.czasopisma.viamedica.pl/fm/article/download/19322/15215)
7. Horton DB, Lieberman G; An Unusual Presentation of Diaphragmatic Hernia. (Internet) [Last accessed on January 2015]. Available from [www.bidmc.harvard.edu/LearningLab/gastro/Horton.pdf](http://www.bidmc.harvard.edu/LearningLab/gastro/Horton.pdf)
8. Miller BJ, Martin IJ; Bochdalek hernia with hemorrhage in an adult. *Can J Surg* 1993; 36(5):476-8.
9. Betremieux P, Dabadie A, Chapuis M, Pladys P, Treguier C, Fremond B *et al.*; Late presenting Bochdalek hernia containing colon: misdiagnosis risk. *Burg J Pediatr Surg* 1995; 5(2):113-5.
10. Thomas S, Kapur B; Adult Bochdalek hernia: clinical features, management and results of treatment. *Jpn J Surg* 1993; 22(1):114-9.
11. Shin MS, Mulligan SA, Baxley WA, Ho K-J; Bochdalek hernia of diaphragm in the adult. *Chest* 1987; 92:1098.
12. Vega TM, Maldonado HR, Vega TG, Vega TA, Liévano AE, Velázquez MP; Late-onset congenital diaphragmatic hernia: A case report. *International Journal of Surgery Case Reports* 2013; 4:952-54. (Internet)[Last accessed on 13 January 2015]. Available from <http://www.sciencedirect.com/science/article/pii/S2210261213002605>
13. Giannoulis K, Sutton R; Bochdalek hernia presenting in adult life: report of an unusual case and review of the literature. *Annals of Gastroenterology* 2004; 17(1):109-12.
14. (Internet)[Last accessed on January 2015]. Available from [www.annalsgastro.gr/index.php/annalsgastro/article/download/275/243](http://www.annalsgastro.gr/index.php/annalsgastro/article/download/275/243)
15. Lin ST, Moss DM, Henderson SO; A case of Morgagni hernia presenting as pneumonia. *J Em Med* 1997; 15: 297-301.
16. Gayer G, Bilik R, Vardi A; CT diagnosis of delayed presentation of congenital diaphragmatic hernia simulating massive pleura pneumonia. *Eur Radiol* 1999; 9:1672-1674.
17. Taskin M, Zengin K, Unal E, Eren D, Korman U; Laparoscopic repair of congenital diaphragmatic hernias. *Surg Endosc* 2002; 16:869.
18. Willemsse P, Schutte PR, Plaisier PW; Thoracoscopic repair of a Bochdalek hernia in an adult. *Surg Endosc* 2003; 17:162.
19. Silen ML, Canvasser DA, Kurkchubasche AG, Andrus CU, Naunbeim KS; Video-assisted thoracic surgical repair of a foramen Bochdalek hernia. *Ann Thorac Surg* 1995; 60(2):448-50.