

Uncommon Benign Disease of the Salivary Glands Necrotizing Sialometaplasia (NS): Review of the Literature and Serie of 2 Cases

N. Belhaj^{1*}, R. Bencheikh¹, M. A. Benbouzid¹, L. H. Essakalli²

¹Department of Otorharyngology Head and Neck Surgery CHU Ibn Sina Rabat, Morocco

²Departement of Otorhinolaryngology Head and Neck Surgery CHR Guelmim, Morocco

DOI: [10.36347/sjmcr.2024.v12i05.102](https://doi.org/10.36347/sjmcr.2024.v12i05.102)

Received: 24.04.2024 | Accepted: 27.05.2024 | Published: 31.05.2024

*Corresponding author: N. Belhaj

Department of Otorharyngology Head and Neck Surgery CHU Ibn Sina Rabat, Morocco

Abstract

Case Report

Necrotizing sialometaplasia (NS) is an uncommon self limiting commonly confused with malignant epithelial entities; the aim of this article is to report series of two cases of necrotizing sialometaplasia: the first case is in 45 year old women with clinical documentary; the second case is an 37 year old man. Data from an extensive review of literature including clinical; imaginologic and microscopic features are provided. Information on diagnostic and prognostic factors is offered and comprehensibly discussed.

Keywords: necrotizing sialometaplasia; salivary gland lesion; oral squamous cell carcinoma.

Copyright © 2024 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

INTRODUCTION

Necrotizing sialometaplasia is an uncommon self limiting confused with malignant entities; the first was described by Dun lap and braker. Afterwards Branon et al using data for theirs own cases and review of the literature; described extensively its clinicopathological features [1-3]. During the following years this entity has been found in different locations [4-8]. And since Abrams *et al.*, first description; the associated etiologic factors of NS have been extensively discussed [2-4, 8]. The most accept theory suggested ischemia as the main etiologic factor. Age range in NS patient varies from (1.5-83 years old) man are more commently affected with male-female ratio 1.95; size from 0.5to 4 cm.

SERIES OF CASES

Observation 1

A 43 year old man ;he denied alcohol-smoking habits; he was first viewed by his general physician and he told him that probably that was malignant lesion; because it is an ulcerated lesion and rapidly growing .The patient was viewed two day ago and the daytime clinical examination finds an ulcerates lesion with fibrinous deposits; the diametr's lesion was 2cm in large ;the

patient report that the lesion appeared pearing during the previous day; and reported that non traumatic episode had occurred recently in the involved area (Figure 1).

Panoramic and occlusals radio-graphs were taken and review confirmed that non bone involvement was present behind the ulcers.

Under regional anesthesia; an incisional biopsy measuring 1.1cm /1.1 cm was taken from the right ulcer and immersing in formalin for sent to pathological laboratry.

The patient was decharged by analgesic and antibiotics covered. The presence of hyperplasic stratified squamous epithelium with long rete ridges and pseudoepitheliomatous hyperplasia (Figure 2); the bulk of lesion was composed of metaplastic squamous epithelial cells deriving from ductual epithelium; no noeplasic changes were observed; no tuberculosis lesions were observed.

TPHA-VDRL and HIV serology were negative. Two week later the lesion was discovered with fibronoid pseudo-membranes; one month late the ulcers decreased in size to approximatively hal of the origin size.



Figure 1: Ulcerated lesion with fibrinous deposits




Figure 2: HE.40 metaplastic squamous epithelial cells deriving from ductal epithelium

Observation 2

45 year old women; she consulted his dentist for painful lesion located in the oral palate that bleeding upon food contact; the patient use oral-washes and buccal miconazol. She denied alcohol-smoking habits. This lesion appear 2months.

Non-improvement and height growth; the patient was addressed in our formation.

The clinical examination found an ulcero-budding lesion 3cm in the large diameter (Figure 3).

Under an local anesthesia; a biopsy was realised; after the patient was decharged by analgesic and antibiotiques covered. The result of the histopathological stud show the same aspect of the first case (Figure 4 & 5).



Figure 3: Ulcero-budding lesion 3 cm in the large diameter




Figure 4 & 5: Microscopic sections of biopsy showing necrosis within salivary glands

DISCUSSION

NSM is an uncommon self-limiting lesion commonly confused with malignant epithelial neoplasms. Despite having been recognized by Abrams *et al.*, in 1973 and with numerous intraoral and extra oral cases reported, 1-40 no agreement on the etiology for NSM development has been reached to date. Most authors sustain that some kind of physical-chemical or biological aggressions over the local blood vessels will produce ischemic changes, leading to infarction with posterior necrosis, and that NSM is the ductal proliferation in response to injury [2, 8, 20, 21].

Many causes have been suggested as possible etiologic factors including: traumatic, chemical, surgical, and infectious events including alcohol, smoking, and drug abuse [3, 10, 12-16, 22-24].

In other instances, some diseases characterized by vascular disturbances have been associated with NSM, supporting the theory that ischemic necrosis is the main etiologic factor [17-19].

In 1982, Anneroth and Hansen²⁶ analyzed its histopathogenesis and separated NSM into five microscopic stages: infarction, sequestration, ulceration, reparative stage, and healing.

This entity is more frequently found in palate, although it has been also described in other intraoral locations: retro-molar area, labial glands, tongue, floor of the mouth, buccal mucosa, mucobuccal fold, sublingual region, and it was also found in parotid and sub-mandibular glands [3, 4, 8-10].

Also, NSM cases were reported in the head and neck area: nasal cavity, maxillary sinus, tonsillar fossa, incisive canal, larynx, trachea, and lymph nodes [3, 5, 6, 9, 15, 27, 31].

The most commonly observed clinical feature is a palatal, painful or painless, deep seated ulcer with clean, sharp, and sometimes erythematous or rolled margins, measuring from 1 cm to 3.5 cm. Other symptoms are somewhat variable: fever, chills, malaise, anesthesia, and paresthesia, and rarely NSM is found as

a nonulcerative, purulent, diffuse lesion with palpable lymph nodes [3, 10, 12-14, 32].

Our case is unique since one of the lesions measured almost 6 cm. The presence of a unique lesion is the most frequent clinical manifestation of NSM, but as it is in the case presented here, sometimes two lesions are present [3, 14].

Another three rare and unique cases have been reported: one with three ulcers, one with multiple palatal ulcerated lesions, and one with metachronous lesions [11, 13, 32]. Also, several cases with one or two palatal asymptomatic, painless, smooth-surfaced nodules of normal color were reported [8, 11, 14, 18, 23]. Duration of the lesion varies from 3 weeks to 3 months [33].

It is well known that almost all studied NSM cases did not display radiographic alterations, but saucerization of the subjacent bone was observed in four reported cases [1, 3, 11].

Some studies of the features of NSM using magnetic resonance (MR) analysis [34-36] showed this lesion was hyperintense on T2 and hypointense on T1 [34]. Lee *et al.*, [35] reported that under MRI scan, a NSM case associated with an adenoid cystic carcinoma presented a diffuse infiltrating image suggestive of a malignant lesion, and a multislice computed tomographic study showed no bone changes [36].

It is essential that the pathologist should receive an adequate biopsy specimen in order to observe all the diagnostic microscopic changes. Under microscopic examination, the lesion is usually covered by hyperplastic squamous stratified epithelium showing elongated epithelial ridges (pseudoeplitheliomatous hyperplasia). A very important finding is the acinar necrosis featured by disintegration of the mucus-secreting acinar cells with persistence of the acinar outlines accompanied by spaces filled with mucin, cellular debris, and inflammatory infiltrate of variable intensity composed by polymorphonuclear neutrophilic leukocytes, plasma cells, foamy macrophages, and numerous eosinophils.

The most prominent and important diagnostic feature is the presence of ductal squamous metaplasia. It is characterized by a squamous change in the cell lining of the gland ducts. These squamous cells proliferate, and this proliferating lining obliterates the lumen, transforming the ducts into solid masses or nidus of squamous stratified epithelial cells showing eosinophilic cytoplasm with uniform and bland nuclei. It is important to note that mitotic cells are scarce, and of normal appearance with no atypical features. These solid squamous epithelial masses are often observed close to or continuous with the enlarged rete pegs of the superficial epithelium. This finding is suggestive of malignancy and may lead to the common misdiagnosis of squamous cell carcinoma [25]. These epithelial nodular masses are present in variable sizes and number, show smooth outlines, and sometimes are complemented with areas of more or less extensive reactive fibrosis. Sometimes, microscopic features of NSM may lead to a diagnostic pitfall of chronic sialadenitis or degenerative age changes. Occasionally, lesions in poorly controlled diabetic patients with mucormycosis may mimic NSM and should not be confused with the so-called sialometaplasia [26, 30]. Microscopic differential diagnosis should be performed with subacute necrotizing sialadenitis, Sutton disease, syphilis, and tuberculosis [38].

The use of immunohistochemistry may help to distinguish mucoepidermoid carcinoma from NSM lesions [33].

Lesions with similar microscopic appearance have been reported in non-neoplastic breast parenchyma, bronchial mucosa, lung, and sweat glands [27-30] and there are several reports on malignant and benign tumors associated to NSM lesions [3, 6, 34, 36, 39, 40] indicating that NSM could develop as a secondary phenomenon.

Sufficient data on long-term follow-up of NSM lesions is not available to date. In this manuscript we present the clinical features observed during the medium-term follow-up in a patient.

CONCLUSION

Necrotizing sialometaplasia (NS) is a relatively uncommon benign disease of the salivary glands that most commonly occurs in the palate. It is often confused clinically and histopathologically with malignancies, such as squamous cell carcinoma or mucoepidermoid carcinoma; This condition requires histological evidence so as not to miss a cancer.

REFERENCES

1. Abrams, A. M., Melrose, R. J., & Howell, F. V. (1973). Necrotizing sialometaplasia. A disease simulating malignancy. *Cancer*, 32(1), 130-135.
2. Dunlap, C. L., & Barker, B. F. (1974). Necrotizing sialometaplasia: Report of five additional

cases. *Oral Surgery, Oral Medicine, Oral Pathology*, 37(5), 722-727.

3. Brannon, R. B., Fowler, C. B., & Hartman, K. S. (1991). Necrotizing sialometaplasia: a clinicopathologic study of sixty-nine cases and review of the literature. *Oral surgery, oral medicine, oral pathology*, 72(3), 317-325.
4. Ravn, T., Trolle, W., Kiss, K., & Balle, V. H. (2009). Adenosquamous carcinoma of the larynx associated with necrotizing sialometaplasia—A diagnostic challenge. *Auris Nasus Larynx*, 36(6), 721-724.
5. Poulson, T. C., Greer Jr, R. O., & Ryser, R. W. (1986). Necrotizing sialometaplasia obscuring an underlying malignancy: report of a case. *Journal of oral and maxillofacial surgery*, 44(7), 570-574.
6. Arguelles, M. T., Vilorio Jr, J. B., Talens, M. C., & McCrory, T. P. (1976). Necrotizing sialometaplasia. *Oral Surgery, Oral Medicine, Oral Pathology*, 42(1), 86-90.
7. Lima, M., Rocha, L. C., Siqueira, L., Carmo, L. C., & H Filho, J. (2002). Cystic form of necrotizing sialometaplasia in sublingual salivary gland. *Rev Bras Otorrinolaringol*, 68(2), 279-279.
8. Maisel, R. H., Johnston, W. H., Anderson, H. A., & Cantrell, R. W. (1977). Necrotizing sialometaplasia involving the nasal cavity. *The Laryngoscope*, 87(3), 429-434.
9. Randhawa, T., Varghese, I., Shameena, P. M., Sudha, S., & Nair, R. G. (2009). Necrotizing sialometaplasia of tongue. *Journal of Oral and Maxillofacial Pathology*, 13(1), 35-37.
10. Imbery, T. A., & Edwards, P. A. (1996). Necrotizing sialometaplasia: literature review and case reports. *The Journal of the American Dental Association*, 127(7), 1087-1092.
11. Santis, H. R., Kabani, S. P., Roderiques, A., & Driscoll, J. M. (1982). Necrotizing sialometaplasia: an early, nonulcerative presentation. *Oral Surgery, Oral Medicine, Oral Pathology*, 53(4), 387-390.
12. Speechley, J. A., Field, E. A., & Scott, J. (1988). Necrotizing sialometaplasia occurring during pregnancy: report of a case. *Journal of oral and maxillofacial surgery*, 46(8), 696-699.
13. Keogh, P. V., O'Regan, E., Toner, M., & Flint, S. (2004). Necrotizing sialometaplasia: an unusual bilateral presentation associated with antecedent anaesthesia and lack of response to intralesional steroids. Case report and review of the literature. *British dental journal*, 196(2), 79-81.
14. Romagosa, V., Bella, M. R., Truchero, C., & Moya, J. (1992). Necrotizing sialometaplasia (adenometaplasia) of the trachea. *Histopathology*, 21(3), 280-282.
15. Batsakis, J. G., & Manning, J. T. (1987). Necrotizing sialometaplasia of major salivary glands. *The Journal of Laryngology & Otolaryngology*, 101(9), 962-966.
16. Buller, D. L. (1980). Nodular and ulcerated lesions of the hard palate. *Journal of the American Dental Association (1939)*, 101(5), 823-824.

17. Lynch, D. P., Crago, C. A., & Martinez Jr, M. G. (1979). Necrotizing sialometaplasia: A review of the literature and report of two additional cases. *Oral Surgery, Oral Medicine, Oral Pathology*, 47(1), 63-69.
18. Mandel, L., Kaynar, A., & Dechiara, S. (1991). Necrotizing sialometaplasia in a patient with sickle-cell anemia. *Journal of oral and maxillofacial surgery*, 49(7), 757-759.
19. Johnston, W. H. (1977). Necrotizing sialometaplasia involving the mucous glands of the nasal cavity. *Human Pathology*, 8(5), 589-592.
20. Suckiel, J. M., Davis, W. H., Patakas, B. M., & Kaminishi, R. M. (1978). Early and late manifestations of necrotizing sialometaplasia. *Journal of oral surgery (American Dental Association: 1965)*, 36(11), 902-905.
21. Solomon, L. W., Merzianu, M., Sullivan, M., & Rigual, N. R. (2007). Necrotizing sialometaplasia associated with bulimia: case report and literature review. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology*, 103(2), e39-e42.
22. Prabhakaran, V. C., Flora, R. S., & Kendall, C. (2006). Pressure-induced necrotizing sialometaplasia of the parotid gland. *Histopathology*, 48(4), 464-465.
23. Fava, M., Cherubini, K., Yurgel, L., Salum, F., & Figueiredo, M. A. (2008). Necrotizing sialometaplasia of the palate in a cocaine-using patient. A case report. *Minerva Stomatologica*, 57(4), 199-202.
24. Hurt, M. A., Diaz-Arias, A. A., Rosenholtz, M. J., Havey, A. D., & Stephenson Jr, H. E. (1988). Posttraumatic lobular squamous metaplasia of breast. An unusual pseudocarcinomatous metaplasia resembling squamous (necrotizing) sialometaplasia of the salivary gland. *Modern Pathology: an Official Journal of the United States and Canadian Academy of Pathology, Inc*, 1(5), 385-390.
25. Hurt, M. A., Diaz-Arias, A. A., Rosenholtz, M. J., Havey, A. D., & Stephenson Jr, H. E. (1988). Posttraumatic lobular squamous metaplasia of breast. An unusual pseudocarcinomatous metaplasia resembling squamous (necrotizing) sialometaplasia of the salivary gland. *Modern Pathology: an Official Journal of the United States and Canadian Academy of Pathology, Inc*, 1(5), 385-390.
26. Anneroth, G., & Hansen, L. S. (1982). Necrotizing sialometaplasia: the relationship of its pathogenesis to its clinical characteristics. *International journal of oral surgery*, 11(5), 283-291.
27. Goldman, R. L., & Klein, H. Z. (1986). Proliferative sialometaplasia arising in an intraparotid lymph node. *American journal of clinical pathology*, 86(1), 116-119.
28. Hurt, M. A., Diaz-Arias, A. A., Rosenholtz, M. J., Havey, A. D., & Stephenson Jr, H. E. (1988). Posttraumatic lobular squamous metaplasia of breast. An unusual pseudocarcinomatous metaplasia resembling squamous (necrotizing) sialometaplasia of the salivary gland. *Modern Pathology: an Official Journal of the United States and Canadian Academy of Pathology, Inc*, 1(5), 385-390.
29. Pagni, F., Zàrate, A. F., & Urbanski, S. J. (2010). Necrotizing Sialometaplasia of Bronchial Mucosa. *International Journal of Surgical Pathology*, 18(1), 64-65.
30. Zschoch, H. (1992). Mucus gland infarct with squamous epithelial metaplasia in the lung. A rare site of so-called necrotizing sialometaplasia. *Der Pathologe*, 13(1), 45-48.
31. King, D. T., & Barr, R. J. (1979). Syringometaplasia: mucinous and squamous variants. *Journal of Cutaneous Pathology*, 6(4), 284-291.
32. Rossie, K. M., Allen, C. M., & Burns, R. A. (1986). Necrotizing sialometaplasia: a case with metachronous lesions. *Journal of oral and maxillofacial surgery*, 44(12), 1006-1008.
33. Rizkalla, H., & Toner, M. (2007). Necrotizing sialometaplasia versus invasive carcinoma of the head and neck: the use of myoepithelial markers and keratin subtypes as an adjunct to diagnosis. *Histopathology*, 51(2), 184-189.
34. Farina, D., Gavazzi, E., Avigo, C., Borghesi, A., & Maroldi, R. (2008). MRI findings of necrotizing sialometaplasia. *The British Journal of Radiology*, 81(966), e173-e175.
35. Lee, D. J., Ahn, H. K., Koh, E. S., Rho, Y. S., & Chu, H. R. (2009). Necrotizing sialometaplasia accompanied by adenoid cystic carcinoma on the soft palate. *Clinical and Experimental Otorhinolaryngology*, 2(1), 48-51.
36. Suomalainen, A., Törnwall, J., & Hagström, J. (2012). CT findings of necrotizing sialometaplasia. *Dentomaxillofacial Radiology*, 41(6), 529-532.
37. Gnepp, D. R. (1981). Warthin tumor exhibiting sebaceous differentiation and necrotizing sialometaplasia. *Virchows Archiv A*, 391, 267-273.
38. Femopase, F. L., Hernández, S. L., Gendelman, H., Criscuolo, M. I., & López de Blanc, S. A. (2004). Sialometaplasia necrotizante: presentación de cinco casos clínicos. *Medicina Oral, Patología Oral y Cirugía Bucal (Ed. impresa)*, 9(4), 309-320.
39. Wenig, B. M. (1995). Necrotizing sialometaplasia of the larynx: a report of two cases and a review of the literature. *American journal of clinical pathology*, 103(5), 609-613.
40. Franchi, A., Gallo, O., & Santucci, M. (1995). Pathologic Quiz Case 1: Necrotizing sialometaplasia obscuring recurrent well-differentiated squamous cell carcinoma of the maxillary sinus. *Archives of Otolaryngology-Head & Neck Surgery*, 121, 584-586.
41. Russo, A., Dell'Aquila, A., Prota, V., & Sica, G. S. (1998). Necrotizing sialometaplasia of the submandibular gland. Report of a case. *Minerva Stomatologica*, 47(6), 273-277.