

Research Progress on the Tolerance of Abscisic Acid in Plants to Abiotic Stresses

Li Luo¹, Xuefei Li¹, Jiayi Tan¹, Wenrui Liu¹, Fuquan Ye¹, Zhangyuan Hu¹, Jia Tian^{1*}¹College of Life Science and Technology, Heilongjiang Bayi Agricultural University, Daqing 163319, ChinaDOI: <https://doi.org/10.36347/sajb.2024.v12i07.007>

| Received: 15.07.2024 | Accepted: 28.08.2024 | Published: 31.08.2024

*Corresponding author: Jia Tian

College of Life Science and Technology, Heilongjiang Bayi Agricultural University, Daqing 163319, China

Abstract

Review Article

The growth and development of plants are inhibited by abiotic stresses, which is one of the factors limiting crop yield. Abscisic acid (ABA) is an important plant hormone that plays a crucial role in plants' response to abiotic stresses. In this review, abscisic acid was introduced from its discovery, distribution and transportation. The role of abscisic acid in abiotic stresses such as drought stress, salt stress and high temperature stress was discussed. Finally, the summary and prospect were summarized.

Keywords: Abscisic acid, Abiotic stresses, Phytohormone.

Copyright © 2024 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

1. INTRODUCTION

The growth and development of plants are influenced not only by biological factors, but also by various abiotic stresses. Abiotic stresses refer to environmental conditions that are unfavorable for plant growth, including high temperature, low temperature, water shortage, salt, and high light intensity [1]. Abiotic stresses affect plant growth, development, and productivity [2], alter plant morphology [3], and reduces photosynthesis [4]. It has the potential to cause a significant loss of approximately 51%-82% in the yield of major global food crops [5].

Abscisic acid (ABA) is one of the five major plant hormones [6], playing a crucial role in regulating various growth and development processes such as seed dormancy and germination [7]. It is also a key hormone in plant responses to abiotic stresses [8] and is referred as a stress hormone [9]. This review described abscisic acid and summarized its role in responses to drought stress, salt stress, and high temperature stress, thereby highlighting the critical importance of abscisic acid in enhancing plant resistance to abiotic stresses.

2. ABCISIC ACID

2.1 Discovery of abscisic acid

Abscisic acid refers to a class of metabolites known as isoprenoids (also termed terpenoids) [10]. In 1963, Philip Wareing isolated a growth-inhibiting substance from the buds of *Acer platanoides* and named it dormin [11]. In the same year, Frederick Addicott identified a substance that regulates the abscission of young cotton (*Gossypium hirsutum*) fruits and named it abscisin II [12]. Subsequently, it was revealed that the chemical structures of dormin and abscisin II were identical, indicating that they were the same compound, which was designated as abscisic acid [13,14].

2.2 Distribution of abscisic acid

Abscisic acid is widely distributed in all higher plants [15] and is extensively distributed throughout plant tissues and organs [16]. Plant organs such as leaves, roots, flowers, and fruits are capable of synthesizing abscisic acid [17]. In tissues and organs that are about to fall off or enter a dormant state [18], as well as in plants under stressful environmental conditions [19], the content of abscisic acid is relatively high. For instance, under drought stress, the content of ABA in plants increases to enhance their stress resistance [20,21].

2.3 Transportation of abscisic acid

Abscisic acid is mainly transported in a free form [22], capable of long-distance transport [23]. It can be transported over long distances through the phloem and xylem [24]. For example, ABA can undergo long-distance transport through the xylem and phloem in plant shoot and root [25]. Abscisic acid can also undergo short-distance transport, which predominantly takes place between cells, encompassing vascular cells and guard cells, among others [26]. ABA may undergo translocation from the biosynthesis sites to the guard cells [27], such as from vascular tissues to guard cells [28]. The translocation of ABA among cells, tissues, and organs plays a pivotal role in the plant's overall physiological response to stressful conditions [29].

3. ABCISIC ACID AND ABIOTIC STRESSES

3.1 Abscisic acid and drought stress

Drought is a recurrent extreme climate event on land, which is destructive to both the economy and ecology [30]. Drought stress has a significant negative impact on the growth and development of plants, which will make plants appear short, yellow leaves, withered branches and other phenomena.

Abscisic acid is a metabolic anti-transpirant [31], which typically plays a crucial role in plants' response to drought [32]. When plants suffer from drought stress, their ABA content will increase [33]. Under drought stress, abscisic acid (ABA) can regulate stomatal opening and closing. ABA-mediated stomatal closure reduces water loss by decreasing transpiration rate, and also enhances the water absorption and ion exchange capacity of the roots, enabling plants to better cope with arid environments and develop resistance to drought [34,35]. In addition, abscisic acid under drought conditions can also cause changes in the content of osmotic adjustment substances and osmotic adjustment capacity. Research by Zhou Lin *et al*. has shown that exogenous ABA treatment can increase the content of proline, soluble sugar, and soluble protein in tea plants under drought stress, reduce the damage caused by drought stress to tea plant seedlings, and thereby enhance their drought resistance [36]. Studies by Wang Wei *et al*. have shown that exogenous ABA can enhance the osmotic adjustment capacity of maize seedling leaves under drought conditions, which plays a certain role in improving the drought resistance of maize [37]. Abscisic acid can regulate plant stress resistance by regulating the expression of drought-resistant genes [38] and enhancing the activity of antioxidant enzymes in plant cells [39]. Relevant research indicates that under drought conditions, the early seedlings of *Onobrychis viciifolia* soaked in ABA exhibit higher activities of superoxide dismutase (SOD), peroxidase (POD), and catalase (CAT) compared to those not soaked in ABA. This suggested that ABA can enhance the antioxidant enzyme activity of *O. viciifolia* under drought stress, mitigating the oxidative damage caused by drought stress [40]. Under drought stress, ABA also alters plant morphology to resist the damage caused by drought. Studies have found that under drought stress, the plant height, leaf area, root length, and root area of maize seedlings treated with exogenous ABA are significantly increased compared to those without ABA treatment, indicating that ABA can alleviate the damage caused by drought stress to maize seedlings [41].

3.2 ABCISIC ACID AND SALT STRESS

Salt stress is one of the most common abiotic stresses in plants. At the same time, it is also the main abiotic stress that restricts plant growth and development [42]. Under salt stress, plant growth slows down, plant height decreases, and leaves turn yellow [43], which is similar to the situation of drought and water shortage. When plants are subjected to salt stress, the transcription level and expression level of ABA synthesis-related genes increase, leading to a substantial synthesis of ABA in the plant to adapt to salt stress [44]. ABA can maintain water balance and alleviate osmotic stress and ionic stress caused by excessive salt content [6]. The study on seedlings of *Toona sinensis* showed that the salt stress of external application of abscisic acid could effectively inhibit the absorption of salt ions by plants and enhanced their absorption of other nutrient ions, thus achieving ion balance, alleviating the salt stress inhibition of *T. sinensis* seedlings and resisting the harm of salt stress [45]. Under salt stress, ABA can also promote the accumulation of small organic molecules such as proline and soluble sugars in plants, thereby enhancing their salt tolerance [46]. Liu Xu and his colleagues found that exogenous ABA treatment can increase the content of proline and soluble sugar in eggplant seedlings under salt stress, enhance their osmotic adjustment ability, and enable them to better adapt to salt stress environments [47]. ABA can enhance the antioxidant defense system of plants by increasing the activity of oxidases (such as SOD, CAT, POD, etc.) to eliminate ROS and alleviate oxidative damage caused by salt

stress to plants [44]. Related studies showed that the activities of SOD, POD and CAT in tomato seedlings under salt stress for 6 days were significantly lower than those under salt stress with exogenous ABA, which indicated that ABA was involved in scavenging active oxygen and improving the salt tolerance of tomato seedlings [48].

3.3 Abscisic acid and high temperature stress

High temperature stress refers to an abiotic stress that the temperature exceeds the suitable temperature for plant growth and development, thus causes plant damage [49]. High temperature stress will limit plant growth [50], lead to premature senescence and reduce photosynthetic efficiency [51]. Research on wild Kentucky bluegrasses under high temperature stress has found that under such stress, the water content of wild Kentucky bluegrasses leaves treated with abscisic acid is higher than that of untreated leaves, indicating that ABA can mitigate the damage of high temperature stress to wild *Poa pratensis* [52]. Under high temperature stress, abscisic acid increases the activity of antioxidant enzymes and removes ROS, thus reducing oxidative damage [53]. Related studies showed that the accumulation of ROS in rice buds with ABA application was significantly lower than that without ABA application, and the content of H₂O₂ was also reduced, and the ROS scavenging genes such as *OsFe-SOD* and *OsCu/Zn-SOD* were significantly up-regulated [54]. This indicated that ABA increases the activity of antioxidant enzymes to clear excess ROS. Abscisic acid can also bind to receptors on the biofilm through signal transduction, increase the stability of the membrane system, and reduce the degree of damage to the cell membrane by high temperature [55].

4. Summary and prospect

This study reviewed the discovery, transportation, and distribution of abscisic acid, and analyzed its role in plant abiotic stresses, aiming to elucidate the importance of abscisic acid in plant resistance to abiotic stresses. Abscisic acid holds broad application prospects in enhancing plant stress resistance, particularly in drought, salt, and high temperature conditions. In addition, abscisic acid can also enhance the yield and quality of plant fruits. Relevant research has shown that exogenous application of abscisic acid can increase the concentration of carotenoids and soluble sugars, and reduce the concentration of organic acids in tomato fruits, thereby improving tomato fruit quality [56]. In fruit tree production, abscisic acid (ABA) is involved in the ripening process of fruits, which not only affects the appearance characteristics of the fruits but also makes the fruits easier to detach from the branches and facilitate harvesting [57]. Therefore, we can utilize genetic engineering techniques such as CRISPR-Cas9 and related technologies to improve the ABA signal transduction pathway [58,59], knock out or edit genes that are not conducive to ABA synthesis to increase the content of abscisic acid in plants, thereby enabling plants to better adapt to stressful environments and ensuring better yield and quality of plants.

REFERENCES

1. Esmaeili, N., Cai, Y., Tang, F., Zhu, X., Smith, J., Mishra, N., Hequet, E., Ritchie, G., Jones, D., Shen, G., Payton, P., & Zhang, H. (2021). Towards doubling fibre yield for cotton in the semiarid agricultural area by increasing tolerance to drought, heat and salinity simultaneously. *Plant biotechnology journal*, 19(3), 462-476.
2. Kumar, S., Beena, A. S., Awana, M., & Singh, A. (2017). Physiological, biochemical, epigenetic and molecular analyses of wheat (*Triticum aestivum*) genotypes with

- contrasting salt tolerance. *Frontiers in plant science*, 8, 1151.
3. Yang, X. G., Liang, W. H., Qi, Z. G., Ma, W. S., & Shen, Y. Z. (2006). Molecular Mechanisms of Plant Responses to Abiotic Stresses. *Journal of Triticeae Crops*, 26(6), 158-161.
 4. Wang, X. L., & Lv, X. F. (2020). Research progress on mechanism of nitrogen metabolism involved in plant stress resistance. *Guihaia*, 40(04), 583-591.
 5. Nehra, A., Kalwan, G., Gill, R., Nehra, K., Agarwala, N., Jain, P. K., Naeem, M., Tuteja, N., Pudake, R. N., & Gill, S. S. (2024). Status of impact of abiotic stresses on global agriculture. *Nanotechnology for Abiotic Stress Tolerance and Management in Crop Plants*. (pp. 1-21). New York, USA: Academic Press.
 6. Zhang, Y., Xu, X., Zhu, Y. X., & Guan, Y. J. (2015). Progress of Mechanisms of ABA Response to Plant Salt Stress. *Chinese Agricultural Science Bulletin*, 31(24), 143-148.
 7. Tian, X. J., Wang, Z. Y., Li, X. F., Lv, T. X., Liu, H. Z., Wang, L. Z., Niu, H. B., & Bu, Q. Y. (2015). Characterization and functional analysis of pyrabactin resistance-like abscisic acid receptor family in rice. *Rice*, 8(1), 28.
 8. Zhang, F., Wei, Q. H., Shi, J. C., Jin, X., He, Y., Zhang, Y., Luo, Q. C., Wang, Y. S., Chang, J. L., Yang, G. X., & He, G. Y. (2017). *Brachypodium distachyon* BdPP2CA6 interacts with BdPYLs and BdSnRK2 and positively regulates salt tolerance in transgenic *Arabidopsis*. *Frontiers in Plant Science*, 8, 264.
 9. Wei, H., Movahedi, A., Liu, G. Y., Li, Y. X., Liu, S. W., Yu, C. M., Chen, Y. H., Zhong, F., & Zhang, J. (2022). Comprehensive analysis of carotenoid cleavage dioxygenases gene family and its expression in response to abiotic stress in poplar. *International journal of molecular sciences*, 23(3), 1418.
 10. Gupta, K., Wani, S. H., Razaq, A., Skalicky, M., Samantara, K., Gupta, S., Pandita, D., Goel, S., Grewal, S., Hejnak, V., Shiv, A., El-Sabrou, A. M., Elansary, H. O., Alaklabi, A., & Brestic, M. (2022). Abscisic acid: role in fruit development and ripening. *Frontiers in plant science*, 13, 817500.
 11. Eagles, C. F., Wareing, P. F. (1963). Dormancy regulators in woody plants: Experimental induction of dormancy in *Betula pubescens*. *Nature*, 199, 874-875.
 12. Ohkuma, K., Lyon, J. L., Addicott, F. T., & Smith, O. E. (1963). Abscisin II, an abscission-accelerating substance from young cotton fruit. *Science*, 142(3599), 1592-1593.
 13. Cornforth, J. W., Milborrow, B. V., Ryback, G., & Wareing, P. F. (1965). Chemistry and physiology of 'dormins' in sycamore: identity of sycamore 'dormin' with abscisin II. *Nature*, 205(4978), 1269-1270.
 14. Addicott, F. T., Lyon, J. L., Ohkuma, K., Thiessen, W. E., Carns, H. R., Smith, O. E., Cornforth, J.W., Milborrow, B.V., Ryback, G., & Wareing, P. F. (1968). Abscisic acid: a new name for abscisin II (dormin). *Science*, 159(3822), 1493-1493.
 15. Zhang, Y. T., & Shi, F. L. (2020). Research progress on plant architecture formation and forage grass plant architecture. *Acta Prataculturae Sinica*, 29, (9), 203-214.
 16. Ding, B. J., Kong, X. Q., & Dong, H. Z. (2020). Research Progress on the Structure and Function of Abscisic Acid Receptor PYLs. *Molecular Plant Breeding*, 18(20), 6844-6852.
 17. Yang, H. Q., Jie, Y. L. (2001). Biosynthesis of Abscisic Acid and Its Regulation in Higher Plants. *Plant Physiology Journal*, (05), 457-462.
 18. Li, H., Li, B., & Yang, Z. (2020). Effect of exogenous ABA on endogenous hormone content of alfalfa under salt and alkali stress. *Heilongjiang Animal Science and Veterinary Medicine*, (06), 103-106+111.
 19. Chen, H. R., Wu, Z. B., He, F., Cheng, W. Y. (2001). The research progress of plant stress resistance. *Techniques and Equipment for Environmental Pollution Control*, (03), 7-13.
 20. Long, J. T., Liu, D. D., Qiao, W., Wang, Y. W., Miao, Y. J., & Baosai, H. (2024). Response of *Elymus nutans* Griseb. seedling physiology and endogenous hormones to drought and salt stress. *Scientific Reports*, 14(1), 17810.
 21. Zhou, Y. F., Wang, D. Q., Lu, Z. B., Wang, N., Wang, Y. T., Li, F. X., Xu, W. J., & Huang, R. D. (2014). Effects of drought stress on photosynthetic characteristics and endogenous hormone ABA and CTK contents in green-stayed sorghum. *Scientia Agricultura Sinica*, 47(4), 655-663.
 22. Duan, N., Jia, Y. K., Xu, J., Chen, H. L., & Sun, P. (2015). Research Progress on Plant Endogenous Hormones. *Chinese Agricultural Science Bulletin*, 31(2), 159-165.
 23. Yang, Q., Deng, X. J., Liu, T., Qian, J. Y., Zhang, P. H., Zhu, E. G., Wang, J. Q., Zhu, X. X., Kudoyarova, G., Zhao, J. Z., & Zhang, K. W. (2024). Abscisic acid root-to-shoot translocation by transporter AtABCG25 mediates stomatal movements in Arabidopsis. *Plant Physiology*, 195(1), 671-684.
 24. Gorgues, L., Li, X., Maurel, C., Martinière, A., & Nacry, P. (2022). Root osmotic sensing from local perception to systemic responses. *Stress Biology*, 2(1), 36.
 25. Kuromori, T., Sugimoto, E., & Shinozaki, K. (2014). Intertissue signal transfer of abscisic acid from vascular cells to guard cells. *Plant physiology*, 164(4), 1587-1592.
 26. Shen, C., Yang, Y. M., Sun, Y. F., Zhang, M., Chen, X. J., & Huang, Y. Y. (2022). The regulatory role of abscisic acid on cadmium uptake, accumulation and translocation in plants. *Frontiers in Plant Science*, 13, 953717.
 27. Kuromori, T., Seo, M., & Shinozaki, K. (2018). ABA transport and plant water stress responses. *Trends in plant science*, 23(6), 513-522.
 28. Cui, W., Bai, X. S., Wang, J., & Jin, H. (2022). Research Progress of ABA Hormone Regulating Biosynthesis of Disease Resistance Related Plant Secondary Metabolites. *Plant Health and Medicine*, 1(6), 1-11.
 29. Sah, S. K., Reddy, K. R., & Li, J. (2016). Abscisic acid and abiotic stress tolerance in crop plants. *Frontiers in plant science*, 7, 571.
 30. Dai, A. (2011). Drought under global warming: a review. *Wiley Interdisciplinary Reviews: Climate Change*, 2(1), 45-65.
 31. Zhang, X. Y., & Zhang, X. Y. (2014). Anti-transpirant studies and applications in agriculture. *Chinese Journal of Eco-Agriculture*, 22(8), 938-944.
 32. Huang, Q., Jiang, W. Y., & Shu, T. (2024). Genome identification and expression analysis of NHX gene family in *Solanum tuberosum*. *Jiangsu Agricultural Sciences*, (15), 44-52.
 33. Liu, H. Z., Cao, M. X., Chen, X. L., Ye, M. H., Zhao, P., Nan, Y. Y., Li, W., Zhang, C., Kong, L. S., Kong, N. N., Yang, C. H., Chen, Y., Wang, D. D. & Chen, Q. (2019). Genome-wide analysis of the lateral organ boundaries domain (LBD) gene family in *Solanum tuberosum*. *International Journal of Molecular Sciences*, 20(21), 5360.
 34. Guo, H., LV, C. P., Zheng, Z., Liu, F., & Ding, D. (2009). Advances in Research of Drought Resistance of Landscape Plants. *Anhui Agricultural Science Bulletin*, 15(7), 53-55.

35. Muhammad Aslam, M., Waseem, M., Jakada, B. H., Okal, E. J., Lei, Z., Saqib, H. S. A., Yuan, W., & Xu, W. F., Zhang, Q. (2022). Mechanisms of abscisic acid-mediated drought stress responses in plants. *International journal of molecular sciences*, 23(3), 1084.
36. Zhou, L., Xu, H., Zhu, X. J., Chen, X., Wang, Y. H., Fang, W. P., & Li, X. H. (2014). Effect of Abscisic Acid on Physiological Characteristics of Tea Plant under Drought Stress. *Journal of Tea Science*, 34(05), 473-480.
37. Wang, W., Li, D. Q., Zou, Q., & Li, C. X. (2000). Effects of ABA on Osmotic Adjustment of Leaves and Roots of *mays* Under Water Stress. *Plant Physiology Journal*, 36(6), 523-526.
38. Yu, Z. Y., Wang, X., & Zhang, L. S. (2018). Structural and functional dynamics of dehydrins: a plant protector protein under abiotic stress. *International Journal of Molecular Sciences*, 19(11), 3420.
39. Li, W. T., Ning, P., Wang, F., Cheng, X. M., & Huang, X. X. (2020). Effects of exogenous abscisic acid (ABA) on growth and physiological characteristics of *Machilus yunnanensis* seedlings under drought stress. *Chinese Journal of Applied Ecology*, (05), 1543-1550.
40. Ma, X. L., Wang, W. Y., Zhou, H. K., Li, W. J., Li, J., Li, Y., Qiu, Q. H., & Yin, H. X. (2024). Effect of soaking exogenous abscisic acid on *Onobrychis viciifolia* seed germination and physiological mechanism under drought stress. *Chinese Journal of Eco-Agriculture*, 32(11), 1882-1890.
41. Wang, X. M., Cao, L. R., & Lu, X. M. (2021). Effects of Abscisic Acid on Growth and Physiological and Biochemical Characteristics of Maize Seedlings under Drought Stress. *Molecular Plant Breeding*, 19(21), 7193-7201.
42. Ye, H., & Wang, Y. K. (2024). Roles of WRKY transcription factors in regulating leaf senescence. *Plant Physiology Journal*, (06), 905-918.
43. Qi, Q., Ma, S. R. & Xu, W. D. (2020). Advances in the Effects of Salt Stress on Plant Growth and Physiological Mechanisms of Salt Tolerance. *Molecular Plant Breeding*, 18(08), 2741-2746.
44. Liu, L., Zeng, Y. L., & Zhang, F. C. (2009). ABA and Salt Tolerance of Plant. *Plant Physiology Journal*, (02), 187-194.
45. Yao, X. M., Ou, C., Zhang, Y. L., Yang, L. M., Xu, M., Wang, Q. Q., & Qu, C. Q. (2020). Effects of Abscisic Acid on Ion Absorption and Photosynthesis of *Toona sinensis* Seedlings under Salt Stress. *Journal of Northeast Forestry University*, 48(8), 27-32.
46. Liu, Y., Wei, L., Feng, L., Zhang, M., Hu, D., Tie, J., & Liao, W. (2022). Hydrogen sulfide promotes adventitious root development in cucumber under salt stress by enhancing antioxidant ability. *Plants*, 11(7), 935.
47. Liu, X., Lin, B. Y., Li, C. X., Wu, H. Q., Zu, Y. D., & Zhong, L. M. (2020). Effects of exogenous abscisic acid on physiological characteristics of eggplant seedlings under salt stress. *Journal of Henan Agricultural University*, 54(02), 231-236+268.
48. Zhao, X. P., Yang, L., Yang, S. Y., & Tang, S. H. (2010). Effects of Exogenous ABA on Physiological Characteristics of Tomato Seedlings under Salt Stress. *Journal of Anhui Agricultural Sciences*, (27), 14833-14835.
49. Jiang, S., Wu, L. Y., Zhao, B. S., Huang, J. H., Jiang, Y. Z., Jiao, Y., & Huang, J. (2024). Molecular Mechanism of Heat Stress Tolerance in Plants: A Review. *Chinese Agricultural Science Bulletin*, (09), 132-138.
50. Wang, Y. W., Quan, S. J., Ma, H., Liu, D. H., & Xie, F. H. (2019). Mechanism of action of trehalose on plant tissues and animal cells: a review. *Jiangsu Agricultural Sciences*, 47(2), 14-18.
51. Wang, W. W., Shen, F., Wu, Y. C., Mei, Y., Zu, Y. X., Chen, C. J., Wan, H. J., & Zheng, J. Q. (2022). Bio synthesis of melatonin and its role in plant stress: a review. *Jiangsu Agricultural Sciences*, 50(1), 1-6.
52. Lei, Y. W., Bai, X. M., Wang, T., Lv, Y. W., & Lei, S. H. (2015). Alleviation Effect of ABA on the Wild Kentucky Bluegrasses under Heat Stress. *Acta Agraria Sinica*, 23(1), 89-94.
53. Tao, Z. Q., Yan, P., Zhang, X. P., Wang, D. M., Wang, Y. J., Ma, X. L., Yang, Y. S., Liu, X. W., Chang, X. H., Sui, P., & Chen, Y. Q. (2022). Physiological Mechanism of Abscisic Acid-Induced Heat-Tolerance Responses to Cultivation Techniques in Wheat and Maize. *Agronomy*, 12(7), 1579.
54. Yang, Y. Y., Chen, X., Chen, Q. Z., Lu, F., Xu, C., Yang, H. T., Su, P. P., & Liu, X. L. (2021). Priming Effects of Abscisic Acid on High Temperature Stress Tolerance in Rice at Seed Germination Stage. *Acta Agriculturae Boreali-Sinica*, 36(3), 185-194.
55. Zhang, H. (2013). Role of Abscisic Acid in Plant Hardiness Physiology. *Journal of Anhui Agricultural Sciences*, (2), 490-491.
56. Barickman, T. C., Kopsell, D. A., & Sams, C. E. (2016). Abscisic acid impacts tomato carotenoids, soluble sugars, and organic acids. *HortScience*, 51(4), 370-376.
57. Yang, X. Q., & Dou, K. (2024). Effects of Plant Hormones on Fruit Tree Growth and Environmental Adaptation. *Journal of Fruit Resources*, (03), 112-114.
58. Park, S. Y., Fung, P., Nishimura, N., Jensen, D. R., Fujii, H., Zhao, Y., Lumba, S., Santiago, J., Rodrigues, A., Chow, T. F. F., Alfred, S. E., Bonetta, D., Finkelstein, R., Provart, N. J., Desveaux, D., Rodriguez, P., McCourt, P., Zhu, J. k., Schroeder, J. I., Volkman, B. F., & Cutler, S. R. (2009). Abscisic acid inhibits type 2C protein phosphatases via the PYR/PYL family of START proteins. *Science*, 324(5930), 1068-1071.
59. Ma, Y., Szostkiewicz, I., Korte, A., Moes, D., Yang, Y., Christmann, A., & Grill, E. (2009). Regulators of PP2C phosphatase activity function as abscisic acid sensors. *Science*, 324(5930), 1064-1068.