

Bilateral Iris and Chorioretinal Coloboma: A Case Report

Youssef Ghallab^{1*}, Soukaina Daidai¹, Houssein Okeih¹, Falak Zamani¹, Zakaria Chaibi¹, Youssef Bouabid¹, Adil El khoyaali¹, Yassine Mouzari¹, Abdelbarre Oubaaz¹

¹Ophthalmology Service of the Mohamed V Military Instruction Hospital (HMIMV), Rabat, Morocco

DOI: <https://doi.org/10.36347/sjmcr.2025.v13i08.011>

| Received: 06.07.2024 | Accepted: 10.08.2024 | Published: 07.08.2025

*Corresponding author: Youssef Ghallab

Ophthalmology Service of the Mohamed V Military Instruction Hospital (HMIMV), Rabat, Morocco

Abstract

Case Report

Coloboma is a congenital anomaly characterized by a defect in one or more structures of the eye. It results from a failure in the closure of the colobomatous fissure, which typically occurs between the 5th and 7th week of embryonic development. We present the case of a 19-year-old patient from a non-consanguineous marriage, with no notable pathological history, who consulted for a progressive decrease in bilateral visual acuity. Examination with a slit lamp and funduscopy revealed bilateral iris and chorioretinal coloboma in both eyes. A thorough physical examination was performed to rule out other associated malformations, and it returned normal. Regular ophthalmological monitoring is necessary to detect any risk of retinal detachment, the main complication associated with coloboma. It is also essential to evaluate the possibility of systemic involvement, which can have serious implications for overall prognosis.

Keywords: Iris Coloboma, Chorioretinal Coloboma, Malformation, Eye.

Copyright © 2025 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

INTRODUCTION

Coloboma is a developmental anomaly of the eye that occurs during embryonic development, typically between the 5th and 7th weeks of gestation. This defect results from the failure of the fetal fissure to close and manifests as gaps in the inferonasal quadrant of the eye, where tissue is missing. Ocular coloboma is often associated with systemic anomalies when caused by chromosomal abnormalities. It can affect various structures of the eye, leading to gaps in the iris, choroid, retina, lens, eyelid, or optic nerve. Depending on the location of these gaps, several types of colobomas are distinguished [1]. Coloboma can affect one eye or both eyes simultaneously. The clinical diagnosis of chorioretinal coloboma is generally straightforward when accompanied by involvement of the anterior segment of the eye [2].

PATIENTS AND METHODS

This is a case of a 19-year-old female patient from a non-consanguineous marriage, with no notable pathological history, who presented to our clinic with a progressive decrease in bilateral visual acuity over the past 2 years. On clinical examination, visual acuity was 4/10 in the right eye (OD) and 3/10 in the left eye (OG). Examination of the anterior segment revealed a deformed pupil in both eyes, with an inferior nasal notch causing a loss of its round appearance (FIGURE 1). Fundus examination showed an inferonasal chorioretinal coloboma in the right eye, sparing the optic disc, characterized by a well-defined, whitish area with few blood vessels. In the left eye, there was a papillary coloboma extending into the interpapillary-macular and nasal regions (FIGURE 2-3). In light of these findings, a thorough physical examination was performed to rule out other associated malformations (cardiac, genitourinary, ENT, etc.), and the results were normal.




Figure 1: Inferonasal iris coloboma in both eyes




Figure 2: Fundus appearance of the right eye showing an inferonasal chorioretinal coloboma sparing the optic disc




Figure 3: Fundus appearance of the left eye showing a papillary coloboma extending into the interpapillary-macular and nasal regions

DISCUSSION

Congenital ocular colobomas result from a failure of closure of the fetal fissure during organogenesis [1]. The prevalence of colobomas is 0.5 to 0.7 per 10,000 births [2]. During embryonic development, the eye naturally features a fissure known as the colobomatous fissure, typically located in the lower and slightly nasal position, extending from the optic nerve to the pupillary margin. This fissure should normally close once eye development is complete. However, incomplete closure of this fissure can lead to partial or complete absence of certain eye structures, resulting in colobomas [3]. These anomalies can affect one or both eyes. Colobomas may arise from genetic mutations or exposure to toxic environmental factors. They are a significant cause of visual impairment and childhood blindness. Congenital ocular colobomas are frequently observed in various multisystemic syndromes, often accompanied by neurological, craniofacial, or other systemic developmental defects. Additionally, acquired colobomas can also form as a result of traumatic injury or following ocular surgical procedures [1-4]. The clinical diagnosis of chorioretinal coloboma is generally straightforward when accompanied by involvement of the anterior segment of the eye. However, it can be challenging to establish when it is isolated, especially in young children. In such cases, the presence of a whitish lesion in the fundus may suggest various retinal pathologies, including tumor-related, infectious, or genetic causes, among others [5].

Management of coloboma includes two main aspects:

Ophthalmological Care:

For iris coloboma, the use of colored (cosmetic) or corrective contact lenses may be recommended to reduce glare. In cases of chorioretinal coloboma with associated retinal detachment, surgical intervention might be necessary, though it is complex due to the delicate nature of the eye. Regular ophthalmological monitoring is crucial to detect complications, especially retinal detachment, which is a major concern [6].

Systemic Evaluation:

While most coloboma cases are idiopathic and sporadic, various modes of inheritance have been

described, sometimes associated with different syndromes such as CHARGE syndrome, Meckel-Gruber syndrome, Sjogren-Larson syndrome, Lenz syndrome, and Catel-Manzke syndrome, among others. Therefore, a comprehensive physical examination is essential to identify any associated malformations or neurological disorders and to assess for multisystemic involvement [7]. In our patient's case, the physical examination revealed no additional abnormalities.

CONCLUSION

Coloboma is a congenital malformation that can affect various ocular structures, influencing visual prognosis. The standard management for any coloboma primarily involves regular clinical and paraclinical monitoring. This approach allows for the early detection and prompt treatment of potential complications.

REFERENCES

1. Bakiri, I., Charvin, E., & Fort, A. (2022). *Les colobomes oculaires*. Sciences du Vivant [q-bio].
2. Nelson, L., Bartiss, M., & Gunton, K. (2012). *Color Atlas & Synopsis of Clinical Ophthalmology: Wills Eye Institute Pediatric Ophthalmology*. 1st ed. Philadelphia: Wolters Kluwer Health.
3. Syndicat National des Ophtalmologistes de France. Les colobomes. Available at: <http://www.snof.org/encyclopedia/colobomes>.
4. Denis, D., Girard, N., Levy Mossyconacci, A., Berbis, J., & Mantonti, F. (2013). Ocular coloboma and results of brain MRI: preliminary results. *J Fr Ophtalmol*, 36(3), 210–20.
5. El Ibrahim, F., Oudbib, M., & Baddou, T. (2018). Interest of imaging in the early diagnosis of unilateral chorioretinal coloboma. *International Journal of Innovation and Applied Studies*, 24(1), 247-250.
6. Hamma, A. (2016). A documented case of iris and chorioretinal coloboma. *Batna J Med Sci*, 3(2), 115-117.
7. Olitsky, S., & Nelson, L. (2012). *A Color Handbook of Pediatric Clinical Ophthalmology*. 1st ed. London: Manson Publishing.