

The Consumption of Iceberg Lettuce May Reduce the Adhesion of Dietary Fat to the Mucus Surface of the Stomach Barrier Lining Decreasing the Risk of Triggering Acute Gastroesophageal Reflux

Thomas J. Hurr^{1*}

¹South Australian Reflux Research Unit, Adelaide, SA, Australia

DOI: <https://doi.org/10.36347/sjams.2025.v13i11.003>

| Received: 17.09.2025 | Accepted: 31.10.2025 | Published: 03.11.2025


*Corresponding author: Thomas J. Hurr

South Australian Reflux Research Unit, Adelaide, SA, Australia

Abstract

Original Research Article

Iceberg lettuce (IL) leaves contain surface epicuticular waxes that can adsorb significant quantities of dietary fat per unit weight. Fats such as olive oil, butter, lamb fat and lard readily adsorb onto IL with soft solid fats contained in lamb fat and lard, showing strong adhesion and more resistance to removal on rinsing in a water stream. Dishwashing detergents containing surfactants prevent the adsorption of both liquid and solid fats to the lettuce leaf surface, unless at very low concentrations, highlighting their significant influence on surface properties. Based on the adhesion of fats to the IL leaf surface, it is proposed the consumption of IL could reduce acute gastroesophageal reflux (GER) and the development of gastroesophageal reflux disease (GERD). In diets containing fat, the adhesion to IL reduces the amount of fat available to adhere to the mucus surface of the stomach barrier lining, protecting the lining and reducing access by trigger foods to underlying surfaces that could potentially initiate GER.



Graphical Abstract

Iceberg lettuce leaves have approximately twice the surface area of the stomach for a 50g serve and 500ml meal volume and due to the waxy surface, can adsorb significant amounts of dietary fat. Lettuce may also adsorb on the mucus layer of the stomach barrier lining but is not regarded as a trigger food. Lettuce with its large surface area to adsorb fat, can keep fat suspended in solution, protecting the lining from fat adsorption and reduce the probability that fatty trigger foods can access underlying surfaces and cause acute gastroesophageal reflux.

Keywords: lettuce, GERD, gastroesophageal reflux, contact angle mapping, fat, trigger foods, lipophilic.

Copyright © 2025 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution **4.0 International License (CC BY-NC 4.0)** which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

1. INTRODUCTION

It has been reported that the prevalence of gastroesophageal reflux disease (GERD) is almost 10% of the world population and up to 20% of populations in western countries [1]. The diagnosis and management of gastroesophageal reflux (GER) and GERD commonly involve the use of proton pump inhibitors (PPIs) but

doubts about their safety and long-term use have emerged [2]. Research into managing GERD through dietary choices rather than the use of PPIs is being undertaken, but it is not clear if any single key element in the diet is most essential [3]. A recent two case report found consuming iceberg lettuce (IL) reduced the

severity of acute GER but no explanation why this could occur, was given and so is undertaken in this report [4].

1.1 Green vegetables, fat and reflux

It is well established that eating leafy green vegetables is beneficial to health and it has been recommended one cup $\approx$ 75g, should be consumed daily [5]. It is also well established that diets high in fat, are both a potential risk in maintaining good health and in developing GERD [6-8]. Although total fat and total vegetables have been considered individually for the risk of generating GER, the role of the dietary fat/vegetable ratio or fat/leafy green vegetable ratio was not found reported.

Cruciferous vegetables, like cabbage, kale, broccoli, radish and rocket, are part of the *Brassica* genus which includes mustards and have been associated with an increased risk of GER and irritable bowel syndrome (IBS) [7]. Lettuce is an annual plant of the *Lactuca* genus and has not generally been associated with causing GER and may be a better option for leafy green vegetables in the diet than cruciferous ones, for some individuals with GERD [7,9].

The dietary lipid/protein ratio and meal acidity have been considered important factors associated with reflux and used to develop a scale for the refluxogenic potential of foods [9]. However, the role surfactants (SF) which are a natural component of foods and the vegetable/fat interactions present during digestion, were not found reported [10-14]. Although it is known that certain foods can trigger GER (trigger foods) and can have refluxogenic scores, the chemical and physical mechanisms of interaction with the mucus layer of the stomach barrier lining or underlying structures, that may initiate and maintain the reflux process, are unknown [8,15]. The relationship between dietary components and the effect on GERD symptoms is reportedly due primarily to decreased lower esophageal sphincter (LES) tone but includes transient LES relaxations, irritation of the esophageal mucosa, increase in gastric distention, altered gastric motility and increased gastric acid production [15]. Inflammation of the stomach barrier lining or underlying structures was not reported to be associated with acute GER or GERD [15].

1.2 The surface of lettuce

A scanning electron microscopy study found leaf surfaces have complex epicuticular wax projections with various crystalline structures that protect the leaf from environmental stresses [16]. The IL leaf surface has wax projections containing 70-85% 1-hexacosanol (a C(26) hydrocarbon chain length alcohol with melting point $\approx$ 76°C) distributed in varying amounts depending on both the location on a leaf and the leaf location within the lettuce [17]. The wettability of IL leaf surfaces is indicated by water contact angles (CA) from 60-100° showing both hydrophilic and hydrophobic character,

where CA values between 0-90° indicate a hydrophilic surface and CA values greater than 90° indicate a hydrophobic surface [17,18].

1.3 Dietary fats

Dietary fats, oils and waxes come from both plant and animal sources and include the glycerolipids as triacylglycerols, diacylglycerols and monoacylglycerols (fat), fatty acids, fatty alcohols, glycerophospholipids, sphingolipids, sterol lipids, prenol lipids, saccharolipids and polyketides [17,19]. The amount of dietary fat consumed each day is reported to be on average 81-87g/day with median values as saturated (32-35g/day), mono (28-31g/day) and polyunsaturated fats (12-13g/day) for both adult men and women with or without GERD [7]. Substituting monosaturated fat or polyunsaturated fat with saturated fats was not found to increase the risk of developing GERD [7]. Gastric lipases contribute to 10-30% of the overall adult dietary fat lipolysis and the stomach can absorb short and medium chain (<12 carbon atoms) fatty acids requiring 70-90% to be passed on to the small intestine for further digestion [20].

1.4 Surfactants in the diet and the gut

Surfactants (SF) are produced by the body and used in the digestive system to emulsify fats for transport and digestion [20-23]. Surfactants are also found to occur naturally in foods, are added to commercially prepared foods, used in dishwashing detergents (DWD) and toothpastes [11-14]. Surfactants occur in saliva with a surface tension of 53-65mN/m, gastric fluid 43-54mN/m, the small intestine as bile salts 25-28mN/m, which may also be present in gastric fluid, and in DWD 22-35 mN/m [20-24].

An in vitro study to show how SF influence the digestibility of the fat tricaprylin, in the presence of the SF polysorbate 80, found that at low SF concentrations and when in the presence of other types of SF (bile salts and phospholipids) that the adsorbed SF molecules can be displaced to allow lipolysis in a concentration dependent manner [25].

Research involving the use of SF to prevent GERD was not found to be reported, possibly because lingual and gastric SF are not available for purchase and were not found included in standard (INFOGEST) in vitro gastric digestion methods [26]. This is not surprising given that the SF composition of saliva and gastric fluid are yet to be fully determined but are reported to consist of SF associated proteins A, B, C, D, fatty acids and triglycerides, pepsin, pulmonary SF like phospholipids involved in mucociliary clearance and possibly bile salts from the small intestine [20-23,27-29].

Investigation with DWD has found SF can remain on dishware after rinsing at concentrations of 0.15-0.4%, which can be above the concentration of SF required to emulsify fats in solution and can cause both gut epithelial inflammation, barrier damage and alter the

gut microbiome [14,15]. As SF has a major impact on the surface properties of fat, are available in DWD and likely to be present in the human diet, DWD were used as the source of SF in the present study.

1.5 The gut barrier system

The gut barrier system is supported by complex protective mechanisms including mucus layers, a constantly renewing epithelial boundary, tight cell junctions and the gut microbiome [30]. The mucus layers are created by various *MUC* genes expressed throughout the body, which form mucin gels that coat all non-keratinized wet epithelial surfaces [30-34]. The stomach barrier lining consists of gastric epithelial cells coated with a dense mucus (mucins MUC5AC, MUC6) [33,34]. Particles with concentrated regions of positive charge or hydrophobicity, can bind and enrich in the outer mobile gastric mucus which is cleared rapidly once binding occurs [33]. It is also reported that particles in the trachea are formed into mucin MUC5B bundles which are then coated with mucin MUC5AC, as part of the mucociliary escalator and mostly swallowed [27].

Many visual models have been created to show how high fat diets increase intestinal permeability, modulate the composition and function of the mucus barrier layer, stimulating proinflammatory signalling, inducing epithelial cell stress and enriching the gut microflora [30-34]. Earlier visual models show the gastric mucus coated with a surface layer containing dipalmitoyl phosphatidylcholine (DPPC), an important pulmonary SF found in scrapings of the stomach wall and in the gastric fluid [28,29].

The DPPC surface layer was found to be hydrophobic with a water CA > 90° in one model and with water CA 60-80° in another model, reduced after exposure to nonsteroidal anti-inflammatory drugs (NSAIDs) to a water CA < 40° due to gastric inflammation [28,29].

If the gastric mucosal layer has hydrophobic surface regions due to the presence of pulmonary SF, then it is likely dietary lipophilic fat would be readily adsorbed (dispersive-dispersive attraction between hydrocarbon molecules) and possibly in a comparable way as found for the IL surface, which also has hydrophobic surface regions and a similar water CA ≈ 60-100° [17,28,29,35].

This report investigates the relationship between IL/fat/SF adsorption and proposes that IL has a protective role by adsorbing dietary fat, reducing the probability that fat will adhere to the mucus surface of the stomach barrier lining, allowing mucosal integrity to be maintained for longer during the digestion of a fatty meal and lower the risk of developing GER and GERD.

2. MATERIALS AND METHODS

2.1. Foods and chemicals.

Iceberg lettuce (IL), olive oil (OO), butter (BU), lamb fat (LF) and lard (LD), water soluble food colouring, pasta (wheat lasagne sheets), apple (royal gala), cucumber (Lebanese), bread (wholemeal flour), minced lamb, cheese (mature), lean beef, spinach and 2 branded household DWD were purchased from local supermarkets. Lamb fat (LF) from different minced lamb samples, bought over 8 months, was extracted by heating to generate multiple samples of fat with several samples combined. Hydrochloric acid (HCl) and Sudan 111 (lysochrome dye) were purchased from Chem Supply (Adelaide). The IL was used directly as purchased without washing while the apples and cucumbers were washed first. Pasta was soaked in water for 30 minutes before use. The lean beef was pan fried without oil. Dish washing detergents as DWD1 contain coconut based anionic and non-ionic SF, DWD2 contain cationic (polyquaternium-7, cocamidopropyl betaine), anionic (sodium laureth sulphate) and non-ionic (capryloyl /caproyl methyl and lauroyl /myristoyl methyl glucamides) SF.

2.2. Hydrophilic/lipophilic dye solutions, rinsing and quantifying fat adsorption

All prepared solutions and rinsing or washing was carried out at 37°±3°C. To replicate a gastric solution, water or 0.1M HCl, without the inclusion of pepsin, mineral salts or lingual or gastric SF or lipases, was used [25]. To test the adhesive strength of the adsorbed fats to the IL leaf surface after agitation in prepared solutions for 30 seconds (s), the leaves were held vertically in air for 30s to determine how much fat (grams g) remains adhered under gravity and then rinsed in a water stream (WS) of running warm water for 30s at a flow rate of 42±2ml/sec (1260±60ml /30s) and again held vertically in air for 30s to again determine how much fat remains adhered.

The amount of fat remaining adhered was termed the retention weight (RW). The WS was thought an aggressive way to determine the strength of IL/fat adhesion and although not the same process as gastric digestion, does allow visual observation of changes to the IL surface. The WS lacked careful temperature control, as solid LF and LD can have a softening or melting point around 37-38°C, with solid fat more resistant to removal and potentially increased the error in measuring the RW.

To prepare a solution to show the macroscopic hydrophilic nature of IL, approximately ≈ 0.5ml of green or blue food colouring dye was added to 90ml of 0.1M HCl. To prepare a solution to show the macroscopic lipophilic nature of the IL, approximately ≈ 0.005g of solid Sudan 111 (inclusion of solvents could dissolve epicuticular wax) was added to 10ml of either OO, BU, LF or LD to form a red liquid dye which can also

contained some solids (LF or LD) and undissolved red dye particles. To determine the macroscopic hydrophilic/lipophilic surface properties of IL or other foods, the 2 dyed aqueous and fat solutions were mixed to form a solution containing 10% fat, shaken for 30s and allowed to stand for 10 minutes. At 37°C, the OO and BU were both liquids, LF consisted of both liquid and soft solids and LD consisted of both liquid and small particulate soft crystalline fat and with all fats, having a lower density than water, floating on the aqueous surface.

To determine the RW on the IL surface (after 30s under gravity and 30s in a WS), samples with the same surface area (SA) were required and cut from both inner and outer leaf regions of the IL as 5cm x 5cm = 25cm² pieces. Samples with an initial weight (w_0) were agitated for 30s in the solutions, removed, held vertically, allowed to drain in air for 30s and re-weighed (w_1).

$$RW (g) \approx w_1 - w_0 \quad (1)$$

The samples were then held under a WS (standard deviation (SD) used for all numerical values of at least 3 repeat measurements) for 30s allowed to drain for 30s and re-weighed (w_2).

$$RW (g) \approx w_2 - w_0 \quad (2)$$

To calculate the average RW (ARW) so that comparison between samples could be made, the average initial weight $w_0(av)$ of IL samples with the same SA, was determined from 66 samples of 25cm² pieces of IL leaf which varied from 0.99 g for the thinner outer leaf to 4.51g for the thicker inner leaf near the core with $w_0(av) \approx 2.24 \pm 0.89g/25cm^2$.

To convert the RW of material adhered after dipping or rinsing, to ARW as g/100g IL, using the average weight of a 25cm² section of IL;

$$ARW (g/100g IL) \approx 100 \times (w_1 - w_0) / w_0(av) \quad (3)$$

or for samples rinsed in a WS

$$ARW (g/100g IL) \approx 100 \times (w_2 - w_0) / w_0(av) \quad (4)$$

In this report, a region on the IL surface was arbitrarily assigned hydrophilic if it retained water, lipophilic if it retained fat and hydrophobic if it did not retain water, when removed from solutions containing water, fats, 0.1M HCl or DWD and allowed to drain in air for 30s [18,35]. Stable dry patches could also form on the IL leaf surface on the break-up of the solutions, which drained as rivulets, when the surfaces were held vertically, as noted previously for adiabatic surfaces and considered in this report as regions that do not show strong hydrophilic or lipophilic surface properties, under these conditions [36].

2.3. The use of dish washing detergents /surfactants.

It was recognised that the surface properties of fats are greatly influenced by the presence of SF, even at very low concentrations and as human gastric SF is not available and at least some SF effects should be investigated, DWD was used. Commercial DWD contain SF at concentrations of ≈ 10 -20% or 100-200g pure SF/litre so 0.1% DWD solutions used in this report, would contain 1g of the 10-20% DWD/litre or 0.1 – 0.2 g pure SF/litre [24]. Mixed SF solutions, as found in DWD, can have a critical micelle concentration (CMC) between 0.1-0.32g/litre [24]. The DWD that contain both amphoteric and non-ionic surfactant mixtures, can promote CMC lowering and the CMC values of the DWD was reported not to vary by more than 5% between 17-42°C [24].

To make the solutions containing DWD, pure fats were first dyed with $\approx 0.005g$ of solid Sudan 111 and then added to the 0.1M HCl/(0.1 – 0.5%) DWD solutions, shaken for 30s resulting in an emulsion of red fat droplets initially but within minutes formed a creaming layer (floating dispersed phase of the emulsion), allowed to stand for 10 minutes and to this solution the IL were added. If the fats were first shaken in water with 0.1-0.5% DWD1 or DWD2, the fat would not adsorb the solid Sudan 111 even after 8 hours, indicating the dye could not accessed lipophilic surface sites and that the fat was in a stable emulsion. For very dilute DWD solutions as 0.1 M HCl/10% OO/0.05% DWD, Sudan 111 would readily dissolve in OO despite the presence of DWD, to give a red floating coalesced layer of fat, indicating that a stable dispersion had not formed. No significant differences were observed when using either DWD1 or DWD2 when forming the fat solutions.

2.4. Hydrophilic /lipophilic nature of common food surfaces

To compare the adhesion of fat on IL with other common foods, intact samples IL, cheese, sliced apple, sliced cucumber, bread, pasta, dry pan fried beef and spinach samples ($\approx 25cm^2$) were agitated in either 0.1M HCl/10% LF or 0.1M HCl/10% LF/0.1-0.5% DWD2 solutions for 30s, removed, held vertically and allowed to drain in air for 30s then agitated in water-bath (a WS would separate the bread matrix) for 30s to rinse any un-adsorbed fat from the surface, held vertically and allowed to drain in air for 30s again and re-weighed.

2.5. The retention weight of absorbed or adsorbed fats

To determine if water, acid solutions, SF or fats were absorbed into the interior of the intact IL leaf rather than adsorbed on the surface, samples were rinsed and then wiped with tissue paper and re-weighed to determine if an increase in the initial weight had occurred.

2.6. The ratio of lettuce: stomach surface area.

The IL: stomach surface area (SA) ratio should give an indication of the potential significance of IL as a competitive surface for the adsorption of fats. Magnetic resonance imaging of the stomachs of 12 healthy volunteers, after consuming 500cm³ of soup, found the maximum macroscopic stomach SA = 536±25 cm² (≈ 23 x 23cm) initially, decreasing over time, as the food was digested [37]. The macroscopic SA, using both sides of the IL leaf, can be calculated from $w_o(av) \approx 2.24g/25cm^2$ such that the SA of 100g IL is $(100/2.24) \times 25 \times 2 \approx 2232cm^2$ (≈ 47 x 47cm) or for 1 serve of IL the SA is ≈ 1674cm²/75g IL (≈ 41 x 41cm) and for a smaller serve of 50g the SA ≈ 1116cm²/50g IL (33.5 x 33.5cm). The ratio of IL: stomach SA, for 1 serve of IL and a 500ml meal, is calculated as IL: stomach SA ≈ 1674:536 ≈ 3:1. For a smaller serve of 50g IL then IL: stomach SA = 1116:536 ≈ 2:1. If the meal size is reduced proportionally to the amount of IL consumed, the ratio of IL: stomach SA would be similar.

3. RESULTS

3.1 Mapping the hydrophilic/lipophilic surface of intact IL

The intact IL leaf surfaces, when agitated in the dye solutions are shown to contain both hydrophilic (aqueous 0.1 M HCl dyed green), lipophilic (fat dyed red) and regions that appear smooth which do not hold either liquid, Figure 1. The aqueous hydrophilic green dye also absorbed into the cut surface edges of the IL, indicating a hydrophilic interior, Figure 1A-D. The use of mixed green and red dyed water/fat solutions to map surface hydrophilic/lipophilic surface sites was not found previously reported for food surfaces, only the use of water contact angles (sessile drop) to determine the local surface hydrophilic/hydrophobic nature. From the appearance of the IL leaf, the red dyed LF and LD were the more adhesive and more resistant to removal under a WS, compared to OO and BU, Figure 1.




Figure 1: The adsorption of 0.1M HCl/10% fat solutions to the IL leaf surface showing hydrophilic (green) and lipophilic (red) surface sites after removal from solution and after rinsing in a WS. The weighed samples of IL (25cm²) were dipped in green dyed 0.1M HCl with 10% red dyed fat solution, agitated for 30s, then held vertically in air for 30s to drain, and re-weighed (A, C, E, G) followed by rinsing in a WS and again held vertically in air for 30s to drain and again re-weighed (B, D, F, H). A. OO, ARW = 17g/100g IL; B. OO, ARW = 6.7g/100g IL; C. BU, ARW = 20g/100g IL; D. BU, ARW = 5.8g/100g IL; E. LF, ARW = 28g/100g IL; F. LF, ARW = 10g/100g IL; G. LD, ARW = 41g/100g IL; H. LD, ARW = 8.9g/100g. The ARW values show the combined weights of the adsorbed 0.1M HCl/fat solution but it can be seen from the amount of red dye, a significant proportion of the ARW value can be associated with fat adsorption particularly for the solid LF and LD

3.2 The adsorption of fats to IL

The average retention weight (ARW) values for the intact IL leaves are given for water, 0.1M HCl solution, pure fat (OO, BU, LF and LD) and 0.1M HCl/10% fat solutions includes the use of 0.05-0.5% DWD1, Table 1. It is found that the ARW values for water (11±3.0g/100g IL) or 0.1M HCl (11±2.1g/100g IL) are similar, showing the adsorption of water to the IL surface was not significantly influenced by the ionic strength or pH of the solution and under the more dynamic conditions of a WS, the ARW are also similar, but lower, Table 1. The ARW values for pure liquid fats,

OO and BU (ARW ≈ 18-19±4.0g/100g IL) were also surprisingly similar for OO and BU in 0.1M HCl (ARW ≈ 17-18±2.9g/100g IL) without the expected increase in weight due to the additional adsorption of 0.1M HCl, Figure 1A, 1C, Table 1. This result indicates the 0.1M HCl solution may change the surface properties of OO, BU or IL possibly by restructuring the behaviour of specific surface molecules selectively exposing hydrophilic or lipophilic parts [35].

For pure LF, the ARW ≈ 43±34g/100g IL has a large standard deviation, reflecting variations in the

amounts of fat that adsorbed from mixtures of both liquid and solid LF, with lower values resulting from predominantly liquid fat adsorption, which could drain from the surface when held vertically, to higher values when solid fats adsorb, that resisted draining. Solid LF was also more readily retained on rinsing in a WS (ARW $\approx 30 \pm 27$ g/100g IL) than the liquid only fats (OO and BU, Table 1). Lamb fat from the LF/0.1M HCl solutions could adsorb as a large soft solid ball (ARW $\approx 102 \pm 49$ g/100g IL) which resists removal in a WS (ARW $\approx 54 \pm 33$ g/100g IL, Table 1). The variations in the amount of LF that adsorb, may be due to the range in the melting points for animal fat, which depend on the origin from the animal body, with back fat 30–40°C, leaf fat 43–48°C and mixed fats 36–45°C.³⁸ Lard, like LF, is composed of both liquid and solid fats with ARW $\approx 38 \pm 13$ g/100g IL and after rinsing in a WS, ARW $\approx 17 \pm 7.1$ g/100g IL for the LD/0.1M HCl solutions but with lower standard deviation values, due to the more consistent composition of a commercial product than for LF, extracted from different mince samples.

The ARW values for IL agitated in water/0.5% DWD1 (7.1 ± 0.45 g/100g IL) or in 0.1M HCl/0.5% DWD1 (5.2 ± 0.93 g/100g IL) were similar but lower than the ARW when only water or 0.1M HCl solutions were present (table 1). The lower ARW values when DWD1 was present may be due to the difference in surface

tension between water and water/SF solutions, with the lower surface tension SF solutions draining from the IL surface more efficiently. When water/0.5% DWD1 and 0.1M HCl/0.5% DWD1 were rinsed in a WS, the ARW values increased, possibly due to increased wetting of the surface regions between the epicuticular waxes.

The ARW values for IL agitated in 0.1M HCl/10% fat (OO, BU, LF, LD)/0.1–0.5% DWD1 solutions and then held vertically in air, were lower than those in the absence of DWD1, with fat adsorbing as part of an emulsion and readily rinsed from the IL surface within seconds in a WS. The same results were also found for 0.1M HCl/10%OO/0.1% DWD2 solutions, where on rinsing in a WS, all adsorbed dyed OO was removed from the IL leaf surface, Figure 2A, 2B.

Only when the DWD1 was at a very low concentration of 0.05% could dyed OO adsorb on the IL leaf surface and remain adsorbed, after rinsing in a WS, indicating the significant influence of SF on IL/fat interactions, Table 1. Epicuticular wax structures remained intact on the IL despite rinsing in DWD2 as OO could re-adsorb from a 0.1M HCl/10%OO solution on to the IL surface (figure 2C–2G). Scanning electron microscopy (SEM) images of soybean leaves show SF solutions can partially remove some but not all epicuticular wax from these leaf surfaces [39].

Table 1: The strength of water and fat adhesion to IL is indicated by ARW (g/100g IL) for samples held vertically in air and under a WS (equations 3 and 4). The liquids used were water, 0.1M HCl, pure fats (OO, BU, LF and LD), fats as 10% solutions with 0.1M HCl with or without DWD1. The IL was agitated in the solutions for 30s, removed, held vertically in air for 30s and allowed to drain and re-weighed then held under a WS for 30s, then again allowed to drain for 30s and again re-weighed. The ARW values for the pure fats (without water) indicate fat only retention prior to rinsing in a WS, all other ARW values include the weight of adsorbed water or DWD1, with the amount of dyed fat adsorbed being visually apparent. After rinsing in a WS, green dye is removed but dyed fats (red) can remain adsorbed, with the fats containing solids (LF and LD) showing the greatest resistance to removal

Water, 0.1M HCl, fat or 0.1M HCl/fat solutions at 37°C	IL agitated in solutions then held vertical in air for 30s. ARW (g/100g IL)	IL held vertically under a WS for 30s then held vertically in air for 30s. ARW (g/100g IL)
water	11 ± 3.0 (green dye)	8.5 ± 5.3 (water only)
HCl	11 ± 2.1 (green dye)	5.5 ± 1.4 (water only)
OO pure	18 ± 4.0 (red dye)	8.9 ± 3.1 (red dye)
BU pure	19 ± 1.8 (red dye)	7.9 ± 5.4 (red dye)
LF pure	43 ± 34 (red dye)	30 ± 27 (red dye)
LD pure	33 ± 7.3 (red dye)	17 ± 5.4 (red dye)
OO/0.1M HCl	18 ± 2.7 (green and red dye)	8.5 ± 2.8 (green and red dye)
BU/0.1M HCl	17 ± 2.9 (green and red dye)	7.5 ± 3.5 (green and red dye)
LF/ 0.1M HCl	102 ± 49 (green and red dye)	54 ± 33 (green and red dye)
LD/0.1M HCl	38 ± 13 (green and red dye)	17 ± 7.1 (green and red dye)
Water/0.5% DWD1	7.1 ± 0.45 (green dye)	9.7 ± 4.1 (water only)
HCl/0.5% DWD1	5.2 ± 0.93 (green dye)	8.0 ± 3.5 (water only)
OO/0.1M HCl/0.05% DWD1	27 ± 4.8 (green and red dye)	15 ± 9.3 (green and red dye)
OO/0.1M HCl/0.1% DWD1	11 ± 2.3 (green and red dye)	18 ± 3.7 (water only)
OO /0.1M HCl/0.5% DWD1	8.3 ± 2.3 (green and red dye)	11 ± 2.4 (water only)
BU/0.1M HCl/0.5% DWD1	7.9 ± 1.9 (green and red dye)	11 ± 5.1 (water only)
LF/ 0.1M HCl/0.5% DWD1	9.5 ± 2.1 (green and red dye)	16 ± 8.5 (water only)
LD/0.1M HCl/0.5% DWD1	11 ± 0.52 (green and red dye)	19 ± 2.1 (water only)

3.3 Fat adsorption or absorption

The water, pure fats, fats/0.1M HCl and fats in DWD solutions that had adsorbed onto the IL and were

liquids at 37°C, could be wiped from the IL leaf surface using tissue paper (ARW $\approx$ 0g/100g IL) showing water, fats and DWD did not significantly absorb into the bulk of the IL.

3.4. The adsorption of fats on common food surfaces.

The IL, cheese, apple, cucumber, bread, pasta, meat and spinach were agitated for 30s in a blue dye solution of 0.1M HCl/10% LF dyed with Sudan 111, held vertically in air for 30s, allowed to drain and washed in a water bath at 37°C, Figure 3. A LF sample that strongly adhered to IL was chosen to test the adhesion of fat to the other food surfaces, Figure 3A.




Figure 2: The influence of OO adsorption to IL from OO/DWD2 solutions (A-B) and on rinsed in DWD2 (C-G). A. a sample of IL ($\approx$ 25 cm²) agitated for 30s in green dyed 0.1M HCl/ red dyed 10%OO/0.1% DWD2, then held vertically in air for 30s, B. the sample in (A) was then rinsed in a WS and allowed to drain for 30s. The sequence A-B shows OO weakly adsorbed from DWD2 solution and is easily rinsed from the IL surface C. a sample of IL ($\approx$ 25 cm²) agitated for 30s in green dyed 0.1M HCl/ red dyed 10%OO (no DWD2), then held vertically in air for 30s showing both water and OO adsorption, D. the sample shown in (C) was agitated in DWD2 for 30s held in air and allowed to drain for 30s showing some OO was removed. E. the sample shown in (D) was then rinsed in a WS held in air and allowed to drain for 30s showing the OO rinsed away with red particles on the IL surface from undissolved solid Sudan 111. F. the sample shown in (E) was re-agitated in the original 0.1M HCl/10%OO solution (no DWD2) used to generate sample C, held in air and allowed to drain for 30s shows OO can re-adsorb. G. the sample in (F) was re-rinsed in a WS then held in air and allowed to drain for 30s showing most of the OO rinsed away. The sequence (C-G) shows that OO adsorbed to the IL surface in (C) is partly removed from the surface by DWD2 in (D) and completely removed in a WS in (E) but DWD2 does not remove the surfaces epicuticular waxes as shown in (F) as OO can re-adsorb on to the IL surface with residual OO still remaining after rinsing in a WS in (G).

The macro-nutrient composition and assigned hydrophilic or lipophilic surface properties of the foods, based on visual observations of the amounts of dye adsorbed, are shown in Table 2 [40]. The IL leaf strongly adsorbed LF and this was not easily washed from the surface in a water-bath (ARW $\approx$ 53g/100g IL, Figure 3A). Cheese adsorbed mainly aqueous blue dye indicating a mostly hydrophilic surface with some small lipophilic red patches, Figure 3B. The lipophilic apple and cucumber skins could adsorb LF, but the fat could

readily be wiped from the smooth skin surface, Figure 3C, 3D. Bread showed a strong adsorption of blue dye in the inner regions and red dyed LF on the crust where fat (grease) had been used on the baking container, while pasta had a completely hydrophilic surface, Figure 3E, 3F. The dry pan-fried beef was found to be hydrophilic despite containing protein, known to consist of amino acids, some of which contain hydrophobic chemical groups [41].

Table 2: Values for the macro-nutrient composition of foods from Food Standards Australia and New Zealand, Australian food composition database and the classification of food surface properties based on the appearance of dyes after adsorption from solutions containing both blue dyed 0.1M HCl, showing hydrophilic surface regions and red dyed LF, showing lipophilic surface regions, Figure 3 [40]. The weight % values may not add to 100%

due to minerals and other micro-components. ¹Note IL does not have a value recorded for fat as amounts are presumably not significant for macro-nutrient compositions. ²g/100cm³ converted to g/100g; ³estimated values

Food	Water weight %	Carbohydrate (starch, sugars and fibre) weight %	Protein weight %	Fat weight %	Classification of surface properties using dyed aqueous /fat solutions
iceberg lettuce	96	2	0	0 ¹	hydrophilic /lipophilic
cucumber (Lebanese)	96	3	0	0	hydrophilic except skin
spinach (fresh raw)	92	2.5	3	0	hydrophilic /hydrophobic
apple (royal gala)	87	12	0	0	hydrophilic except skin
pasta (white wheat)	63	28	5	0	hydrophilic
bread (wholemeal flour)	39	46	10	2-3	hydrophilic
cheese (mature)	34	0	25	34	hydrophilic /lipophilic patches
beef (dry fried)	65	0	32	3	hydrophilic
olive oil (OO)	0	0	0	100 ²	lipophilic
butter (BU)	15	0	1	81	lipophilic
lamb fat (LF)	0	0	0	100 ³	lipophilic
lard (LD)	0	0	0	100 ³	lipophilic




Figure 3: Common foods ($\approx 25\text{cm}^2$) agitated in aqueous 0.1M HCl(blue)/10% LF (red) solution for 30s, allowed to drain in air for 30 s, washed in a water bath for 30s, then held in air and allowed to drain for 30s, at 37°C. The adsorption of aqueous blue dyed 0.1M HCl solution shows hydrophilic surface regions and red dyed LF show the lipophilic surface regions. A. IL, ARW $\approx 53\text{g}/100\text{g}$ IL after washing, B. cheese, C. apple, D. cucumber, E. bread, F. pasta, G. fried beef and H. spinach, RW $\approx 59\text{g}/100\text{g}$ spinach after washing (RW = $w_2 - w_0 / w_0$ where $w_0 = 1.29\text{g}$, $w_2 = 2.05\text{g}$)

Spinach, like IL, has a surface which contains epicuticular waxes and was found to adsorb LF in patches with some smooth sections that did not strongly adsorb either water or fat, Figure 3H. If DWD2 (0.1-0.5%) was included in the fat solutions, none of the foods adsorbed significant amounts of LF after washing in a water bath.

4. DISCUSSION

This study has demonstrated that by using mixed dyed aqueous/fat solutions, the hydrophilic/lipophilic surface properties of food can be identified, Figure 1-3. Raw leafy green vegetables like lettuce with an outer protective layer of epicuticular waxes, can adsorb significant quantities of dietary fat.

Apart from spinach, other common foods tested do not maintain the adsorption of any significant amounts of dietary fats from the 0.1M HCl /fat solutions after rinsing in a water bath, despite some containing protein and fat (cheese and beef, Table 2, Figure 3).

Intact IL can adsorb dietary fats, with soft solid LF and LD showing higher ARW values and greater resistance to removal under a WS, than the liquid fats, OO or BU at 37 °C. Figure 1, Table 1. The ARW values for the 4 pure fats (OO, BU, LF, LD, drained in air for 30s) can be added together as $18+19+43+33 \approx 28.2\text{g}/100\text{g}$ IL or $21\text{g}/75\text{g}$ IL to give an average value for lipophilic fat adsorption to IL, Table 1. If the average consumption of fat is $\approx 81\text{-}87\text{g}/\text{day}$ then 1 serve of IL (75g) could adsorb $\approx 21/84 \approx 25\%$ of average daily fat

consumption [7]. Although only a very approximate result and the influence of dietary SF unclear, it still indicates the potential significant value IL may have in adsorbing and suspending lipophilic dietary fat as fat/IL particles in the gastric fluid, reducing fat exposure and possible adhesion or absorption to the mucus surface of the stomach barrier lining. In the presence of DWD, both liquid and solid fats poorly adsorb on to the IL leaf surface and the fats that did adsorb, were readily removed on rinsing under a WS unless the concentration of DWD was very low, Table 1, Figure 2A, 2B. It was unexpected that the SF in DWD, designed to clean surfaces, would be so powerful in preventing the adsorption of both liquid and solid fats to the IL leaf surface, even at low concentrations, suggesting studies into the adsorption of fats to food surfaces should consider the influence of SF, Figure 2A, 2B.

A model, showing how fat and trigger foods interact in the presence and absence of IL with the mucus surface of the stomach barrier lining, is proposed in the Graphical Abstract. If 50g of IL in the diet is consumed with a meal of volume 500ml, then the IL has approximately twice the SA as the stomach and so can potentially adsorb twice as much dietary fat, assuming the kinetics of adsorption are similar, which may be probable given the gastric mucosal surface may contain an outer hydrophobic layer with a similar CA as the IL surface [17,28,29]. The waxy surface of IL may adsorb on the mucus surface of the stomach barrier lining, but IL is not recognised as a trigger food and due to the large lipophilic surface area of IL, reduce the amount of fat that can adsorb on the mucus layer. The IL/fat and fat that adhere and slough off the mucus surface together with the IL/fat particles that do not adhere to the mucosal surface, progress to the small intestine to be digested by bile salts and lipases [23,24]. Benefits of consuming IL may also extend to diets that consume foods with low natural SF content or lack sufficient lingual or gastric SF to emulsify fats, which could otherwise reduce the concentration of adhesive fat deposits on the gastric mucosal surface.

If commercially prepared SF from DWD were present, the fats would be dispersed in the gastric solution, but SF may also damage the mucosal layer and potentially the underlying epithelial cells together with changes to the mucosal microbiome [13,14]. It is likely SF have an important role in the digestion of fat in the gastric environment but must be of the appropriate chemical composition and concentration to be beneficial [32,42].

It is still unknown if the interactions between lettuce, fat, SF and the mucus surface have a role in reducing GER or GERD. Further studies could consider particulate IL leaves, fat/SF/gastric mucosa interactions, gastric SF, the gut/lung axis and the role of lipases.

5. CONCLUSION

The large lipophilic surface area per unit weight of IL, lacking in other common foods, gives IL the potential to adsorb significant quantities of dietary fat, with solid fats more resistant to removal on rinsing in water than liquid fats. It was found that DWD can prevent the adsorption of fat to the IL leaf surface, except at very low concentrations, indicating a potentially significant role of SF in dietary fat/food interactions. An explanation why IL in the diet could reduce GER and GERD is proposed, based on the competitive adsorption of dietary fats between the IL leaf and the stomach barrier lining. The strong adhesion of solid fats to IL may allow fats to pass from the gastric environment to the small intestine, minimizing irritation and adhesion to the mucus layer of the stomach barrier lining, lowering the probability trigger foods can access underlying structures and cause GER and GERD. The ability of IL surfaces to adsorb dietary fats may prove to be a supporting factor in emerging evidence that the consumption of vegetables, which includes lettuce and leafy vegetables, can be negatively correlated with the risk of developing GERD.

Funding The author has not received funding from this work.

Competing Interests The author declares that there are no competing interests.

Abbreviations: ARW average retention weight, BU butter, CA contact angle, DWD dishwashing detergent, GER gastroesophageal reflux, IL iceberg lettuce, LD lard, LF lamb fat, OO olive oil, RW retention weight, s second, SA surface area, SF surfactant, WS water stream.

REFERENCES

1. Li, N., Yang, WL., Cai, MH. *et al.*, Burden of gastroesophageal reflux disease in 204 countries and territories, 1990–2019: a systematic analysis for the global burden of disease study 2019. BMC Public Health. 2023; 23:582-595.
2. Katz PO, Dunbar KB, Schnoll-Sussman FH *et al.*, ACG Clinical guideline for the diagnosis and management of gastroesophageal reflux disease. Am J Gastroenterol. 2022;117:27-56.
3. Lechien JR, Crevier-Buchman L, Distinguin L *et al.*, Is diet sufficient as laryngopharyngeal reflux treatment? a cross-over observational study. Laryngoscope. 2022; 132:1916-1923.
4. Hurr TJ. Iceberg lettuce reduces the severity of gastroesophageal reflux symptoms: two case reports. Sch J Med Case Rep. 2023; 7:1373-1376.
5. Eat for health Australian dietary guidelines summary. Australian government national health and medical council. 2013:1-153. www.eatforhealth.gov.au
6. Zhang M, Hou ZK, Huang ZB *et al.*, Dietary and lifestyle factors related to gastroesophageal reflux disease: A systematic review. Ther Clin Risk Manag. 2021; 17:305-323.

7. Wang SE, Hodge AM, Dashti SG *et al.*, Diet and risk of gastroesophageal reflux disease in the Melbourne collaborative cohort study. *Pub Health Nutr.* 2021; 24:5034-5046.
8. Heidarzadeh-Esfahani N, Soleimani D, Hajiahmadi S *et al.*, Dietary intake in relation to the risk of reflux disease: a systematic review. *Prev Nutr Food Sci.* 2021; 26:367-379.
9. Lechien JR, Bobin F, Mouawad F *et al.*, Development of scores assessing the refluxogenic potential of diet of patients with laryngopharyngeal reflux. *Eur Arch Otorhinolaryngol.* 2019; 276:3389-3404.
10. Yang X, Gil MI, Yank Q *et al.*, Bioactive compounds in lettuce: highlighting the benefits to human health and impacts of preharvest and postharvest practices. *Compr Rev Sci Food Saf.* 2022; 21:4-45.
11. De S, Malik S, Ghosh A *et al.*, A review on natural surfactants. *RSC Advances.* 2015; 5:65757-65767.
12. Dembitsky VM. Astonishing diversity of natural surfactants: 1. Glycosides of fatty acids and alcohols. *Lipids.* 2004; 39:933-53.
13. Ogulur I, Pat Y, Aydin T A *et al.*, Gut epithelial barrier damage caused by dishwasher detergents and rinse aids. *J Allergy Clin Immunol.* 2023;151:469-484.
14. Gerasimidis K, Bryden K, Chen X *et al.*, The impact of food additives, artificial sweeteners and domestic hygiene products on the human gut microbiome and its fibre fermentation capacity. *Eur J Nutr.* 2020; 59:3213-3230.
15. Özenoğlu A, Anul N, Özçelikçi B. The relationship of gastroesophageal reflux with nutritional habits and mental disorders. *Hum Nutr Metab.* 2023;33:1-7.
16. Barthlott W, Neinhuis C, Cutler D *et al.*, Classification and terminology of plant epicuticular waxes, *Bot J Linnean Soc.* 1998;126:237–260.
17. Ku K, Chiu Y, Shen C *et al.*, Leaf cuticular waxes of lettuce are associated with reduced attachment of the foodborne pathogen *Salmonella* spp. at harvest and after postharvest storage. *Food Sci. Tech* 2020; 117:108657,1-6.
18. Law KY. Definitions for hydrophilicity, hydrophobicity, and super hydrophobicity: getting the basics right. *J. Phys. Chem. Lett.* 2014; 5:686–688.
19. Ratnayake WM, Galli C. Fat and fatty acid terminology, methods of analysis and fat digestion and metabolism: a background review paper: review. *Ann Nutr Metab.* 2009; 55:8-43.
20. Gittings S, Turnbull N, Henry B *et al.*, Characterisation of human saliva as a platform for oral dissolution medium development. *Eur J Pharm Biopharm.* 2015; 91:16-24.
21. Rivera KR, Pessi J, Andersson V *et al.*, Characterizing interspecies differences in gastric fluid properties to improve understanding of in vivo oral drug formulation performance. *Eur J Pharm Sci.* 2023; 183:106386,1-11.
22. Acevedo-Fani A, Singh H. Biophysical insights into modulating lipid digestion in food emulsions. *Prog Lipid Res.* 2022; 85:101129,1-14.
23. Macierzanka A, Torcello-Gómez A, Jungnickel C *et al.*, Bile salts in digestion and transport of lipids. *Adv Colloid Interface Sci.* 2019; 274:102045,1-14
24. Blagojević S, Blagojević SM, Pejić ND. Performance and efficiency of anionic dishwashing liquids with amphoteric and nonionic surfactants. *J Surfactants Deterg.* 2016; 19:363-372.
25. Jacobsen AC, Kabelev A, Sinko PD *et al.*, Intrinsic lipolysis rate for systematic design of lipid-based formulations. *Drug Deliv Transl Res.* 2023;13:1288-1304.
26. Mulet-Cabero AI, Egger L, Portmann R *et al.*, A standardised semi-dynamic in vitro digestion method suitable for food – an international consensus. *Food Funct.* 2020; 11:1702-1720.
27. Ermund A, Meiss LN, Rodriguez-Pineiro AM *et al.*, The normal trachea is cleaned by MUC5B mucin bundles from the submucosal glands coated with the MUC5AC mucin. *Biochem Biophys Res Commun.* 2017; 492:331-337.
28. Hills BA. Gastric surfactant and the hydrophobic mucosal barrier. *Gut.* 1996; 39:621-4.
29. Lichtenberger LM, Zhou Y, Dial EJ *et al.*, NSAID injury to the gastrointestinal tract: evidence that NSAIDs interact with phospholipids to weaken the hydrophobic surface barrier and induce the formation of unstable pores in membranes. *J Pharm Pharmacol.* 2006; 58:1421-8.
30. Rohr MW, Narasimhulu CA, Rudeski-Rohr TA *et al.*, Negative effects of a high-fat diet on intestinal permeability: a review, *Advances in Nutrition.* 2020; 11:77–91.
31. Gulhane M, Murray L, Lourie R *et al.*, High fat diets induce colonic epithelial cell stress and inflammation that is reversed by IL-22. *Sci Rep.* 2016; 6:28990,1-16.
32. Lock JY, Carlson TL, Wang CM *et al.*, Acute exposure to commonly ingested emulsifiers alters intestinal mucus structure and transport properties. *Sci Rep.* 2018; 8:10008,1-13
33. McShane A, Bath J, Jaramillo AM *et al.*, Mucus. *Curr Biol.* 2021; 31: R938-R945.
34. Subramanian DA, Langer R, Traverso G. Mucus interaction to improve gastrointestinal retention and pharmacokinetics of orally administered nano-drug delivery systems. *J Nanobiotechnol.* 2022;20:362.
35. Wang Y, Wang Q, Wang B *et al.*, Modulation of solid surface with desirable under-liquid wettability based on molecular hydrophilic-lipophilic balance. *Chem Sci.* 2021; 12:6136-6142.
36. El-Genk MS, Saber HH. Minimum thickness of a flowing down liquid film on a vertical surface. *Int J Heat Mass Transfer.* 2001; 44:2809-2825.
37. Bertoli D, Mark EB, Liao D *et al.*, A novel MRI-based three-dimensional model of stomach volume,

- surface area, and geometry in response to gastric filling and emptying. *Neurogastroenterol Motil.* 2023;35: e14497,1-15.
38. Sharma H, Giriprasad R, Goswami M. Animal fat processing and its quality control. *J Food Process Technol.* 2013; 4:1-5.
 39. Hagedorn O, Fleute-Schlachter I, Mainx HG *et al.*, Surfactant-induced enhancement of droplet adhesion in superhydrophobic soybean (*Glycine max* L.) leaves. *Beilstein J Nanotechnol.* 2017; 8:2345-2356.
 40. Food standards Australia and New Zealand. Australian food composition database - Release 2.0, 2022.
 41. Bergfreund J, Bertsch P, Fischer P. Adsorption of proteins to fluid interfaces: role of the hydrophobic subphase. *J Colloid Interface Sci.* 2021; 584:411-417.
 42. Gadhave A. Determination of hydrophilic-lipophilic balance value. *Int J Sci Res.* 2014; 3:573-578.