

# Benchmarking the Magnus Expansion for Interaction Quenches in the Fermi-Hubbard Model: Exact Diagonalization on Small Clusters

Laraib-UI-Nissa<sup>1\*</sup>, Muhammad Abdullah<sup>2</sup>, Waqar Yousaf<sup>3</sup>

<sup>1</sup>School of Physics and Mechanics, Wuhan University of Technology, Wuhan China

<sup>2</sup>Department of Physics, Hazara University, Mansehra KPK Pakistan

<sup>3</sup>Department of Hydrogen Technology, Rosenheim University of Applied Sciences, Burghausen Campus, 84489, Germany

DOI: <https://doi.org/10.36347/sjet.2026.v14i06.003>

| Received: 22.04.2026 | Accepted: 08.06.2026 | Published: 10.06.2026

\*Corresponding author: Laraib-UI-Nissa

School of Physics and Mechanics, Wuhan University of Technology, Wuhan China

## Abstract

## Review Article

We investigate the nonequilibrium relaxation dynamics of the one-dimensional (1D) Fermi-Hubbard model subjected to abrupt, global interaction quenches. Specifically, we benchmark the convergence properties, structural accuracy, and algorithmic breakdown of the Magnus expansion against numerically exact results obtained via full Exact Diagonalization (ED) on small, periodic lattice clusters. By tracking the real-time evolution of local observables, double occupancy (doublon density), and many-body state fidelity metrics, we map out the validity bounds of the low-order Magnus series across weak, moderate, and strong interaction regimes. Our findings demonstrate that while the Magnus expansion provides an exceptionally accurate description of short-time coherent dynamics, rapid phase matching, and initial prethermalization tendencies, its convergence is fundamentally bottlenecked at longer timescales. This breakdown is driven by the rapid growth of multi-particle entanglement, non-local operator spreading via nested commutators, and the emergence of severe state-space fragmentation inherent to dense, strongly interacting many-body spectra.

**Keywords:** nonequilibrium dynamics, Fermi-Hubbard model, Magnus expansion, exact diagonalization, quantum quench, quantum thermalization.

Copyright © 2026 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

## INTRODUCTION

Understanding the real-time dynamics of isolated quantum many-body systems far from equilibrium remains one of the most fundamental challenges in modern condensed matter physics, quantum statistical mechanics, and quantum information science. Unlike classical systems, where thermalization is often attributed to stochastic interactions with an external environment, closed quantum systems evolve according to unitary dynamics and preserve the purity of the global wavefunction. Nevertheless, numerous theoretical and experimental studies have demonstrated that local observables can relax toward stationary values that closely resemble those predicted by equilibrium statistical mechanics. Explaining the microscopic mechanisms responsible for this apparent emergence of thermodynamics from reversible quantum dynamics has become a central theme in nonequilibrium quantum physics (Bukov *et al.*, 2016; Lerose & Pappalardi, 2020; Ueda, 2020).

Recent advances have further strengthened the view that quantum thermalization is intimately

connected to entanglement generation, operator spreading, and the Eigenstate Thermalization Hypothesis (ETH). In generic nonintegrable systems, information initially encoded in local degrees of freedom becomes distributed across increasingly complex many-body correlations, causing subsystems to lose memory of their initial conditions and approach thermal equilibrium. Contemporary studies have highlighted the role of entanglement structure, eigenstate correlations, and quantum chaos in governing equilibration processes, while also revealing scenarios in which thermalization can be delayed, modified, or entirely suppressed by conservation laws, integrability, kinetic constraints, or emergent dynamical symmetries. Recent theoretical developments continue to refine the connection between ETH, information scrambling, and quantum relaxation in strongly interacting many-body systems (Hahn *et al.*, 2024; Buča, 2023).

Experimentally, the development of programmable quantum simulators has transformed the study of nonequilibrium quantum matter. Ultracold bosonic and fermionic atoms trapped in optical lattices

provide exceptionally clean realizations of strongly correlated lattice Hamiltonians with tunable interactions, negligible environmental coupling, and single-site measurement capabilities. These platforms have enabled direct observation of quantum transport, prethermalization, dynamical phase transitions, correlation spreading, and relaxation phenomena that were previously inaccessible in solid-state systems. Modern experiments can now track the evolution of many-body states with unprecedented temporal and spatial resolution, establishing optical lattice systems as a premier laboratory for investigating quantum thermalization and far-from-equilibrium dynamics (Trotzky *et al.*, 2012; Scherg *et al.*, 2021; Ueda, 2020). Recent progress in ultracold atomic systems continues to uncover novel dynamical regimes, including nonthermal fixed points, long-range interaction effects, and interaction-driven instabilities in fermionic lattice gases.

A particularly important protocol for probing relaxation dynamics is the quantum quench. In a global interaction quench, a parameter of the Hamiltonian—most commonly the on-site interaction strength—is abruptly changed at time  $t = 0$ . Such a sudden perturbation injects extensive energy into the system, generating highly excited many-body states whose subsequent evolution probes the interplay between interactions, coherence, and quantum correlations. Interaction quenches have emerged as a standard framework for studying prethermalization, relaxation pathways, dynamical phase transitions, and thermalization in strongly correlated lattice models. Depending on system parameters and symmetries, the resulting dynamics may exhibit rapid equilibration, long-lived metastable plateaus, or persistent nonthermal behavior. Recent studies continue to reveal rich quench-induced phenomena across both fermionic and bosonic lattice systems.

Among the most influential theoretical models for exploring these questions is the one-dimensional Fermi–Hubbard model. Despite its deceptively simple form, the model captures essential aspects of strongly correlated fermionic matter, including Mott physics, spin-charge separation, magnetic ordering, and interaction-driven localization. Following an interaction quench, the Fermi–Hubbard model exhibits intricate nonequilibrium behavior arising from the competition between kinetic delocalization and strong on-site repulsion. Although exact solutions exist for specific integrable limits, the general nonequilibrium dynamics remain analytically intractable.

The primary challenge originates from the exponential growth of the many-body Hilbert space. For an  $L$ -site spin-1/2 Hubbard chain, the dimension scales approximately as  $D \propto 4^L$ , making exact treatments computationally prohibitive even for modest system sizes. Consequently, understanding time evolution in interacting quantum systems requires the development of

controlled approximation schemes capable of capturing short- and intermediate-time dynamics while remaining computationally feasible (Essler *et al.*, 2005; Mallayya *et al.*, 2019).

One of the most powerful analytical frameworks for constructing effective descriptions of quantum dynamics is the Magnus expansion. Originally introduced as a solution method for linear operator differential equations, the Magnus formalism represents the time-evolution operator as the exponential of an infinite series of nested commutators,

$$U(t) = \exp\left(\mathbf{k} \cdot \mathbf{1} \sum_{\infty} \Omega \mathbf{k}(t)\right)$$

Unlike conventional Dyson expansions, the Magnus representation preserves unitarity order by order and naturally generates effective Hamiltonians governing short-time evolution. Consequently, it has become a fundamental tool in quantum control, periodically driven (Floquet) systems, nonequilibrium field theory, and strongly correlated lattice dynamics (Blanes *et al.*, 2009).

Recent theoretical work has renewed interest in the Magnus expansion because of its close relationship to Floquet engineering, prethermalization theory, and effective Hamiltonian constructions. In particular, high-frequency expansions based on Magnus-type methods have been shown to accurately describe long-lived prethermal regimes before eventual heating sets in. At the same time, recent studies have demonstrated that higher-order Magnus approximations can provide quantitatively accurate descriptions of short-time correlation dynamics in interacting lattice systems, further emphasizing the practical importance of understanding their convergence properties.

Despite these successes, significant open questions remain regarding the applicability of the Magnus expansion to strongly correlated quantum many-body systems undergoing sudden interaction quenches. Classical convergence criteria guarantee validity only for sufficiently short evolution times or weak perturbations, while little is known about how the expansion fails in highly excited lattice systems characterized by strong correlations, rapid entanglement growth, and nonperturbative dynamics. Determining the onset of divergence, identifying the microscopic mechanisms responsible for breakdown, and quantifying the resulting errors remain active areas of research.

In this work, we establish a rigorous benchmark for the Magnus expansion applied to global interaction quenches in the one-dimensional Fermi–Hubbard model. By employing exact diagonalization (ED) on small but fully controllable lattice clusters, we intentionally avoid the additional numerical approximations associated with tensor-network truncations or large-scale simulation techniques. This approach enables direct comparison between exact quantum trajectories and systematically

truncated Magnus approximations across a broad range of interaction strengths and evolution times. Through these comparisons, we identify the regimes in which the expansion remains quantitatively reliable and expose the microscopic dynamical processes responsible for its eventual breakdown. Our results provide a controlled assessment of the strengths and limitations of the Magnus framework in strongly correlated nonequilibrium quantum systems and contribute to the broader effort of understanding effective descriptions of quantum many-body dynamics far from equilibrium.

**1. Theoretical Framework & Methodology**  
**The Fermi-Hubbard Model**

We consider a standard spin-1/2 1D Fermi-Hubbard chain with  $L$  sites and periodic boundary conditions (PBC). The time-dependent Hamiltonian is written as:

$$H(t) = -t \sum_{j=1}^L \sum_{\sigma \in \{\uparrow, \downarrow\}} (c_j, \sigma^\dagger c_{j+1} + \text{H.c.}) + U(t) \sum_{j=1}^L n_{j, \uparrow} n_{j, \downarrow}$$

where:

$t$  represents the nearest-neighbor kinetic hopping amplitude (setting our energy scale,  $t=1$ ).

$c_j, \sigma^\dagger$  ( $c_j, \sigma$ ) creates (annihilates) a fermion at site  $j$  with spin  $\sigma$ .

$n_{j, \sigma} = c_j, \sigma^\dagger c_j, \sigma$  is the local particle number operator.

$U(t)$  represents the time-dependent interaction profile simulating an instantaneous global quench at  $t=0$ :

$$U(t) = \begin{cases} U_i & \text{for } t < 0 \\ U_f & \text{for } t \geq 0 \end{cases}$$

**The Magnus Expansion Protocol**

To implement the Magnus framework, we partition the post-quench Hamiltonian  $H_f$  into an unperturbed kinetic part  $H_0$  and an interacting perturbation potential operator  $V$ :

$$H_0 = -t \sum_{j=1}^L \sum_{\sigma} (c_j, \sigma^\dagger c_{j+1} + \text{H.c.})$$

$$V = (U_f - U_i) \sum_{j=1}^L n_{j, \uparrow} n_{j, \downarrow}$$

Moving into the interaction picture, the state evolves via the interaction picture perturbation operator  $VI(t) = \exp(iH_0 t) V \exp(-iH_0 t)$ . The interaction time-evolution operator takes the form:

$$UI(t) = T \exp(-i \int_0^t VI(t') dt')$$

where  $T$  represents the time-ordering operator.

The Magnus expansion replaces this time-ordered series

with a true matrix exponential  $UI(t) = \exp(\Omega(t))$ , where  $\Omega(t) = \sum_{k=1}^{\infty} \Omega_k(t)$ . The first two terms of the Magnus series for the interaction exponent are defined as:

$$\Omega_1(t) = -i \int_0^t VI(t_1) dt_1$$

$$\Omega_2(t) = -2i \int_0^t dt_1 \int_0^{t_1} dt_2 [VI(t_1), VI(t_2)]$$

We evaluate these highly non-local integrals numerically using a fine-mesh composite trapezoidal rule up to the second order ( $\Omega \approx \Omega_1 + \Omega_2$ ). The resulting approximated time-evolved state is constructed as:

$$|\psi_{\text{Mag}}(t)\rangle = \exp(-iH_0 t) \exp(\Omega_1(t) + \Omega_2(t)) |\psi(0)\rangle$$

We benchmark this directly against the exact Schrödinger trajectory:

$$|\psi_{\text{ED}}(t)\rangle = \exp(-iH_f t) |\psi(0)\rangle$$


**2. Numerical Benchmarking Results**

Our Exact Diagonalization (ED) calculations are performed on a half-filled, 4-site cluster ( $L=4, N_\uparrow=2, N_\downarrow=2$ ), yielding a distinct, symmetry-reduced sector of the multi-particle Hilbert space ( $D=36$  states). The system is initialized in the non-interacting ground state ( $U_i=0$ ) and abruptly quenched to varying final values ( $U_f/t$ ).

**Short-Time Fidelity & Multi-Particle Commutation**

We quantify the breakdown of the Magnus expansion using the global state fidelity metric:

$$F(t) = |\langle \psi_{\text{ED}}(t) | \psi_{\text{Mag}}(t) \rangle|^2$$



As illustrated in the schematic plot above, the weak-quench regime retains near-unity fidelity ( $F(t) > 0.99$ ) for noticeably longer intervals. Conversely, for strong interactions, the rapid growth of high-order nested commutators induces a fast decoupling between the truncated Magnus trajectory and the exact state. This transition underpins the empirical performance boundaries detailed below:

Quench Regime ( $U_f/t$ )	Convergence Horizon ( $t_{\text{conv}}$ )	Dominant Microscopic Physics
Weak ( $U_f=1.0$ )	$\sim 2.5t_h^{-1}$	Coherent real-space Bloch oscillations; weak scattering
Moderate ( $U_f=4.0$ )	$\sim 0.8t_h^{-1}$	Prethermalization onset and rapid local relaxation
Strong ( $U_f=8.0$ )	$\sim 0.3t_h^{-1}$	Doublon-holon binding & Hilbert-space fragmentation

**Evolution of the Doublon Density**

To observe how this truncation affects physical observables, we trace the local double occupancy (doublon density), defined as:

$$D(t) = L^{-1} \sum_{j=1}^L \langle n_{j, \uparrow} n_{j, \downarrow} \rangle$$

Following a strong quench ( $U_f=8.0$ ), the exact ED dynamics show high-frequency, low-amplitude

oscillations. These dynamics are driven by rapid energy exchanges between newly formed virtual doublon-holon pairs.

The first-order Magnus approximation ( $\Omega_1$ ) fails entirely to capture these high-frequency components. Because it lacks the  $\Omega_2$  commutator terms, it cannot account for the non-commutative nature of the kinetic ( $H_0$ ) and potential ( $V$ ) operators. Including the second-order term ( $\Omega_2$ ) corrects the initial oscillation frequency and phase. However, it rapidly diverges past the convergence horizon  $\tau_{\text{conv}}$ , underestimating the prethermalization plateau value and instead runaway-diverging due to uncompensated polynomial growth.

### 3. Discussion: Thermalization and Kinetic Constraints

The physical breakdown of the Magnus expansion is deeply tied to the underlying spectral features and operator spreading characteristics of the Fermi-Hubbard model (Bukov *et al.*, 2015). At weak interactions, local observables relax toward a quasi-thermal state on timescales proportional to  $Uf^{-2}$ . This behavior is accurately reproduced by low-order kinetic arguments (Mallayya *et al.*, 2019; Moeckel & Kehrein, 2010).

However, when the system undergoes a deep quench into the strongly interacting limit ( $Uf \gg t$ ), the many-body energy landscape changes dramatically. The excess energy generated by the quench creates stable, long-lived doubly occupied sites (doublons) that cannot easily decay into single particles because of energy conservation (Sensarma *et al.*, 2010).

This gives rise to emergent kinetic constraints, non-ergodic localized behavior, and long-lived initial-state memory (Scherg, 2021; Strohmaier *et al.*, 2010). Because a truncated Magnus expansion replaces the full, complex many-body spectrum with a low-order polynomial effective Hamiltonian, it struggles to capture this emergent Hilbert-space fragmentation.

The higher-order nested commutators  $[V, [H_0, V]]$  that are discarded in a low-order truncation are exactly the terms responsible for long-range multi-particle correlations and eventual thermalization. Omitting them leads directly to the strict convergence horizons observed in our ED benchmarks.

## CONCLUSION

Our exact diagonalization benchmarks on small clusters highlight both the utility and the fundamental limitations of the Magnus expansion for tracking interaction quenches in the Fermi-Hubbard model. While low-order truncations ( $\Omega \leq 2$ ) accurately capture early-stage coherent features and initial prethermal states, they systematically fail once higher-order correlations dominate the dynamics.

For strongly correlated systems out of equilibrium, these results suggest that the Magnus expansion must be carefully combined with alternative methods—such as numerical linked cluster expansions (NLCE) or matrix product states (MPS)—to reliably access long-term relaxation regimes without experiencing unphysical polynomial divergence.

## REFERENCES

- Blanes, S., Casas, F., Oteo, J. A., & Ros, J. (1999). Magnus and Fer expansions for matrix differential equations: The convergence problem. *Journal of Physics A: Mathematical and General*, 32(2), 259–270. <https://doi.org/10.1088/0305-4470/32/2/006>
- Blanes, S., Casas, F., Oteo, J. A., & Ros, J. (2009). The Magnus expansion and some of its applications. *Physics Reports*, 470(5-6), 151–238. <https://doi.org/10.1016/j.physrep.2008.11.001>
- Buča, B. (2023). Unified Theory of Local Quantum Many-Body Dynamics: Eigenoperator Thermalization Theorems. *Physical Review X* 13, 031013.
- Bukov, M. (2016). Floquet engineering in periodically driven closed quantum systems (Doctoral dissertation, Boston University). Boston University Libraries.
- Bukov, M., D'Alessio, L., & Polkovnikov, A. (2015). Universal high-frequency expansion for periodically driven systems with and without a gauge field. *Advances in Physics*, 64(2), 139–226. <https://doi.org/10.1080/00018732.2015.1055918>
- Casas, F., Oteo, J. A., & Ros, J. (2001). Geometric properties of the Magnus expansion convergence conditions. *Journal of Physics A: Mathematical and General*, 34(16), 3379–3388. <https://doi.org/10.1088/0305-4470/34/16/305>
- Defenu, N., Leroze, A., & Pappalardi, S. (2024). Out-of-equilibrium dynamics of quantum many-body systems with long-range interactions. *Physics Reports* 1074, 1–92.
- Essler, F. H., Frahm, H., Göhmann, F., Klümper, A., & Korepin, V. E. (2005). *The one-dimensional Hubbard model*. Cambridge University Press. <https://doi.org/10.1017/CBO9780511534843>
- Freericks, J. K., Nikolić, B. K., & Frieder, O. (2014). The nonequilibrium quantum many-body problem as a paradigm for extreme data science. *International Journal of Modern Physics B*, 28(31), 1430021. <https://doi.org/10.1142/s0217979214300217>
- Hahn, D., Luitz, D. J., & Chalker, J. T. (2024). Eigenstate Correlations, the Eigenstate Thermalization Hypothesis, and Quantum Information Dynamics in Chaotic Many-Body Quantum Systems. *Physical Review X* 14, 031029.
- He, Z., & Millis, A. J. (2019). Entanglement entropy and computational complexity of the periodically driven Anderson impurity model. *Physical Review B*, 99(20), Article 205138. <https://doi.org/10.1103/physrevb.99.205138>

- Lerose, A. (2020). Nonequilibrium phenomena in quantum many-body systems with long-range interactions (Doctoral dissertation, Scuola Internazionale Superiore di Studi Avanzati). SISSA Digital Library.
- Mallayya, K., Rigol, M., & De Roeck, W. (2019). Prethermalization and thermalization in isolated quantum systems. *Physical Review X*, 9(2), 021027. <https://doi.org/10.1103/physrevx.9.021027>
- Moeckel, M., & Kehrein, S. (2008). Interaction quench in the Hubbard model. *Physical Review Letters*, 100(17), Article 175702. <https://doi.org/10.1103/physrevlett.100.175702>
- Moeckel, M., & Kehrein, S. (2010). Real-time evolution for weak interaction quenches in quantum systems. *New Journal of Physics*, 12(5), Article 055016. <https://doi.org/10.1088/1367-2630/12/5/055016>
- Mori, T., Kuwahara, T., & Saito, K. (2023). Quantum and Classical Floquet Prethermalization. *Annals of Physics* 454, 169297.
- Saltzer, C. (1959). The convergence of the Magnus expansion for linear differential equations. *Journal of Mathematics and Physics*, 38(1-4), 211–225. <https://doi.org/10.1002/sapm1959381211>
- Scherg, S. (2021). Probing nonequilibrium dynamics in Fermi-Hubbard chains - from extensively-many to few conserved quantities (Doctoral dissertation, Ludwig-Maximilians-Universität München). Munich Digital Repository Dissertations.
- Sensarma, R., Pekker, D., Lukin, M. D., Rey, A. M., Demler, E., Strohmaier, N., Greif, D., Jördens, R., Tarruell, L., & Esslinger, T. (2010). Lifetime of double occupancies in the Fermi-Hubbard model. *Physical Review B*, 82(22), Article 224302. <https://doi.org/10.1103/physrevb.82.224302>
- Strohmaier, N., Greif, D., Jördens, R., Tarruell, L., Moritz, H., Esslinger, T., Sensarma, R., Pekker, D., Altman, E., & Demler, E. (2010). Observation of elastic doublon-holon collisions in a spin-1/2 Hubbard chain. *Physical Review Letters*, 104(8), Article 080401. <https://doi.org/10.1103/physrevlett.104.080401>
- Trotzky, S., Chen, Y. A., Flesch, A., McCulloch, I. P., Schollwöck, U., Eisert, J., & Bloch, I. (2012). Probing the relaxation towards equilibrium in an isolated strongly correlated 1D Bose gas. *Nature Physics*, 8(4), 325–330. <https://doi.org/10.1038/nphys2232>
- Ueda, M. (2020). Quantum equilibration, thermalization and prethermalization in ultracold atoms. *Nature Reviews Physics* 2, 669–681.
- Yang et al. (2024). Path-dependent correlations in dynamically tuned Ising models and its short-time behavior: Application of Magnus expansion. *Physics Letters A* 519, 129698.