



Comparative *in-Vitro* Antimicrobial Study of Ethanolic Seed Extract of *Syzygium cumini* (L.) and Methanolic Leaves Extract *Tecomaria capensis*

Prasanta Dey^{1*}, Partha Chakraborty², Tejendra Bhakta³

¹Dept. of Pharmaceutical Chemistry, C. L. Baid Metha College of Pharmacy, Thoraipakkam, Chennai-97, India

²Dept. of Microbiology, Tripura Institute of Paramedical Science, Agartala, Tripura (W), 799014, India

³Regional Institute of Pharmaceutical Science & Technology, Agartala, Tripura (W), 799005, India

Research Article

Article History

Received: 06 October, 2012

Accepted: 18 October, 2012

Published: 25 November, 2012

Corresponding author

Prasanta Dey

Email: deypasantadey@yahoo.com

Mob: +91-7418380592

Abstract – The comparative *in-vitro* anti-bacterial activity was carried out between ethanolic seed extract of *Syzygium cumini* (L.) and methanolic leaves extract *Tecomaria capensis* against *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Enterococcus faecalis* using different concentration of 2, 4, 6 and 8 mg/ml and STD ciprofloxacin 5 mcg. Between these two plants extract the methanolic leaves extract *Tecomaria capensis* shows more anti-bacterial activity than the other.

Keywords – *Syzygium cumini*, *Tecomaria capensis*, anti-bacterial activity

Introduction

Medicinal plants can able to protect from different deadly microbial species. These plants are used in different forms under indigenous systems of medicine like Ayurveda, Sidha and Unani [1]. Rapid and overdose of synthetic drugs can cause resistance development in the body [2]. So, in comparison to synthetic drugs herbal medicines are safer [3]. According to WHO, 80% of the world's inhabitants problem should be treated by medicinal herbal drug for their primary health care [1].

Syzygium cumini (L.) belongs to family Myrtaceae [3, 4, 5]. It is commonly known as Jamun (Hindi) [3], Naaval (Tamil) [5]. The seeds of *S. cumini* are widely considered to have anti-diabetic properties [3, 4]. The fruits [6] and leaves [7] of *S. cumini* have unique anti-oxidant activity. The seeds have significant anti-inflammatory [8], anticancerous [9] and central nervous system [5] activities.

Tecomaria capensis belongs to family bignoniaceae also known as Cape-honeysuckle [10] is a fast growing, scrambling shrub which may grow up to 2-3m high and spread more than 2.5m. Plant is used traditionally to relieve pain and sleeplessness [11]. Dried powdered bark infusions are taken for sleeplessness [12], reported to induce sleep [13] and also as anti-plasmodial activity against *Plasmodium falciparum* [14].

Materials and Methods

Plant material

The leaf of *tecomaria capensis* and seed of *syzygium cumini* are collected from various parts of Tripura. These parts are dried, powdered and extracted by methanol and ethanol respectively. The extract was collected, evaporated and kept in desiccators.

Drugs and chemical

The standard drug ciprofloxacin was used for comparison. Methanol and ethanol used in the experiment were of analytical grade.

In-vitro antibacterial activity (Well-diffusion method)

The antibacterial activities of ethanolic seed extract of *Syzygium cumini* (L.) and methanolic leaves extract *Tecomaria capensis* was carried out against different gram positive and gram negative bacteria were carried out by Agar well diffusion method on nutrient agar plates [15]. Test organisms used were *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Enterococcus faecalis*. Different concentrations (2 mg/ml- 8 mg/ml) of seed and leaves extract were made and the zone of inhibition was calculated for each concentration. Ciprofloxacin (5mcg) were used as the standards.

Results and Discussion

Four varieties of bacteria were tested for their sensitivity to ethanolic seed extract of *Syzygium cumini* (L.) and methanolic leaves extract *Tecomaria capensis*, antibacterial potency was determined by the well diffusion method. Table 1, 2 and 3 presents diameters of inhibition zones (clear zones around wells) exerted by the various concentrations of ethanolic seed extract of *Syzygium cumini* (L.), methanolic leaves extract *Tecomaria capensis* and standard ciprofloxacin. Between these two plants extract the methanolic leaves extract *Tecomaria capensis* shows more anti-bacterial activity than the other.

Both the plants contain phenolic compounds. Most of the studies on the mechanism of phenolic compounds focused on their effects on cellular membranes. Actually, phenolics not only attacked cell wall and cell membrane, thereby destroying its permeability and releasing of intracellular constituents (ribose, Na glutamate, etc.) but also interfered with membrane function e.g. electron transport, nutrient uptake, protein and nucleic acid synthesis and enzyme activity. That was, active compounds might have several invasive targets which could lead to inhibit the bacteria. Furthermore, leakage of intracellular material was a general phenomenon induced by many antibacterial substances.

Table 1: Antibacterial activity of ethanolic seed extract of *S. cumini*

Sl. No	Bacteria	Zone of inhibition in mm			
		2 mg/ml	4 mg/ml	6 mg/ml	8 mg/ml
1	<i>Escherichia coli</i>	8.1 ± 0.1	8.2 ± 0.1	8.4 ± 0.1	8.5 ± 0.09
2	<i>Staphylococcus aureus</i>	5.2 ± 0.7	5.4 ± 0.10	5.7 ± 0.12	5.9 ± 0.07
3	<i>Pseudomonas aeruginosa</i>	8.1 ± 0.1	8.2 ± 0.1	8.3 ± 0.11	8.6 ± 0.11
4	<i>Enterococcus faecalis</i>	6.2 ± 0.10	6.4 ± 0.08	6.6 ± 0.07	7 ± 0.28

Table 2: Antibacterial activity of methanolic leaves extract *tecomaria capensis*

Sl. No	Bacteria	Zone of inhibition in mm			
		2 mg/ml	4 mg/ml	6 mg/ml	8 mg/ml
1	<i>Escherichia coli</i>	15.8 ± 0.1	19.6 ± 0.1	35.5 ± 0.1	37.7 ± 0.1
2	<i>Staphylococcus aureus</i>	13.4 ± 0.1	18 ± 1.22	34.8 ± 0.06	37.6 ± 0.08
3	<i>Pseudomonas aeruginosa</i>	11.6 ± 0.08	15.9 ± 0.12	33.5 ± 0.09	34.7 ± 0.11
4	<i>Enterococcus faecalis</i>	14.5 ± 0.1	20.8 ± 0.06	33.5 ± 0.24	36.6 ± 0.15

Table 3: Antibacterial activity of standard antibiotics

Bacteria	Zone of inhibition in mm
	Ciprofloxacin (5mcg)
<i>Escherichia coli</i>	35.6 ± 0.27
<i>Staphylococcus aureus</i>	30.6 ± 0.18
<i>Pseudomonas aeruginosa</i>	30.5 ± 0.41
<i>Enterococcus faecalis</i>	39.3 ± 0.38

Conclusion

Above experiment shows the anti-bacterial activity of both the plant extracts. But methanolic leaves extract *tecomaria capensis* shows more anti-bacterial activity than the other. So, *tecomaria capensis* is a good anti-microbial agent.

References

- NK, Kumar R and Tripathi P, Global promotion of herbal medicine: India's opportunity, Current Science, 2004, 86: 37-41.
- Zampini IC, Cuello S, Alberto MR, Ordonez RM, Almeida RD, Solorzano E, Isla MI, Antimicrobial activity of selected plant species from the Argentine puna against sensitive and multiresistant bacteria, Journal of Ethnopharmacology, 2009, 124: 499-505.
- Modi DC, Patel JK, Shah BN and Nayak BS, Pharmacognostic Studies of the Seed of *Syzygium cumini* Linn., Pharma Science Monitor, 2010, 1: 20-26.
- Gowri SS and Vasantha K, Free Radical Scavenging and Antioxidant Activity of Leaves from Agathi (*Sesbania grandiflora*) (L.) Pers., American-Eurasian Journal of Scientific Research, 2010, 5: 114-119.
- Kumar A, Padmanabhan N and Krishan MRV, Central Nervous System Activity of *Syzygium cumini* Seed., Pakistan Journal of Nutrition, 2007, 6: 698-700.
- Zhang LL and Lin YM, Antioxidant tannins from *Syzygium cumini* fruit, African Journal of Biotechnology, 2009, 8: 2301-2309.
- Ruan ZP, Zhang LL and Lin YM, Evaluation of the Antioxidant Activity of *Syzygium cumini* leaves, Molecules, 2008, 13: 2545-2556.
- Kumar A, Ilavarasan R, Jayachandran T, Decaraman M, Kumar RM, Aravindhan P, Padmanabhan N and Krishan MRV, Anti-inflammatory activity of *Syzygium cumini* seed, African Journal of Biotechnology, 2008, 7: 941-943.

9. Parmar J, Sharma P, Verma P, Sharma P and Goyal PK, Chemopreventive Action of *Syzygium cumini* on DMBA -induced Skin Papillomagenesis in Mice, Asian Pacific Journal of Cancer Prevention, 2010, 11: 261-265.
10. USDA, ARS, National Genetic Resources Program, *Germplasm Resources Information Network* - (GRIN) [Online Database], National Germplasm Resources Laboratory, Beltsville, Maryland.
11. South African National Biodiversity Institute's (SANBI) Integrated Biodiversity System, information.
12. Roberts, M; *Indigenous Healing Plants*, Southern Book Publishers, Halfway House, 1990.
13. Hutchings A., Scott, A.H., Lewis, G., Cunningham, A.B., *Zulu Medicinal Plants: An Inventory*. University of Natal Press, Pietermaritzburg, 1996.
14. P. Pillaya, V.J. Maharaj, P.J. Smith, Investigating South African plants as a source of new antimalarial drugs, *Journal of Ethnopharmacology*, 2008, 119: 438-454.
15. Clinical and Laboratory Standards Institute, *Performance Standards for Antimicrobial Susceptibility Testing. Fifteenth Information Supplement M100-S15*, Wayne, PA, USA, 2005.