

Fishing Activity on Solomougou Dam Lake (Northern Côte d'Ivoire): Synthesis of Recent Data and Additional Information

Moustapha DIABY^{1*}, Jean-Paul Aka AGNISSAN², Koffi Dongo KOUASSI³, Konan N'DA⁴¹Maître-Assistant, Institut de Gestion Agropastorale, Université Peleforo GON COULIBALY, Côte d'Ivoire²Attaché de Recherche, Unité de Formation et de Recherches des Sciences Biologiques, Université Peleforo GON COULIBALY, Côte d'Ivoire³Maître de Conférences, Unité de Formation et de Recherches des Sciences Biologiques, Université Peleforo GON COULIBALY, Côte d'Ivoire⁴Professeur titulaire, Unité de Formation et de Recherches des Sciences de la Nature, Université Nangui ABROGOUA, Côte d'IvoireDOI: [10.36347/sajb.2022.v10i06.001](https://doi.org/10.36347/sajb.2022.v10i06.001)

| Received: 18.04.2022 | Accepted: 26.05.2022 | Published: 09.06.2022

*Corresponding author: Moustapha DIABY

Maître-Assistant, Institut de Gestion Agropastorale, Université Peleforo GON COULIBALY, Côte d'Ivoire

Abstract

Review Article

In Côte d'Ivoire, like other continental waters, many dam lakes in the North are exploited for fishing purposes. This activity provides local communities fish for consumption and income through the economic activities it generates. Therefore, sustainable management of these fisheries is necessary. For this purpose, fishing on the Solomougou Dam Lake was investigated in 2018 in order to analyse the current state of exploited fish resources. The investigation provided data on production factors with reference to fishermen's sociological profiles and the type of fishing gears used, on the ichthyological composition of catches and on stocks exploitation parameters of abundant species caught. This paper is a review of those previous studies. It is completed on the one hand using information on conservation and the vulnerability status of the whole species and on the other hand using stocks exploitation parameters of a second group of species secondarily abundant in catches.

Keywords: Fishing activity, fish stocks, Solomougou Dam Lake, Côte d'Ivoire.

Copyright © 2022 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

INTRODUCTION

Côte d'Ivoire has various aquatic environments. The Southern part of the country is bordered by the Atlantic Ocean with a continental shelf covering 12,000 km². There are also lagoons with a total area of 1,200 km² (Golé Bi *et al.*, 2005). The country also has four artificial lakes that serve as hydroelectric dams of 1,700 km² including rivers (FAO, 2005). In the North, the body of water includes many dam lakes, estimated at 210. They have been created since the 1970s to develop agriculture and livestock breeding (Le Guen, 2002), and some of them are now exploited for fishing.

Fishing in these dam lakes is of great importance for local communities. It provides them with fish, one of the main animal protein sources in their diet. Indeed, although the Northern region is known as a cattle-breeding area in the country, it has been noted that the price of beef is constantly rising in recent years. Therefore, fishing activities should be monitored for perpetuating fish supply to many economically weaker populations in this region.

The Solomougou Dam Lake is one of the largest fisheries in northern Côte d'Ivoire (Figure 1). Located between 09° 18' 09" N latitude and 05° 37' 36" W longitude, it was created in 1973 for rice cultivation (Cecchi *et al.*, 2007). The water surface is ranked fifth in the country with an area of 500 hectares, Morisson Dam Lake being the largest with its 3,700 hectares (Traoré, 1996). During the last three years, fishing activities in this reservoir have been the subject of several studies. These include the fishermen's sociological profile as well as fishing gears and fishing techniques used (Diaby *et al.*, 2020a), the dynamics of the fishing activity throughout the year (Kouassi *et al.*, 2019), the fish composition of catches (Kouassi *et al.*, 2020a), the biological parameters related to the length-weight relationship of fish species (Kouassi *et al.*, 2020b) or the stocks exploitation parameters of abundantly species caught (Diaby *et al.*, 2020b; Diaby *et al.*, 2020c).

This work is a review of those previous studies. It also provides additional information on the conservation status of the fish species present in this

fishery. The latter data have been obtained from the IUCN list (IUCN, 2021) and the Fishbase site (Froese and Pauly, 2021). Species vulnerability levels were retrieved from the Froese and Pauly (2021) database. Brodie (2010) defines vulnerability as sensitivity of organisms to impacts caused by fishing activities or other factors that threaten their existence, such as habitat loss. Additional information also concerns stocks

exploitation parameters of a second group of species secondarily abundant in catches. The methods used are identical to those used by Diaby *et al.* (2020b) and Diaby *et al.* (2020c). Indeed, taking these species into account in fisheries management must also be a priority. They are captured for consumption, and they contribute to the biological diversity, and consequently to the proper functioning of these ecosystems.




Fig-1: Geographical location of the Solomougou Dam Lake in Northern Côte d'Ivoire

FISHERMEN'S SOCIOLOGICAL PROFILE

In total, forty-six (46) fishermen have been registered in this fishery from October 2018 to September 2019. Twenty-six (26) of them were working individually and the other twenty (20) organized in seven (7) teams including two (2) teams of two (2) fishermen each, four (4) of three (3) fishermen each and only one (1) of four (4) fishermen. So, this represents a total of thirty-three (33) fishing units, fishing unit representing individual fisherman or fishermen in association operating one fishing gear. Fishermen's sociological profile indicated that 95.65% of them are of Malian origin, while the others are Ivorians. Their ages range from 13 to 61 years old and 95.74% of them have not been to school. Regarding their activities outside fishing, except two (2) fishermen practicing trading and butchering in addition to the fishing, the others do not have alternative livelihoods (Kouassi *et al.*, 2019).

FISHING GEARS TYPES

The fishermen used six fishing gear types including sparrow hawk, longline, trap, bamboo trap,

seine net and gillnet. Sparrow hawks, constituted in monofilament, have bar length of 20 to 60 millimetres and opening diameters from 3 to 5 meters. The main line of longlines is 300 to 500 meters long, on which are attached at least 3,000 hooks spaced generally 10 centimetres apart. The traps are of an artisanal model made by weaving liana with average height of 1 meter and an opening in the upper part. As for Bamboo traps, they are made up of 1-meter-long hollowed china bamboo pieces mounted on a 500 meters long rope. Seine nets had nets with bar length of 16 millimetres, their total lengths ranging from 500 to 800 meters. Gillnets have bar length from 14 to 40 millimetres. Most of them used 30 millimetres ones (31.94%) followed by 20 millimetres (22.93%). Those with bar length smaller than 20 millimetres represent only 22.91% against 32.64% which have meshes between 20 and 30 millimetres. In addition, the lengths of these gears vary between 50 and 200 meters and the depth is from 1 to 2 meters. However, the most used have lengths from 100 to 150 meters (40.28%) and height between 1.10 and 1.50 meters (67.36%) (Diaby *et al.*, 2020a).

RELATIVE ABUNDANCES OF FISHING GEAR TYPES

Among the 321 fishing gears enumerated during the study period, traps (47.35%) and gillnets (44.86%) were the dominant ones, representing over 90% of fishing gears deployed. The least used are bamboo traps (0.31%) and seine nets (1.25%), while longlines and sparrow hawks have respective proportions of 2.80% and 3.43%. Relatively to the thirty-three (33) fishing units, 15.15% of them use more than one fishing gear type, including the association gillnet and longline (3.03%), sparrow hawk and gillnet (6.06%) and trap and sparrow hawk (6.06%). As for the others operating with only one fishing gear type, they use mostly gillnets (45.45%) and seine nets (12.12%), sparrow hawks and traps 9.09% each, longline (6.06%) and bamboo trap (3.03%) (Diaby *et al.*, 2020a).

FISHING GEARS PERIODS OF USE

Two fishing gear types were used throughout the year regardless of the change in water depth. These were gillnets and sparrow hawks. However, the first were less used from August to October during the flood period of the reservoir and the second ones from April to September. The other fishing gears were characterized by their periods of use limited to a few months. Traps were not used from April to June in the period of low water, contrary to seine nets exclusively used during this period. As for longlines and bamboo traps, used respectively during 7 and 5 months, their periods of use were disparate and do not seem to be related to variations in water depth (Kouassi *et al.*, 2019).

TEMPORAL VARIATION OF FISHING ACTIVITY

As a reminder, forty-six (46) fishermen grouped into thirty-three (33) fishing units have been identified. Fishing is practiced throughout the year. However, the site remains less frequented from August to October with only four (4) to seven (7) fishing units. From November, its colonization gradually increases to reach the peak in March with 21 units corresponding to 33 fishermen in total. Outside this period, the number of fishing units varies between eleven (11) and seventeen (17).

The length of stay of fishermen on the site varies between one (01) and twelve (12) months.

Anyway, most of the fishermen stayed for less than 6 months. In fact, 24.24% stayed for only one month while 30.30% stayed there for four months. A second group of fishermen, fewer in number, stayed longer on the site for a period of 8 months for most of them. In any case, only two fishermen remained on the site for most of the year, 11 months for one and 12 months for the other (Kouassi *et al.*, 2019).

FISHERY EXPLOITATION LEVEL

The total number of fishing units (33) and the area of the fishery (500 hectares or 5 km²) corresponds to a density of 6.6 fishing units per km². As some fishing units use more than one fishing gear type, those using longlines, bamboo traps and seine nets are at a density lower than 1 per km² each. The others using traps, sparrow hawks and gillnets are respectively at 1, 1.4 and 3.6 per km². In terms of the number of fishing gear types, bamboo traps (0.2 per km²), seine nets (0.8 per km²), longlines (1.8 per km²) and sparrow hawks (2.2 per km²) can be deployed in more places. The most numerous are gillnets (28.8 per km²) and traps (30.4 per km²) (Kouassi *et al.*, 2019).

FISH COMPOSITION, RELATIVE ABUNDANCES AND TEMPORAL DISTRIBUTION OF SPECIES

In total, fourteen (14) families, twenty-five (25) genera and thirty-seven (37) species were recorded. As for families, the most diverse are Cichlidae, Mormyridae and Cyprinidae with ten, six and five species respectively. On the other hand, the most abundant are the Cichlidae (41.87%), followed by the Alestidae (14.47%) and the Mochokidae (12.84%). At the species level, the three most representative species in proportions varying between 10% and 15% are *Brycinus imberi* (14.41%), *Synodontis schall* (12.76%) and *Oreochromis niloticus* (11.62%) (Table I).

The different species were grouped into fourteen (14) accidental, one (01) incidental and twenty-two (22) constant species. The former was encountered between one (01) and three (3) months during the year, as opposed to four (4) for the latter. As for the so-called constant species, they appeared during the year between seven (7) and twelve (12) months (Kouassi *et al.*, 2020a).

Table-I: Composition, relative abundances and temporal distribution of species: Accid., accidental; Acces., accessory and Const., constant. (Kouassi *et al.*, 2020a)

Families	Species	Number	Relatives Abundances (%)	
			Per species	Per family
	<i>Brycinus imberi</i> (Const.)	1,866	14.41	
Alestidae	<i>Brycinus nurse</i> (Accid.)	8	0.06	14.47
Anabantidae	<i>Ctenopoma patherici</i> (Const.)	27	0.21	0.21
	<i>Chromidotilapia guntheri</i> (Const.)	248	1.92	
	<i>Hemichromis bimaculatus</i> (Accid.)	6	0.05	
	<i>Hemichromis fasciatus</i> (Const.)	906	7.00	
	<i>Oreochromis niloticus</i> (Const.)	1,505	11.62	

Cichlidae	<i>Sarotherodon galilaeus</i> (Const.)	1,160	8.96	41.87
	<i>Sarotherodon melanotheron</i> (Accid.)	18	0.14	
	<i>Tilapia busumana</i> (Accid.)	2	0.02	
	<i>Tilapia guineensis</i> (Accid.)	21	0.16	
	<i>Tilapia mariae</i> (Const.)	1,115	8.61	
	<i>Tilapia zillii</i> (Const.)	441	3.41	
Clariidae	<i>Clarias gariepinus</i> (Const.)	1,036	8.00	
	<i>Heterobranchus longifilis</i> (Const.)	129	1.00	9.00
Claroteidae	<i>Chrysichthys maurus</i> (Const.)	252	1.95	
	<i>Chrysichthys nigrodigitatus</i> (Const.)	529	4.08	6.03
Cyprinidae	<i>Barbus sacratu</i> (Accid.)	8	0.06	
	<i>Barbus wurtzi</i> (Accid.)	7	0.05	
	<i>Labeo coubie</i> (Const.)	213	1.64	2.97
	<i>Labeo parvus</i> (Const.)	138	1.07	
Malapteruridae	<i>Raiamas senegalensis</i> (Const.)	18	0.14	
	<i>Malopterurus electricus</i> (Accid.)	4	0.03	0.03
Mochokidae	<i>Synodontis nigrita</i> (Accid.)	10	0.08	
	<i>Synodontis schall</i> (Const.)	1,653	12.76	12.84
Mormyridae	<i>Marcusenius furcidens</i> (Const.)	322	2.49	
	<i>Marcusenius senegalensis</i> (Accid.)	1	0.01	
	<i>Marcusenius ussheri</i> (Const.)	253	1.95	8.78
	<i>Mormyrus rume</i> (Const.)	314	2.42	
Notopteridae	<i>Petrocephalus Bovei</i> (Const.)	245	1.89	
	<i>Pollimyrus isidori</i> (Accid.)	2	0.02	
	<i>Papyrocranus afer</i> (Accid.)	7	0.05	0.05
Osteoglossidae	<i>Heterotis niloticus</i> (Const.)	29	0.22	0.22
Polypteridae	<i>Polypterus endlicheri</i> (Accid.)	5	0.04	0.04
Protopteridae	<i>Protopterus annectens</i> (Accid.)	5	0.04	0.04
Schilbeidae	<i>Schilbe intermedius</i> (Const.)	439	3.39	
	<i>Schilbe mystus</i> (Acces.)	8	0.06	3.45
Total		12,950	100.00	100.00

CONSERVATION STATUS AND VULNERABILITY OF SPECIES

In the Solomougou Dam Lake, fish species are mainly benthopelagic (43.24%) and demersal (48.65%), and secondarily pelagic (8.11%). According to the classification of the International Union for Conservation of Nature (IUCN), 97% of the lake's fish population belong to the category of Least Concern (LC). Only one species, *Marcusenius furcidens*, is Near Threatened (NT). In addition, the different species belong to seven (7) levels of vulnerability. Those with "low" and "low to moderate" vulnerability levels are the most represented, respectively 27.03% and 29.73% of the species, followed by "moderate" (18.92%) and "moderate to high" (10.81%) vulnerabilities. Species with "high" (5.41%), "high to very high" (5.41%) and "very high" (2.70%) vulnerabilities are less abundant (Table II).

EXPLOITATION PARAMETERS OF SPECIES STOCKS

In Solomougou Dam Lake, stocks exploitation parameters of the nineteen (19) fish species studied are variable. The longest asymptotic lengths (L_{∞}) are noted for *Clarias gariepinus* (47.78 cm) and *Mormyrus rume* (43.58 cm) and secondarily for *Heterobranchus*

longifilis, *Labeo coubie*, *Marcusenius furcidens*, *Marcusenius ussheri* and *Schilbe intermedius* with values between 30 cm and 40 cm. In contrast, growth rates (K) are higher for *Mormyrus rume*, *Petrocephalus Bovei*, *Schilbe intermedius*, *Hemichromis fasciatus*, *Sarotherodon galilaeus* and *Synodontis schall* (K between 7 and 10). These growth parameters provide longevities greater than five years for nine species, with those of *Chromidotilapia guntheri*, *Heterobranchus longifilis*, *Labeo parvus* and *Marcusenius ussheri* being the maximum, ranging from nine to ten years.

Mortality parameters were used to obtain exploitation rates. Three species were overexploited. These were *Chromidotilapia guntheri* (E=0.54), *Chrysichthys maurus* (E=0.78) and *Schilbe intermedius* (E=0.60). These exploitation rates, higher than reference points E_{10} , E_{50} or E_{max} , confirm this level of exploitation. Four other species, represented by *Marcusenius furcidens*, *Marcusenius ussheri*, *Mormyrus rume* and *Tilapia zillii* are close to their optimal exploitation level. Indeed, with their exploitation rates close to E=0.5 and above E_{50} , their stocks have therefore reached critical level. The stocks of the remaining twelve species are underexploited (Table III).

Table-II: Conservation status, vulnerability and habitat of fish species encountered in the period January-December 2019 in the Solomougou Dam Lake (Northern Côte d'Ivoire) (IUCN, 2021; Froese and Pauly).

Species	Conservation status	Vulnerability	Habitat
<i>Brycinus imberi</i>	Least Concern (LC)	Low vulnerability (10%)	Demersal
<i>Brycinus nurse</i>	Least Concern (LC)	Low to moderate vulnerability (29%)	Pelagic
<i>Barbus sacratus</i>	Least Concern (LC)	Moderate vulnerability (42%)	Benthopelagic
<i>Barbus wurtzi</i>	Least Concern (LC)	Moderate vulnerability (40%)	Benthopelagic
<i>Labeo coubie</i>	Least Concern (LC)	Moderate vulnerability (39%)	Benthopelagic
<i>Labeo parvus</i>	Least Concern (LC)	Moderate to high vulnerability (51%)	Benthopelagic
<i>Raiamas senegalensis</i>	Least Concern (LC)	Low to moderate vulnerability (32%)	Benthopelagic
<i>Protopterus annectens</i>	Least Concern (LC)	Moderate vulnerability (42%)	Demersal
<i>Heterotis niloticus</i>	Least Concern (LC)	Moderate to high vulnerability (55%)	Pelagic
<i>Marcusenius furcidens</i>	Near Threatened (NT)	Moderate vulnerability (35%)	Demersal
<i>Marcusenius senegalensis</i>	Least Concern (LC)	Moderate vulnerability (35%)	Demersal
<i>Marcusenius ussheri</i>	Least Concern (LC)	Low to moderate vulnerability (35%)	Demersal
<i>Mormyrus rume</i>	Least Concern (LC)	High vulnerability (63%)	Demersal
<i>Petrocephalus bovei</i>	Least Concern (LC)	Low vulnerability (13%)	Demersal
<i>Pollimyrus isidori</i>	Least Concern (LC)	Low vulnerability (11%)	Demersal
<i>Papyrocranus afer</i>	Least Concern (LC)	Moderate vulnerability (38%)	Demersal
<i>Ctenopoma petherici</i>	Least Concern (LC)	Low to moderate vulnerability (31%)	Benthopelagic
<i>Chromidotilapia guntheri</i>	Least Concern (LC)	Low vulnerability (25%)	Benthopelagic
<i>Hemichromis bimaculatus</i>	Least Concern (LC)	Low vulnerability (19%)	Benthopelagic
<i>Hemichromis fasciatus</i>	Least Concern (LC)	Low vulnerability (14%)	Benthopelagic
<i>Oreochromis niloticus</i>	Least Concern (LC)	Low to moderate vulnerability (30%)	Benthopelagic
<i>Sarotherodon galilaeus</i>	Least Concern (LC)	Low to moderate vulnerability (35%)	Demersal
<i>Sarotherodon melanotheron</i>	Least Concern (LC)	Low vulnerability (16%)	Demersal
<i>Tilapia busumana</i>	Least Concern (LC)	Low vulnerability (25%)	Demersal
<i>Tilapia guineensis</i>	Least Concern (LC)	Low vulnerability (19%)	Benthopelagic
<i>Tilapia mariae</i>	Least Concern (LC)	Low to moderate vulnerability (28%)	Demersal
<i>Tilapia zillii</i>	Least Concern (LC)	Low to moderate vulnerability (27%)	Benthopelagic
<i>Polypterus endlicheri</i>	Least Concern (LC)	Moderate to high vulnerability (49%)	Demersal
<i>Clarias gariepinus</i>	Least Concern (LC)	Very high vulnerability (79%)	Benthopelagic
<i>Heterobranchus longifilis</i>	Least Concern (LC)	High to very high vulnerability (69%)	Demersal
<i>Chrysichthys maurus</i>	Least Concern (LC)	Moderate to high vulnerability (46%)	Demersal
<i>Chrysichthys nigrodigitatus</i>	Least Concern (LC)	High vulnerability (60%)	Demersal
<i>Malapterurus electricus</i>	Least Concern (LC)	High to very high vulnerability (74%)	Benthopelagic
<i>Schilbe intermedius</i>	Least Concern (LC)	Low to moderate vulnerability (29%)	Pelagic
<i>Schilbe mystus</i>	Least Concern (LC)	Low to moderate vulnerability (33%)	Demersal
<i>Synodontis nigrita</i>	Least Concern (LC)	Low to moderate vulnerability (29%)	Benthopelagic
<i>Synodontis schall</i>	Least Concern (LC)	Low vulnerability (21%)	benthopelagic

Table-III: Stocks exploitation parameters of species. L_{∞} , asymptotic length; K, growth rate; t_0 , age at zero length; ϕ' growth index in length; t_{max} , longevity; Z, total mortality; M, natural mortality; F, fishing mortality; E, exploitation rate; L_{25} , L_{50} and L_{75} , lengths at which 25%, 50% and 75% of specimens are captured; E_{50} , exploitation rate reducing virgin biomass by half; E_{max} , exploitation rate at maximum relative yield per recruit; E_{10} , exploitation rate at which the marginal increase in relative yield per recruit is 10% of its value at $E = 0$; Y'/R, relative yield per recruit; B'/R, relative biomass per recruit. Highlighted parameters are from Diaby *et al.* (2020b) and Diaby *et al.* (2020c).

	Growth parameters			ϕ'	T_{max}	Mortality parameters			E	Lengths probabilities capture			Reference points for Y'/R and B'/R		
	L_{∞} (cm)	K/an	t_0 (an)			Z	M	F		L_{25}	L_{50}	L_{75}	E_{10}	E_{50}	E_{max}
<i>Brycinus imberi</i>	18.38	1.10	-0.16	2.570	2.73	3.25	2.12	1.13	0.35	9.34	9.91	10.40	0.720	0.375	0.832
<i>Chromidotilapia guntheri</i>	20.4	0.310	-0.60	2.111	9.68	1.97	0.91	1.06	0.54	9.84	10.41	10.97	0.650	0.370	0.749
<i>Chrysichthys maurus</i>	25.73	0.390	-0.44	2.412	7.69	4.61	0.99	3.62	0.78	10.04	10.62	11.19	0.560	0.339	0.699
<i>Clarias gariepinus</i>	47.78	0.390	-0.37	2.950	7.69	1.04	0.84	0.20	0.20	15.42	16.34	17.26	0.466	0.285	0.581
<i>Heterobranchus longifilis</i>	38.33	0.30	-0.52	2.644	10.00	0.96	0.75	0.21	0.22	15.05	16.35	17.65	0.567	0.343	0.699
<i>Labeo coubie</i>	33.08	0.390	-0.41	2.630	7.69	1.05	0.93	0.12	0.12	16.51	17.87	19.23	0.655	0.376	0.799
<i>Labeo parvus</i>	29.93	0.30	-0.56	2.429	10.00	1.05	0.80	0.25	0.24	11.92	12.62	13.31	0.569	0.339	0.699
<i>Marcusenius furcidens</i>	32.03	0.520	-0.31	2.727	5.77	1.91	1.13	0.78	0.41	12.69	13.30	13.91	0.550	0.327	0.688
<i>Marcusenius ussheri</i>	33.08	0.31	-0.52	2.530	9.68	1.53	0.80	0.73	0.48	13.05	13.60	14.16	0.555	0.328	0.699

<i>Mormyrus rume</i>	43.58	0.760	-0.19	3.159	3.95	2.22	1.33	0.89	0.40	17.61	19.84	22.07	0.570	0.348	0.682
<i>Petrocephalus Bovei</i>	13.13	0.850	-0.24	2.166	3.53	3.22	2.00	1.22	0.36	10.02	10.64	11.27	0.650	0.397	0.799
<i>Schilbe intermedius</i>	32.03	0.730	-0.22	2.874	4.11	3.55	1.41	2.14	0.60	11.12	11.33	11.55	0.453	0.313	0.579
<i>Tilapia zillii</i>	27.83	0.640	-0.26	2.695	4.69	2.53	1.34	1.19	0.47	8.94	10.48	12.01	0.500	0.268	0.613
<i>Hemichromis fasciatus</i>	19.95	0.96	-0.02	2.582	3.13	2.69	1.92	0.77	0.29	9.91	10.36	10.82			
<i>Oreochromis niloticus</i>	24.15	0.42	-0.01	2.389	7.14	1.26	1.06	0.20	0.16	15.95	17.77	19.6			
<i>Sarotherodon galilaeus</i>	25.2	0.84	-0.01	2.727	3.57	1.94	1.65	0.29	0.15	8.43	8.59	8.75			
<i>Tilapia mariae</i>	18.9	0.64	-0.01	2.359	4.69	1.73	1.50	0.23	0.13	8.89	9.12	9.35			
<i>Chrysichthys nigrodigitatus</i>	24.15	0.60	-0.01	2.544	5.00	1.66	1.34	0.32	0.19	11.8	12.05	12.3	0.601	0.329	0.699
<i>Synodontis schall</i>	22.05	0.73	-0.01	2.550	4.11	2.11	1.56	0.55	0.26	10.67	11.17	11.67	0.553	0.359	0.699

CONCLUSION

This study is an overview of fishermen, fishing gear types, fish species and levels of exploitation of their stocks in the Solomougou Dam Lake. It provides information to decision-makers for better control of fishing activity on this reservoir.

Some illegal fishing gears are used, resulting in overexploitation of three species, four others being close to their optimal exploitation level. However, the exploitation regime of this fishery, characterised by its low frequentation over a good period of the year, preserves the stocks of many species. Thus, the use of regulatory fishing gears could be accompanied by an increase in fishing effort to increase yields.

In any case, regular assessment of fishing gears on fish stocks should be done to ensure sustainable fishing activity in this fishery

REFERENCES

- Brodie, B.W. (2010). Occurrence, vulnérabilité à la pêche et fonction écologique des coraux, des éponges et des griffons hydrothermaux dans les eaux canadiennes. Secrétariat canadien de consultation scientifique / Région de la capitale nationale / Pêches et Océans, Ontario K1A 0E6, 56.
- Cecchi, P., Léveque, C., & Aubertin, C. (2007). L'eau en partage: Les petits barrages de la Côte d'Ivoire. Paris, IRD Edition, 295 p.
- Diaby, M., Kouassi K.D., Diaha N.C., & N'da, K. (2020c). Population Dynamics of *Chrysichthys nigrodigitatus* (Lacepede, 1803) and *Synodontis schall* (Bloch & Schneider, 1801) from Solomougou's Dam Lake (Côte d'Ivoire). *International Journal of Innovative Studies in Aquatic Biology and Fisheries*, 6(2), 11-19.
- Diaby, M., Kouassi, K.D., Akadje, C.M.A., & N'da, K. (2020b). Exploitation level of the four major Cichlid fish from the Solomougou dam lake (Korhogo, Côte D'Ivoire). *European Journal of Scientific Research*, 157(4), 440 – 449.
- Diaby, M., Kouassi, K.D., Konan, K.S., & N'Da, K. (2020a). Typology of fishermen and fishing gears from the Solomougou dam lake (Korhogo, Côte d'Ivoire). *International Journal of Fisheries and Aquatic Studies*, 8(5), 355-359.
- FAO (2005). Gouvernement de la République de Côte d'Ivoire. Appui à la mise en œuvre du NEPAD-PDDAA, TCP/IVC/2903 (I) (NEPAD Réf 05/25 F). Profil de projet d'investissement bancaire. Appui à la pêche continentale et lagunaire, 6 p.
- Froese, R., & Pauly, D. (2021). FishBase. World Wide Web electronic publication. www.fishbase.org, version 06/2021, consultée le 27 /8/2021.
- Golé Bi, G.G., Koffi, K.J.D.D., & Dadi, S.G. (2005). Contribution socio-économique de la pêche artisanale en Côte d'Ivoire. Programme pour les moyens d'existence durables de la pêche en Afrique de l'ouest. Document FAO (GCP/INT/735/UK), 49 p.
- Kouassi, K.D., Diaby, M., Amon, Y.N., & N'Da, K. (2019). Dynamic of fishing activity on the Solomougou dam lake (Korhogo, Côte d'Ivoire). *International Journal of Agriculture, Environment and Bioresearch*, 4(6), 298-307.
- Kouassi, K.D., Diaby, M., Amon, Y.N., & N'da, K. (2020b). Length-Weight relationship and condition factor for 22 freshwater fish species from Solomougou Dam Lake (Korhogo, Côte d'Ivoire). *International Journal of Fisheries and Aquatic Studies*, 8(1), 172-175.
- Kouassi, K.D., Diaby, M., Soro, Y., & N'Da, K. (2020a). Faune ichtyologique du lac de barrage Solomougou (Korhogo, Côte d'Ivoire). *International Journal of Biological and Chemical Sciences*, 14(7), 2528-2537.
- Le Guen, T. (2002). Les barrages du Nord de la Côte d'Ivoire: développement socio-économique et état sanitaire des populations. Doctorat nouveau régime en Géographie, Université de Bretagne Occidentale, Brest, France, 467 p.
- Traoré, K. (1996). État des connaissances sur les pêcheries continentales ivoiriennes. Rapport de consultation Avril 1996, PROJET FAO TCP / IVC / 4553. <http://www.fao.org/3/ag188f/AG188F00.htm#TOC>, consulté le 12 juin 2021.
- UICN. (2021). IUCN Red List of Threatened Species. Union Internationale de la Conservation de la Nature (IUCN). Version 2011-1. <http://www.iucnredlist.org>. (Page consultée le 27 /8/2021).