

Impact of Climate Change and Vulnerability Status of Dairy Farming System in Dang, Nepal

Acharya Y¹, Dhakal S.C.²

¹Technical Officer, Nepal Agricultural Research Council (NARC), Kathmandu, Nepal.

²Asst. Professor, Agriculture and Forestry University (AFU), Rampur, Chitwan.

*Corresponding Author

Acharya Y

Email: yogendraacharya9@gmail.com

Abstract: The study was conducted in Dang district to analyse the farmer's perception, impact and vulnerability of dairy farming system in the context of changing climatic condition. Total 120 livestock farmers were selected from two VDCs using purposive simple random sampling. Primary data were collected through household survey and Focus Group Discussion using structured and pretested interview schedule. Descriptive statistics, integrated vulnerability approach based on principle component analysis and indexing were used to analyse the collected data. Agriculture was major source of household income for majority of respondent households (67.5%) following foreign employment (17.5%) and off farm activities (15%). The average livestock holding was nearly 16.81 livestock standard unit that contribute 31 percent to annual household income. A large majority of respondents (89.0%) observed the weather variation. Farmers perceived incidence of livestock disease (95%), decrease in milk yield (75.0%) and unavailability of feed and grasses (66.7%) as the major impact of climate change on livestock production. Most of respondents felt that extreme hot, drought spells, and floods were most prioritized climatic hazards in study area. Integrated vulnerability approach based on principle component analysis was used to analyze the livelihood vulnerability of livestock farmers. The vulnerability index of Dang district was found to be 4.4 which signified high risk of climate change in the district. The study concluded that farming practices and livelihood options of large portion of livestock farmers were affected negatively. Based on results, awareness raising program, effective adaptation strategy like livestock insurance, institutional enrollment and extension service with new technologies are recommended to enhance the adaptive capacity of vulnerable livestock communities in changing climatic scenario.

Keywords: Vulnerability, dairy farming system, climate change, Principal component analysis, Dang

INTRODUCTION

Livestock being integral part of Nepalese farming system has substantial role in the national economy. Climate change as evidenced by increased climatic variability and temperature, is affecting the livelihoods of rural poor and marginal farmers in Nepal [1]. Therefore, climate change has been a real threat to livelihood of Nepalese farmers due to the problematic impact on their crop and livestock production system. The climate change would likely result in considerable adverse impacts on breeding pattern, fertility, repeated breeding, productivity, disease incidence, and reduced feed conversion efficiency in warmer regions like terai part in general and western dry arid terai in particular of Nepal. Changing climatic condition clearly plays a key role in poor livestock performance [2]. However, there are considerable gaps in research findings to know how climate change affects livestock production system and the livelihoods of farmers in the context of large microclimatic variation in Nepal.

National target for achieving growth rate in livestock sector during twenty years periodic plan is 6.1% in 2015 from 2.9% at base year period (APP,

1995). But the livestock sector being highly vulnerable to climate change and due to other technical and institutional obstacles related to it the target rate could not be achieved. Studies on climate variability at local level are very sporadic in spite of the accepted fact that the micro-level community vulnerability and adaptation assessment is more relevant than the mega-scale. Therefore, assessing vulnerability provides a starting point for the determination of effective means of promoting remedial action to limit impacts by supporting coping strategies and facilitating adaptation [3]. Identification of impact of climate change helps to formulate appropriate coping strategies and further recommends for policy makers to give due attention to this phenomenon so as to make farmers sustain in this farming sectors. This study will help to identify previous research gaps and impact of climate change on livestock production and its effect on livelihood in specific location along with types of impact associated. The findings from this proposed study will be useful to support timely improvement of these policies and formulation of new policies after identifying the level of vulnerability in livestock sector.

MATERIALS AND METHODS

This research was conducted in two VDCs, namely Lalmatiya and Urahari of Dang district of Nepal. Two VDCs were selected purposively in consultation with DLSO based on vulnerability to climate change impact and commercialization of livestock farmers. From these two VDCs, each two wards were selected randomly using simple random sampling techniques. Sampling frame was prepared taking the name list of livestock farmers of selected two ward of each VDC and simple random sampling technique was used to select the respondents in the size 120 from two different VDCs under study. Various sources and technique were used for collection of necessary information. In this study, both the primary and secondary data were collected. The methodologies consisted of household survey, review of previous studies, and interviews with key informants, and also direct observation by the researchers. Livestock farmers were the primary source of information. The pre-tested interview schedule was administered to the respondents to collect primary information. All randomly selected participants were visited and interviewed. Primary data were also collected through Focus Group Discussion (FGD) and Key Informant Survey (KIS) using interview guide. Meteorological data like temperature, precipitation and relative humidity (RH) of the proposed study area from 1981 to 2013 was taken from Department of Hydrology and Meteorology, Kathmandu. Other secondary information related to livestock production performance, livelihood and other socio economic information of country and study area were collected from various published materials and reports of different National and International organization. District agriculture information was collected from District Agriculture Development Office and District Livestock Service Office. After collection of necessary information it was coded and entered to SPSS software for analysis. Data analysis was done by using statistical packages for social sciences (SPSS V.20) and STATA 12. Mean, standard deviations, frequency, percentage were used to derive inference needed.

For obtaining the information on the livestock based livelihood vulnerability in the study area, integrated vulnerability assessment approach was used. Vulnerability Assessments are the crucial component to study the livelihood condition of the people. According to Deressa, Hassan, & Ringler, et al. [4], the integrated vulnerability approach is considered to be superior over other approaches and is particularly useful for policy decision. Accordingly, socio-economic and biophysical indicators of vulnerability were employed and classified into adaptive capacity, exposure and sensitivity.

For the analysis of vulnerability through integrated approach, both primary and secondary sources of data were used. The chosen indicators of adaptive capacity, exposure and sensitivity were separately employed for PCA. The loading from the first component of PCA were used as the weights for the indicators. The weights assigned for each indicator varies between -1 to +1, sign of the indicators denoting the direction of relationship with other indicators used to construct the respective index. The magnitude of the weights describe the contribution of each indicator to the value of the index.

Thus, vulnerability was calculated as defined by IPCC (2001),

$$\text{Vulnerability} = \text{Adaptive capacity} - \text{Sensitivity} - \text{Exposure}$$

Equation above can be operationalized as follows:

$$V = \sum_{i=1}^n W_i \cdot X_i - (\sum_{i=1}^n W_i \cdot Y_i + \sum_{i=1}^n W_i \cdot Z_i)$$

Where

$i = 1, 2, 3, 4, \dots, n$ households

V = Vulnerability index

W_i = Weight obtained from first principal component scores of i^{th} variable.

X_i = Adaptive capacity of i^{th} variable

Y_i = Sensitivity of i^{th} variables

Z_i = Exposure of i^{th} variables

The relationship between Vulnerability Index and Vulnerability is negative. The higher value of vulnerability index indicates lesser vulnerability and vice versa condition [5]. But, negative value of the index doesn't imply that the household is not vulnerable at all. The index value gives the comparative ranking of sampled households/ study area rather absolute measurement of vulnerability.

The suitable variables used for calculation of vulnerability index need to be normalized so as to bring the values of variables within the comparable range [6, 7]. Normalization is done by subtracting the mean from the observed value and dividing by the standard deviation for each variable.

$$\text{Normalized value} = \frac{\text{Observed value} - \text{mean}}{\text{Standard deviation}}$$

Climatic hazards and extremities in the study areas were identified with the use of index. The formula given below was used to find index for severity of climatic hazards.

$$I_{\text{prob}} = \sum SiFi/N$$

Where,
 I_{prob} = Index value for intensity of problem
 S_i = Scale value of i th intensity
 F_i = Frequency of i th response
 N = Total number of respondents

latitude and between 80°02' to 82°54' East longitude. It is bordered with Kapilbhatu and Argakhanchi district in the east, Salyan, Rolpa and Piuthan in the North, Uttar Pradesh of India in the South and Surkhet, Banke and Salyan in the West. It extends 90 km east west and 71 km north south and the area of 2955 square kilometer which occupies 2.03% area of Nepal. The elevation varies between 213m. And 2058 m. above mean sea level [8]. There are 116415 households with total population 552583 (CBS, 2011).

RESULTS

Description of the study area

The Dang is one as Inner Terai district of Mid-Western Development Region. Geographically, Dang district has been expanded from 27°36' to 28°29' North

Table-1: Land use pattern of the district

Particulars	Area in (Ha)
Total area	2,95,500
Forest	2,01,900
Cultivable land	69,950
Cultivated land	65,580
Low land	38,600
Upland	26,980
Pastureland	12,950
Others	10,700

Source: DADO, Dang, 2013/2014

Socio-economic and demographic characteristics

The socio-economic characteristics of respondents include age distribution, population and sex distribution, economically active population, education, land utilization pattern, involvement of farmers in local institution, group, access to credit etc. These characteristics are described below.

The average age of the respondent is 44.22 years ranging from 16 to 88 years. Whereas the average age of the household head was 49.01 years and ranges from 24 years to 83 years. The total population of the sampled household was 702, of which average family size of the survey district area was 5.86 which is higher than the national average 4.46 (CBS, 2011). The household were found as patriarchal, out of the 120 household that were sampled in area 65.0 % was male headed family remaining 35.0 % was only the female headed family. The study revealed that the majority of

the population were Chettri (57.5%) followed by the Brahmin (24.2%) and Ethnic/indigenous (13.3%) and other occupational caste as 5.0%.

Occupational and educational status and economically active population

The economically active population, educational and occupational status were studied. The major occupation was categorized as the agriculture, off farm income and foreign employment. From the study the majority of the studied household (67.5%) have the agriculture as their major occupation followed by the foreign employment (17.5%). Only 15 percent household sustain their livelihood mainly from off farm income. The study revealed that the majority of respondents have secondary level education. Relatively fewer respondents (10.8%) have attained higher level education.




Fig-1: Economically active population

The number of economically active population was categorized into three group; less than 15 years, 15-59 year and more than 59 year. The study revealed that more than two third of population (67.1%) were economically active.

Livestock Standard Unit

The average livestock holding size was 16.81 LSU. The minimum holding was found 2.60 LSU while maximum was recorded in 503.80 LSU. The maximum LSU (503.8) in study area may be due to large number of poultry production in commercial scale.

Table-2: Livestock holding (LSU)

Livestock holding (LSU)	VDC		Total
	Lalmatiya	Urahari	
Mean	22.80	10.81	16.81
S.D	81.41	10.34	58.09
Minimum	2.60	2.60	2.60
Maximum	503.80	82.00	503.80

Source: Field Survey, 2015

Perception of respondents towards climate change

About 89 percent had observed the change in the climate and 11 percent did not observed any changes in climate.




Fig-1: Perception of respondents towards the climate change

Climatic and biological hazards prioritized by farmers

Flood, drought, extreme hot and hailstorm were four major climatic problems in the study area of Dang district. Similarly crop disease, livestock internal/external parasite, loss in grass land and loss in forage biodiversity were four major biological hazards.

The result showed that extreme hot (0.67) was major climatic hazard in study area followed by drought (0.580) and least climatic hazard was hailstorm (0.30). Likewise, crop disease pest outbreak, livestock internal and external parasite, loss in grass land and loss in forage biodiversity were major biological hazard.

Table-3: Perception of respondent towards major climatic hazards and prioritization

Climatic Hazards	Index value	Ranking	Biological Hazards	Index value	Ranking
Floods/Riverbank Erosion	0.462	III	Crop disease pest outbreak	0.731	I
Drought/Dry Spell	0.580	II	Livestock internal/external parasite	0.698	II
Extreme Hot	0.673	I	Loss in grass land	0.623	III
Hailstorm	0.304	IV	Lose in forage biodiversity	0.387	IV

Source: Field Survey, 2015

Perception on changing temperature and rainfall

There is a gradual increase in temperature in different ecological zones of the western development region of Nepal [9]. Most of the respondents (92.5%) felt the increase in general temperature level. 95.0% of the respondents felt that the number of hot summer days has increased and 92.5% respondents felt the increase in

degree of hotness of summer days which was also proved by the increasing trend of maximum, average and minimum temperature from Meteorological Station of Dang district. Farmers also added that local community has become more vulnerable to spread of vectors and diseases both to human and livestock affecting livestock production.

Table-4: Perception of respondents towards the change in weather condition

Weather condition	Perception			Don't know
	Increased	Same	Decreased	
Temperature level	111(92.5)	4(3.3)	0(0.0)	5(4.2)
No. of hot days in summer	114(95)	4(3.3)	0(0.0)	2(1.7)
Degree of hotness in summer	111(92.5)	7(5.8)	1(0.8)	1(0.8)
No. of cold days in winter	5(4.2)	14(11.7)	99(82.5)	2(1.7)
Degree of coldness in winter	51(42.5)	40(33.3)	28(23.3)	1(0.8)
Winter rainfall intensity	20(16.7)	31(25.8)	45(37.5)	24(20.0)
Winter rainfall frequency	16(13.3)	36(30)	44(36.7)	24(20.0)
Summer rainfall intensity	41(34.2)	6(5.0)	72(60.0)	1(0.8)
Summer rainfall frequency	8(6.7)	8(6.7)	101(84.2)	3(2.5)




Fig-2: Maximum, Minimum and Average temperature as recorded in Meteorological Station, Dang (1981-2013).

Farmers perceive that there was wide variation in the rainfall pattern as compared to the past. Majority of the farmers, 60% and 84.2% perceived decrease in intensity and frequency of rainfall respectively as

compared to the past years. Majority of the respondents, 36.5% reported decline in frequency and 37.5% reported intensity of the winter rainfall as compare to the past.




Fig-3: Average rainfall as recorded in Meteorological Station, Dang (1981-2013).

Perception on impact of climate change on livestock

The climate change impact is widely felt and its effect is experienced across all the regions of the Nepal. Livestock is the most sensitive farming system felt due to climate change About 95.0 percent respondents felt that the incidence of disease and

parasites in livestock, 66.70 percent respondent felt the unavailability of feed and grass, 75 percent felt decrease in milk production, 40.8 percent felt death of animal, 22.5 percent felt change in appetite of livestock and 49.2 percent felt heat stress in animal (Table 5).

Table-5: Major climate change impact on livestock perceived by respondent

Major impacts on livestock	Total
Unavailability of feed and grass	80(66.7)
Incidence of disease and parasites	114(95.0)
Decrease in milk production	90(75.0)
Heat stress	59(49.2)
Death of animal	49(40.8)
Loss of appetite	27(22.5)

Source: Field Survey, 2015

Barriers to adoption of adaptation strategy

The analysis of barriers to adaptation to climate change based on the perception of respondents in the study area indicated that there were six major constraints to adaptation. These were lack of government support, lack of availability of suitable

adaptation strategy, lack of climate information, lack of appropriate adaptation knowledge, lack of money and poor market access. (Table 6). Around 44 percent of respondents strongly agreed that lack of government support was first most barriers to adoption of adaptation strategies.

Table-6: Perception of respondent towards barrier to adaptation

Reasons	Level of agreement			Neutral
	Strongly agree	Agree	Disagree	
Lack of government support	52(43.3)	36(30.0)	12(10)	20(16.7)
Availability of suitable adaptation strategy	42(35.0)	75(62.5)	1(0.8)	2(1.7)
Lack of climate information	35(29.2)	72(60.0)	7(5.8)	6(5.0)
Lack of technology	34(28.3)	81(67.5)	1(0.8)	4(3.3)
Lack of appropriate adaptation knowledge	30(25)	88(73.3)	2(1.7)	0(0.0)
Lack of credit/Poverty	25(20.8)	33(27.5)	34(28.3)	28(23.3)
Poor market access/transport	8(6.7)	85(70.8)	6(5.0)	21(17.5)

Figures in the parenthesis indicate percent

Source: Field Survey, 2015

Livestock holder's vulnerability to climate change

Adaptive capacity, sensitivity and exposure are major three components of Vulnerability assessment. The adaptive capacity was taken for socio-demographic, economic, institutional, gender inclusion characteristics. Vulnerability indices were obtained by applying Principal Component Analysis on the adaptive capacity, sensitivity, and exposure variables. Principal component analysis is frequently used in research that constructs indices for which there are no well-defined weights. Principal component analysis generated the weights, based on the assumption that there is a common factor that explains the variance in the vulnerability [4].

The result of Principal component Analysis (PCA) showed that eight component with the Eigen value of one or greater than one accounting for 66.64

percent of total variance. The first component had Eigen value of 3.548 contributing 16.89 percentage of cumulative variance followed by the second component with Eigen value of 2.165 accounted for 27.20 percent of cumulative variance. The weights obtained from PCA are given in Table 8 for the indicators of adaptive capacity, exposure and sensitivity respectively. Only the component score of the first component was used weighing the variables for the construction of vulnerability index.

The vulnerability index for the Dang district was found to be 4.4 (Table 7). The calculation of vulnerability index showed that Dang district had positive value indicating that Dang was less vulnerable. However the index being relative value must be used for comparison with indices of other areas.

Table-7: Vulnerability index for study area

District	Index score
Dang	4.4

Table-8: Component scores of coefficient for component first through PCA

Indicators	Types of variables	Component score
Age of household head	Adaptive capacity	-0.093
Sex of household head	Adaptive capacity	0.551
Education of household head	Adaptive capacity	0.566
Economically active member	Adaptive capacity	-0.028
Land size	Adaptive capacity	0.393
Livestock holding	Adaptive capacity	0.266
Information on climate change	Adaptive capacity	0.346
Credit facility	Adaptive capacity	0.321
Institutional membership	Adaptive capacity	0.072
Extension service available	Adaptive capacity	0.262
Training on livestock	Adaptive capacity	0.460
Nature of dairy farming	Adaptive capacity	0.505
Maximum temperature variability	Exposure	0.022
Minimum temperature variability	Exposure	-0.029
Precipitation variability	Exposure	-0.824
Relative humidity variability	Exposure	0.142
Summer temperature	Exposure	0.027
Drought	Exposure	0.347
Flooding	Exposure	0.287
Animal death	Sensitivity	0.318
Income share from natural resource based sector	Sensitivity	0.861
Income share from non-natural based sector	Sensitivity	-0.861
Dependency ratio	Sensitivity	-0.128
Livestock diversification practices	Sensitivity	-0.082
Employment diversification practices	Sensitivity	-0.340

DISCUSSION

The study revealed that about two third (65.0%) of respondent's households were male headed

with average family size of 5.86. Still, agriculture was found to be most reliable source of household income of majority of sampled household (67.5%) followed by

foreign employment (17.5%) and off farm activities (15.0%). Majority of the surveyed households were of Chhetri ethnicity (57.5%). Livestock holding in study area was (16.81 LSU) ranging from 2.60 to 503.80. The maximum LSU (503.80) may be due to commercial poultry farming. Though the percentage of economically active member was more than 66 percent, there involvement in agricultural activity has declined year after year. The lower profit from agriculture as compared to other livelihood options like wage laboring, remittance and service were considered as a main reason for declining economically active member in agriculture. A large portion of respondents felt that there was changes in most of the weather patterns like increase in degree of hotness, temperature level, degree of coldness and fluctuations in rainfall intensity and duration. Farmer perception towards weather parameter was further supported by the analysis of meteorological data of Dang district. Vulnerability, the degree to which a system is susceptible to the adverse effects of climate change, is composed of three components: adaptive capacity, exposure and sensitivity [10]. Integrated vulnerability assessment approach was used to determine the climate change vulnerability index of the study area as it overcomes the shortcomings of other methods.

The vulnerability of the study area was found to be 4.4, which falls in the high risk zone [11]. The livestock vulnerability to climate change of Kapilbastu district, neighbor district of Dang, was found to be 7.41 [12]. So the Dang district vulnerability, when compared to Kapilbastu, is said to be higher. Higher vulnerability may be due to higher incidences of drought and higher variation in climate parameter, illiteracy rate, lack of year round irrigation, higher number of economically active population engaged in study.

CONCLUSION

Livestock farming was one of major source of household income for majority of farming households. Although livestock plays dominating role on the livelihood of marginal and poor people; climate change impact negatively affects the livestock production. Most of the livestock keepers felt that extreme hot, drought, flooding and hailstorm were the perceived climatic hazards and crop disease, livestock disease, loss of forage land and loss of forage diversity were another biological hazard in the study area. Vulnerability of any regions depends upon the adaptive capacity, exposure and sensitivity. The adaptive capacity of the livestock keepers helps to combat the negative effect from the climate change. Age and sex of household head, high education level, number of economically active member, more no. of livestock, institutional membership, more clear weather information, availability of extension service, access to credit were

the adaptive capacity of livestock keepers which were seen as strengthening factors for negative changing climate change scenario. The result revealed that vulnerability index of Dang district being positive, implies less vulnerable as compared to vulnerability index with negative value. However, Dang district has been said to be relatively more vulnerable to climate change if compared to other terai district of western Nepal. Therefore, climate change program or policy launched by government or nongovernment sector should focus on awareness raising activities with suitable adaptation strategy and technology intervention.

ACKNOWLEDGEMENT

We are very much obliged to thank National Agriculture and Development Fund (NARDF) for providing financial support for this study. The author would like to acknowledge all personal and institutional helping hands for different technical and administrative supports. The authors are equally grateful to the farmers of Dang district, Nepal for providing invaluable information and shared their experience with great hospitality.

REFERENCES

1. Dahal, H., & Khanal, D. R. (2010, February). Food security and climate change adaptation framework: issues and challenges. In *second stakeholders workshop on NAPA in agriculture sector, Kathmandu* (Vol. 23).
2. Luborth, J. (2012). Climate change and animal health. *Building Resilience for Adaptation to Climate Change in the Agriculture Sector* 23, 63.
3. Kelly, P. M., & Adger, W. N. (2000). Theory and practice in assessing vulnerability to climate change and facilitating adaptation. *Climate Change* 47(4), 325-352.
4. Deressa, T., Hassan, R. M., & Ringler, C. (2008). Measuring Ethiopian farmers' vulnerability to climate change across regional states. *Environment and Production Technology Division* (p. 8). IFPRI Discussion Paper 00806
5. Madu, C. (Ed.). (2012). *Handbook of total quality management*. Springer Science & Business Media.
6. Nelson, R., Kokic, P., Crimp, S., Martin, P., Meinke, H., S.M., et al. (2010). The Vulnerability of Australian Rural Communities to Climate Variability and Change: Part II- Integrating Impact with Adaptive Capacity. *Environment Science and Policy*, 13, 18-27.
7. Gbetibouo, G. A., & Ringler, C. (2009). *Mapping South African farming sector vulnerability to climate change and variability: A subnational assessment*. International Food Policy Research Institute (IFPRI) and Center for Environmental Economics and Policy in Africa (CEEPA).

-
8. DADO. (2014). *Dang District Profile*. Dang: District Agriculture Development Office.
 9. Manandhar, S., Vogt, D. S., Perret, S. R., & Kazama, F. (2011). Adapting cropping systems to climate change in Nepal: a cross-regional study of farmers' perception and practices. *Regional Environmental Change*, 11(2), 335-348.
 10. IPCC. (2014). *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*.
 11. de Sherbinin, A. (2014). Climate change hotspots mapping: what have we learned?. *Climatic Change*, 123(1), 23-37.
 12. Dev, R. (2014). *Assessment of climate change impact and adaptation strategies for livestock production through gender perspective in Gandaki river basin of Nepal* (Master thesis, Institute of Agriculture and Animal Science, Tribhuvan University, Chitwan, Nepal).