

Research Article

Preliminary Phytochemical Investigation and Antioxidant Activity of *Bougainvillea glabra* choicy Leaves

J. Mariajancyrani^{1*}, G. Chandramohan¹, M. Fathima Beevi², A. Elayaraja¹

¹Department of chemistry, A.V.V.M. Sri Pushpam College, Poondi, Thanjavur-613 503, Tamil Nadu, India.

²Department of chemistry, S.T.E.T. Women's College, Poondi, Mannargudi, Tamil Nadu, India.

Corresponding author

J. Mariajancyrani

Email: mariajancyrani@yahoo.in

Abstract: Antioxidants are an inhibitor of the process of oxidation, even at relatively small concentration and thus have diverse physiological role in the body. The present study was carried out to identify the phytochemicals and evaluate antioxidant activity of chloroform extract of *Bougainvillea glabra* choicy leaves. The antioxidant activity was determined by different methods via DPPH radical scavenging assay, Hydrogen peroxide scavenging assay and reducing power assay. The leaves of chloroform extract contain terpenoids, steroids, flavonoids and Tannins. The antioxidant activity found in DPPH method is more effective than other two methods. Thus the in-vitro studies clearly indicate that the chloroform extract of *Bougainvillea glabra* choicy leaves shows significant antioxidant activity and also a better source of natural antioxidant, which might be helpful in preventing the process of various oxidative stresses.

Keywords: Antioxidant activity, flavonoids, terpenoids, Hydrogen peroxide, Tannins

INTRODUCTION

Plants have always been a part of medicinal science from the beginning of human civilization to the present modern world of synthetic medicines. Even in the presence of variety of effective synthetic drugs, use of medicinal plants for maintaining human health has acquired a lot of importance in the present era [1]. There is a global interest in non-synthetic, natural drugs derived from plant sources, because of low cost, non-toxic nature and availability. Many plants with antioxidant potential possess flavonoids and phenolic compounds [2]. Free radical reactions have been implicated in the pathology of many human disease including atherosclerosis, ischemic heart disease, the aging process, inflammation diabetes and other conditions [3].

The research for medicinal plants discovered various common plants having distinguishable medicinal properties, among which one is *Bougainvillea glabra*. *Bougainvillea glabra* is an ornamental flowering plant from the genus of *bougainvillea*; family *Nyctaginaceae* [4] and a native to Brazil. The genus *bougainvillea* has eighteen species, of which *B. spectabilis*, *B. glabra* and *B. Peruvian* [5] are horticultural important. *Bougainvillea glabra* is a woody climber with thorny thin stems and long branches; also it has papery bracts and smooth leaves, which grows to more than 10 meters of height [6].

B. glabra is one that is great for container plantings and produce brilliant colors. The plant is perfect for arbors and draped long fences where it creates a security barrier by its thorns and twiggy growth [7]. Body has itself antioxidant system, which

reacts with reactive species and neutralizes them. This natural antioxidant system includes enzymes like catalase, superoxide dismutase and glutathione, which protect the body from free radical species and prevent oxidative stress [8]. Synthetic antioxidant like butylated hydroxyl toluene and butylated hydroxyl anisole are carcinogenic in nature. So, there arises a need for natural antioxidant [9]. The antioxidant activity of the hydroalcoholic, acetone extracts of the leaves were evaluated for this plant [10]. With this background, the aim of the present study was to determine the possible phytochemical and antioxidant activity of chloroform extract of *Bougainvillea glabra* choicy leaves.

MATERIALS AND METHODS

Plant Material

Plants were collected from Thanjavur District of Tamilnadu. The botanical identity of the plant of *Bougainvillea glabra* was confirmed by Dr. John Brito, Rapinet Herbarium. St. Joseph's College, Thiruchirappalli.

Preparation of Extract

The powder (5kg) was extracted with 95% ethanol at room temperature for seven days. The extracts were filtered and concentrated under reduced pressure in a rotary evaporator and for further fractionated successively with n-hexane, chloroform, ethyl acetate, acetone, ethanol, butanol and methanol solvents were then removed under reduced pressure. From the fractionation chloroform extract were subjected to screening of phytochemical and antioxidant activity.

Phyto Chemical Analysis

Phytochemical analysis involves the qualitative analysis of herbal plants. The preliminary qualitative tests have been attempted in *Bougainvillea glabra* choicy to find out the presence or absence of certain bio active Compounds. Chemical tests were carried out on the chloroform extract using standard procedures to identify the constituents as described by Harbone [11].

Test for Tannins

About 0.5g of the dried powdered samples was boiled in 20ml of water in a test tube and then filtered. A few drops of 0.1% ferric chloride were added. Presence of brownish color indicated the presence of tannins.

Test for Phlobatannins

Extract boiled with 1% aqueous hydrochloric acid. No red precipitate is formed indicated absence of phlobatannins.

Test for Saponin

About 2g of the powdered sample was boiled in 20ml of distilled water in a water bath and filtered. 10ml of the filtrate was mixed with 5ml of distilled water and shaken vigorously for a stable persistent froth. No change in solution. Saponin is absent.

Test for Flavonoids

Few drops of 1% aluminum solution were added to a portion of extract. A yellow colouration was observed indicating the presence of flavonoids.

Test for Oil and Fat

Stain test

Small Quantity of extract was pressed between two filter papers. An oily stain on filter paper indicates the presence of fixed oil.

Test for Steroids

2ml of acetic anhydride was added to 0.5g ethanolic extract with 2ml sulphuric acid. The color changed from violet to blue indicating the presence of steroids.

Test for Terpenoids (Salkowski Test)

5ml of extract was mixed in 2ml of chloroform, and 3ml of conc. Sulphuric acid was carefully added to form a layer. A reddish brown colouration of the interface was formed to show positive results for the presence of terpenoids.

Test for Alkaloid

Mayer's Test

0.5 ml of extract is added with 1 ml of mayer reagent. Absence of yellow colour indicated absence of alkaloid.

Dragandroff's test

0.5ml of extract is added with 1ml of dragandroff's reagent. Absence of orange red colour indicated absence of alkaloid.

Wagner's Test

0.5ml Sample is treated with 1 ml of wagner reagent. No change in colour indicated absence of alkaloids.

Antioxidant Activity

DPPH method

0.1 ml of the chloroform extract was taken in test tubes. 6 ml of DPPH (diphenyl picryl hydrazyl) solution was added and the tubes kept in dark for one hour. The color was read at 517 nm. The difference in the Optical density of DPPH solution and DPPH solution + sample was calculated. The decrease in OD with sample addition is used for calculation of the antioxidant activity. The activity was compared with BHT (butylated hydroxyl toluene) standard [12]. Free radical scavenging activity was expressed as the inhibition percentage calculated using the formula.

$$\text{Percentage of antioxidant activity} = \frac{[\text{Absorbance of control} - \text{Absorbance of sample}]}{\text{Absorbance of sample}} \times 100$$

Reducing power assay

Reducing power assay of sample was done using published protocols [13]. 1 ml of chloroform extract was mixed with phosphate buffer (2.4 ml 0.2 M pH 6.6) and potassium ferricyanide (2.5 ml). The mixture was incubated at 50 °C for 20 minutes. A portion (2.5 ml) of trichloroacetic acid (10%) was added to the mixture, which was then centrifuged at 3000 rpm for 10 minutes. The upper layer of solution (2.5 ml) was mixed with distilled water (2.5 ml) and Ferricchloride (0.5 ml, 0.1%) and absorbance measured at 700 nm. Increased absorbance of the reaction mixture indicates stronger reducing power. The activity was compared with BHT standard. Scavenging activity was expressed as the inhibition percentage calculated using the formula.

$$\text{Percentage of antioxidant activity} = \frac{[\text{Absorbance of control} - \text{Absorbance of sample}]}{\text{Absorbance of sample}} \times 100$$

Hydrogen peroxide scavenging activity

To determine the Hydrogen peroxide assay of chloroform extract by Umamaheswari and Chatterjee Method [14]. Hydrogen peroxide solution (2 mM/L) was prepared with standard phosphate buffer (pH- 7.4). Different concentration of the extracts in distilled water was added to 0.6 ml of hydrogen peroxide solution. Absorbance was determined at 230 nm after 10 minutes against a blank solution containing phosphate buffer without hydrogen peroxide. The inhibition was calculated. BHT was used as standard. Scavenging activity was expressed as the inhibition percentage calculated using the formula.

$$\text{Percentage of antioxidant activity} = \frac{[\text{Absorbance of control} - \text{Absorbance of sample}]}{\text{Absorbance of sample}} \times 100$$

RESULTS AND DISCUSSION

Table 1: Qualitative analysis of phytochemicals of *Bougainvillea glabra* choicy leaves.

Sl. No.	Phytochemical	Result
1	Alkaloids	Absent
2	Pholobtannin	Absent
3	Tannin	Present
4	Terpenoid	Present
5	Saponin	Absent
6	Flavonoids	Present
7	Steroids	Present
8	Fatty acid	present

Table 2: Antioxidant activity of chloroform extract of *Bougainvillea glabra* choicy leaves.

Sl no.	Method	Control Absorbance	Sample Absorbance	% Antioxidant activity	Standard (BHT) (0.01 mg/ml)
1	DPPH assay	1.586	0.421	73.45	78.74
2	Reducing power assay	1.007	0.538	46.57	69.72
3	Hydrogen Peroxide assay	1.389	0.983	29.22	45.92

The antioxidant activity showed the inhibition percentage is 73.45 %, 46.57 % and 29.22 % respectively. The results observed from chloroform extract of *Bougainvillea glabra* choicy leaves shows higher antioxidant potential and compared to three methods DPPH assay observed high antioxidant activity. The results were compared to the standard BHT, it was only slight difference has been noted. The WHO estimated that 80% of the population of developing countries still relies on traditional medicine, mostly plant drugs for their primary health care needs. Hence, there is an urgent need to study the screening of antioxidant properties of herbs which will be helpful in the treatment of several diseases [15].

Antioxidants are an inhibitor of the process of oxidation, even at relatively small concentration and thus have diverse physiological role in the body. Antioxidants may be synthetic or natural. Synthetic antioxidants such as BHT and BHA have recently been reported to be dangerous for human health. Thus, the

The result of the phytochemical screening and antioxidant activity are given in the Table1, 2. Terpenoids, fatty acids, flavonoids, steroids, Tannins were present in *Bougainvillea glabra* choicy leaves and alkaloids, saponins, Pholobtannin were absent in chloroform extract *Bougainvillea glabra* choicy leaves. search for effective, non-toxic natural compound with antioxidative activity has been intensified in recent years [16]. On the basis of our results, *Bougainvillea* appears to have potential for treatment of oxidative stress related diseases. It should, however, be explored as a functional medicinal plant for isolating the active ingredients along with animal studies in vivo.

Acknowledgement

I wish to express my deep sense of gratitude and most sincere thanks to S. Kumaravel, senior scientist, Indian Institute of Crop Processing Technology, Thanjavur for providing all the facilities and support to finish my work.

REFERENCES

1. Joshny J, Ramya Devi D and Vedha Hari BN; Anti-cancer and Anti-Microbial Activity of Hydro Alcoholic Extract of *Bougainvillea glabra*. International Journal of Chemical and Pharmaceutical Research, 2012; 3(4):79-85.
2. Gupta V, George M, Joseph L, Singhal M and Singh HP; Evaluation of antibacterial activity of *Bougainvillea glabra* ‘snow white’ and *Bougainvillea glabra* ‘choicy’. Journal of chemical and Pharmaceutical Research, 2009; 1(1): 233-237.
3. Tomar RS and Sisodia SS; Estimation of phenolic content, total flavonoids and *in-vitro* antioxidant activity of *annona squamosa* linn and *Bougainvillea glabra* choicy. Journal of Global Pharma Technology, 2013; 5(3).
4. Jawa S, Kumar Y and Khan MSY; Hypoglycemic activity of *Bougainvillea spectabilis* stem bark in normal and alloxan – induced diabetic rats. Asian pacific Journal of Tropical Biomedicine, 2012; S919-S923.
5. Umamaheswari A, Shreevidya R and Nuni A; *In vitro* antibacterial activity of *Bougainvillea spectabilis* leaves extracts. Advances in Biological Research, 2008; 2(1-2):01-05.
6. Joshny J, Devi RD and Vedha Hari BN; Phytochemical and *in-vitro* anthelmintic activity of hydro alcoholic extract of *Bougainvillea glabra*. IJPPS, 2012; 4(2).
7. El-Quesni FEM, Kandil MM and Mona H. Mahgoub; Some studies on the effect of putrescine and paclobutrazol on the growth and chemical composition of *Bougainvillea glabra* L. at Nubariaed. American –Eurasian J.Agric. & Environ.Sci, 2007; 2(5): 552-558.
8. Elumalai A, Chinna Eswariah M, Vidhyulatha Chowdary CH, Ranjitj Kumar, Anusha M and

- Naresh K; Screening of Thrombolytic activity of *Bougainvillea glabra* leaves extract by *In-vitro*. Asian J. Res. Pharm. Sci, 2012;2(4):134-136.
9. Ravi Narayan Venkatachalam, Kanchanlata singh, Thankamani Marar; *Bougainvillea spectabilis*, a good source of antioxidant phytochemicals. RJPBCS, 2012; 3(3):605.
 10. Harborne JB; Phytochemical methods. A guide to modern techniques of Plant Analysis, Chapman and Hall, London; 2005: 182-189.
 11. Gupta V and Rathore DS; Evaluation of AntiOxidant Activity and Determination of Phenolic Content of Leaves Extracts of *Bougainvillea Glabra* 'Snow White'. Pharmacologyonline, 2011; 2: 219234.
 12. Umamaheswari M and Chatterjee TK; *In vitro* antioxidant activities of the fractions of *Coccinnia grandis* L Leaf extract. Afr J Trad Compl Altern Med, 2008; 5: 61-73.
 13. Yildirim A, Mavi A and Kara A; Determination of Antioxidants and Antimicrobial of *Rumex crispus* L. extracts. J. Agric. Food Chem, 2001; 49: 4083-4089.
 14. Gyamfi MA, Yonamine M, Aniya Y; Free-radical scavenging action of medicinal herbs from Ghana *Thonningia sanguinea* on experimentally- induced liver injuries. Gen. Pharmacol, 2002; 32:661-667.
 15. Vani TV, Rajani M, Sarkar S and Shishoo CJ; Antioxidant properties of the ayurvedic formulation Triphala and its constituents. Inter. J. Pharmacognosy, 1997; 35:313-317.
 16. Sahu N and Saxena J; *Bougainvillea glabra* a natural antioxidant: A review. Inventi Rapid: Planta Activa, 2012:135.