Scholars Academic Journal of Biosciences (SAJB)

Sch. Acad. J. Biosci., 2014; 2(2): 132-136

©Scholars Academic and Scientific Publisher (An International Publisher for Academic and Scientific Resources) www.saspublishers.com

ISSN 2321-6883 (Online) ISSN 2347-9515 (Print)

DOI: 10.36347/sajb.2014.v02i02.011

Review Article

Haritaki A Boon To Herbalism - A Review

Dr S. Aruna*1, Dr L.V. Nandakishore2

¹Asst Professor, Dept of computer Science, A.M Jain College, Chennai -114. ²Professor, Dept of Mathematics, Dr MGR Educational and Research Institute University, Chennai-95.

*Corresponding author

Dr S. Aruna

Email: arunale lla palli@yahoo.com

Abstract: The demand for herbal therapeutics is increasing gradually in the world. The Indian system of medicines uses herbal preparations in majority for the management of diseases. Globally a large number of Pharmacological studies have been conducted extensively on various medicinal plants. In recent years a lot of research is being conducted on *Terminalia chebula* (Haritaki). The pharmacological studies of Haritaki were reviewed in this paper.

Keywords: Terminalia chebula, herbal drugs, Indian medicinal plants, pharmacological studies, haritaki, Kadukkai.

INTRODUCTION

According to the World Health Organization (WHO) more than 80% of the people living in the developing countries depend on traditional medicine for their primary health needs [1]. The traditional Indian system of medicines like Ayurveda, Siddha and Unani support the importance of medicinal plants to treat diseases [2].

Plants produce a wide array of bioactive principles and constitute a rich source of medicines [3]. The demand for plant-based therapeutics is increasing in both developing and developed countries due to the growing recognition that they are natural products, non narcotic, easily biodegradable, pose minimum environmental hazards, have no adverse side-effects and are easily available at affordable prices [4].

Terminalia chebula (haritaki) is used in traditional medicine due to the wide spectrum of pharmacological activities associated with the biological active chemicals present in this plant [5]. The name Haritaki in Sanskrit refers to the yellowish dye (harita) that contains the god Siva (Hari, i.e. the Himalayas) and that it cures (harayet) all the diseases [6].

Haritaki is a moderate sized tree grows widely in India, Myanmar, Bangladesh, Iran, Egypt, Turkey, China etc. [5]. It is found throughout India, chiefly in deciduous forests and areas of light rainfall, but occasionally also found in slightly moist forests, flowers appear from April-August and fruits ripen from October-January, fruit is drupe-like, 2—

4.5 cm long and 1.2–2.5 cm broad, blackish, with five longitudinal ridges [7].

SYNONYMS [8]

China – Zhang-Qin-Ge, Hezi

France – Myrobalan in dien

Germany - Myrobalane

India - {Assamese - Shilikha; Bengali - Haritaki; Gujarati-Hirdo, Himaja, Pulo-harda; Hindi - Harre, Harad, Harar; Kannnada - Alalekai; Kashmiri-Halela; Malayalam - Katukka; Marathi - Hirda, Haritaki, Harda, Hireda; Oriya - Harida; Punjabi - Hakeka, Harar; Sanskrit - Haritaki, Abhaya, Kayastha, Siva, Pathya; Tamil - Ammai, Amutam, Aritaki, Pethiyam, Varikkai; Telugu - Karakkaya; Urdu - Halela}

Srilanka – Aralu

Tibet – Harro

Types of haritaki:

In Siddha literature seven types of Haritaki are identified according to their geographical distribution namely Vijayan, Boodhana, Rogini, Abhyan, Amrutha, Boothagi and Sethagi.

Tastes found in the fruit of haritaki:

Among the six tastes except salt other five tastes are found in the fruit of haritaki namely outer skin-pungent, ridge-sour, seed-astringent, stem-bitter, endosperm-sweet.

Haritaki for tridosha:

To Indian system of medicine all diseases are due to the imbalance in tridosha or three bodily humors. Haritaki if taken with salt can cure kapha dosha, with

sugar can cure pitha dosha and when taken with ghee can cure vatha dosha.

Chemical constituents of haritaki

It contains high phenolic content, especially hydrolyzable tannins, anthraquinone, flavonol, carbohydrates, glucose and sorbitol [9], chebulic acid [10], chebulinic acid [11], ellagic acid, gallic acid [12], chebulagic acid [13] etc.,

Pharmacological studies of Haritaki:

Based on the chemical constituents present in haritaki several pharmacological investigations have been conducted in various in vivo and in vitro. Table 1 shows the summary of the findings of some of these pharmacological studies.

Table 1. Pharmacological studies on Haritaki

Pharmacological acitivity	Author/Reference	Extract type	organisms
Antibacterial	Kannan et al.,[4]	Ethanol extract	Salmonella typhi, Staphylococcus aureus, Bacillus subtilis etc.
	Malekzadeh et al.,[14]	Ether, alcoholic, water extract	Helicobacter pylori
Anticancer	Saleem et al.,[15]	70% of methanol	Human(MCF-7), mouse (S115) breast cancer cell lines etc.
Anticaries	Jagtap et al.,[16]	Aqueous extract	Streptococcus mutans
Anticonvulsant	Hogade Maheswar et al.,[17]	Ethanolic, chloroform, Petroleumether aqueous extract	Rats
Antidiabetic	Gandhipuram P et al.,[18]	Ethanol extract	Adult albino male rats
	Rao et al.,[19]	chloroform extract	Streptozotocin induced diabetic rats
Antifungal	Saheb Shinde et al.,[20]	Aqueous, alcoholic, ethyl acetate extract	Aspergillus niger, Aspergillus flavus, Alternaria alternata etc
	Vivek et al.,[21]	70% of methanol, ethylacetate, hexane, chloroform extract	Fusarium oxysporum, Phytophthora capsici, Fusarium solani etc.
Antimutagenic	Grover et al.,[22]	Chloroform, aqueous extract	Salmonella typhimurium
	Kaur et al.,[23]	Acetone,aqueous chloroform extract	Salmonella typhimurium
Antioxidant	Suchalata et al.,[24]	95% of ethanol extract	Adult male albino rats
	Chia-lin chang et al.,[25]	water, methanol & 95% of ethanol extract	Fermented products
Antiulcer	Raju et al.,[26]	Methanolic extract	Wistar albino male rats
Antiviral	Hongbo Ma et al.,[27]	Acetone extract	Swine influenza A virus
	Kim et al.,[28]	Aqueous extract	Hepatitis B virus
Cardioprotective	Suchalata et al.,[29]	95% of ethanol extract	Adult albino male rats
Cytotoxic	Kaur et al.,[30]	Acetone extract	Cancer cell lines
Immunodulatory	Vaibhav Aher et al.,[31]	Alcohol extract/	Male wistar rats
Radioprotective	Jagetia et al.,[32]	Aqueous extract	Rats
Wound healing	Manish PalSingh et al.,[33] Choudhary [34]	Hydroalcoholic extract 90% of ethanol extract	Induced diabetic rats Wistar albino rats
	Choudhary [34]	90% of enfanoi extract	vv istar albillo rats

Toxicological studies on Haritaki:

The inner seed of the fruit of Haritaki is toxic hence removed while preparing therapeutics based on

Haritaki. The acute toxicity study of the 50% alcoholic extract [35], subchronic toxicity study of both powder and water extract [36], acute and chronic toxicity studies of water extract given orally [37] from dried fruits of haritaki demonstrated no toxic effects in mice.

Traditional Uses of haritaki:

The fruit of haritaki has been extensively used in Thai traditional medicine for laxative, carminative, astringent, expectorant, and tonic effects [38]. It is routinely used as traditional medicine by tribes of Tamil Nadu to cure several ailments such as fever, cough, diarrhea, gastroenteritis, skin diseases, candidiasis, urinary tract infection and wound infections [6]. It is used commonly in many Ayurvedic preparations as diuretic and cardiotonic [16]. It is used to prevent aging and impart longevity, immunity [31]. It is reputed to cure blindness and it is believed to inhibit the growth of malignant tumors [15].

Haritaki as Kayakalpa:

It can be used as a kayakalpa (rejuvenating the body and the mind) if taken in evening in prescribed amounts. To act as kayakalpa it should be consumed according to the season as shown in Table 2.

Table 2. Consumption of Haritaki in 6 seasons

Season/Kaalam	Adjuvant
Illavenil (mildly sunny)	Honey
Muthuvenil (intense sunny)	Jaggery
Kar (Cloudy rainy)	Rock salt
Kuthir (Cold)	Sugar
Munpani (Early misty)	Dried ginger
Pinpani (Late misty)	Long pepper

Important preparations of haritaki:

Kadukkai mathirai:Iron deficiency

Moola kudori thylam: For bleeding piles.

Bhavana kadukkai:Anaemia

Kadukkai legiyam: constipation, gas trouble, ulcer

Triphala choornam:Digestive, blood purifier, reduces cholesterol, antioxidant, balances three body humors.

CONCLUSION

The traditional Indian system of medicines like Siddha, Ayurveda and Unani mainly uses medicinal plants for the management of diseases. Recognition for herbal therapeutics is gradually increasing in the world because they are safe and natural products. Extensive research has been carried out on medicinal plants attracting many more scholars to devote themselves for further study. Terminalia chebula (Haritaki) is one of the important herbal drug used for treating many diseases including some varieties of cancer. It is rich in pharmacological chemical constituents. Many investigations have been carried out based on its chemical constituents. A review on pharmacological studies on haritaki are presented in this

paper. These studies justify the claim on ancient literatures that haritaki called as amutham (nector) in tamil is a kayakalpa which balances the three bodily humors or doshas that are the main reasons for illness and stress that herbal medicines treats the cause of the disease rather than suppressing the symptoms.

REFERENCES

- 1. Sharma M, Pandey Govind P; Ethanomedicinal Plants for Prevention and treatment of tumors. International Journal of Green pharmacy, 2009; 3(1): 2-5.
- 2. Beusher N, Bodinet C, Neumann-Haefelin D, Marston A, Hostettmann K; Antiviral activity of African medicinal plants. J. Ethnopharmacol, 1994; 42(2): 101-109.
- 3. Badrul Alam B; Antioxidant, Antimicrobial and Toxicity studies of the Different Fractions of fruits of Terminalia belerica Roxb. Global Journal of Pharmacology, 2011;5(1):07-17.
- 4. Kannan P, Ramadevi SR, Hopper W; Antibacterial activity of *Terminalia chebula* fruit extract", African Journal of Microbiology Research, 2009; 3 (4):180-184.
- Prakash DV, Sree Satya N, Sumanjali Avanigadda S, Vangalapati M; Pharmacological review on Terminalia chebula. International Journal of Research in Pharmaceutical and Biomedical Sciences. 2012; 3 (2):679-683.
- 6. Dash B; Materia Medica of Ayurveda, B Jain Publisher, New Delhi, 1991.
- Flora of china, Terminalia chebula, Vol-13; 311-313.
- 8. The siddha pharmacopoeia of india, part I vol I, First edition, 99 100.
- 9. Creencia E, Eguchi T, Nishimura T, Kakinuma K; Isolation and structure elucidation of the biologically active components of *Terminalia chebula* Retzius (Combretaceae). KIMIKA ,1966;12:1-10.
- 10. Lee HS, Koo YC, Suh HJ, Kim KY and Lee KW; Preventive effects of chebulic acid isolated from *Terminalia chebula* on advanced glycation endproduct-induced endothelial cell dysfunction. Journal of Ethnopharmacol, 2010; 131(3): 567-574.
- 11. Quanbin Han, Jingzheng Song, Chunfeng Qiao, Lina Wong and Hongxi Xu; Preparative isolation of hydrolysable tannins chebulagic acid and chebulinic acid from *Terminalia chebula* by high-speed counter-current chromatography. J. Sep. Sci, 2006;29:1653-1657
- 12. Patel Madhavi G, Patel Vishal R, Patel Rakesh K; Development and Validation of Improved RP-HPLC method for Identification and Estimation of Ellagic and Gallic acid in

- Triphala churna. International Journal of ChemTech Research, 2010;2(3):1486-1493.
- 13. Bharat Reddy D, Reddy TCM, Jyotsna G, Sharan S, Priya N, Lakshmipathi V, Reddanna P; Chebulagic acid a COX–LOX dual inhibitor isolated from the fruits of *Terminalia chebula* Retz induces apoptosis in COLO-205 cell line. Journal of Ethnopharmacology,2009; 124(3): 506-512.
- Malekzadeh F, Ehsanifar H, Shahamat M, Levin M, Colwell RR; Antibacterial activity of black myrobalan (*Terminalia chebula* Retz) against Helicobacter pylori. International Journal of Antimicrobial Agents, 2001; 18(1): 85–88.
- 15. Saleem A, Husheem M, Harkonen P and Pihlaja K; Inhibition of cancer cell growth by crude extract and the phenolics of *Terminalia chebula* Retz fruit. Journal of Ethnopharmacology, 2002; 81(3): 327-336.
- 16. Jagtap AG and Karkera SG; Potential of the aqueous extract of *Terminalia chebula* as an anticaries agent. Journal of Ethnopharmacology,1999; 68(1-3): 299–306.
- 17. Hogade Maheshwar G, Deshpande S.V and Pramod H.J. Anticonsultant activity of *Terminalia chebula* retz. against mes and ptz induced seizures in rats, Journal of Herbal Medicine and Toxicology. 2010; 4(2): 123-126.
- 18. Gandhipuram P, Kumar S, Arulselvan P, Kumar DS, Subramanian SP; Anti Diabetic activity of fruits of Terninalia chebula on Streptozotocin induced Diabetic rats. Journal of Health Science, 2006; 52(3): 283-291.
- 19. Rao NK, Nammi S; Antidiabetic and renoprotective effects of the chloroform extract of *Terminalia chebula* Retz. Seeds in streptozotocin-induced diabetic rats. BMC Complement Altern Med, 2006; 6: 17.
- 20. Saheb S, More SM, Junne SB, Wadje SS; The Antifungal activity of five Terminalia species checked by paper disk method. International Journal of Pharma Research and Development, 2011; 3(2).
- 21. Vivek KB, Rahman A, Shukla S, Shukla S, YassirArafat SM, Hossain AM, Mehta A; In vitro kinetics and antifungal activity of various extracts of *Terminalia chebula* seeds against plant pathogenic fungi. Archives of Phytopathology and Plant Protection,2010; 43(8): 801-809.
- 22. Grover IS and Bala S; Antimutagenic activity of *Terminalia chebula* (myroblan) in Salmonella typhimurium. Indian Journal of Experimental Biology, 1992; 30(4): 339-341.
- 23. Kaur S, Arora S, Kaur K, Kumar S;The in vitro antimutagenic activity of Triphala an

- Indian herbal drug. Food Chem Toxicol, 2002; 40(4): 527-34.
- 24. Suchalatha S, Devi CS; Antioxidant activity of ethanolic extract of *Terminalia chebula* fruit against isoproterenol induced oxidative stress in rats. Indian Journal of Biochemistry and Biophysics, 2005; 42: 246-249.
- 25. Chia-Lin Chang and Che-San Lin; Development of antioxidant activity and pattern recognition extracts and its fermented products, Research Institute of Biotechnology, Hungkuang University. 2010.
- 26. Raju D, Ilango K, Chitra V, Ashish K; Evaluation of Anti-ulcer activity of methanolic extract of *Terminalia chebula* fruits in experimental rats. Journal of Pharmaceutical Science and Research, 2009; 1(3): 101-107.
- 27. Hongbo Ma, Yunpeng Diao, Danyu Zhao, Kun Li and Tingguo Kang; A new alternative to treat swine influenza A virus infection: extracts from *Terminalia chebula* Retz. African Journal of Microbiology Research, 2010; 4(6): 497-499.
- 28. Kim TG, Kang SY, Jung KK, Kang JH, Lee E, Han HM, Kim SH; Antiviral activities of extracts isolated from Terminalis chebula Retz., Sanguisorba officinalis L., Rubus coreanus Miq and Rheum palmatum L. against hepatitis B virus. Phytotherapy research, 2001; 15(8): 718-720.
- 29. Suchalatha S, Devi CS; Protective effect of *Terminalia chebula* against lysosomal enzyme alterations in isoproterenolinduced cardiac damage in rats. Experimental clinical cardiology, 2005; 10(2): 91-95.
- 30. Kaur S, Michael H, Arora S, Harkonen PL, Kumar S; The in vitro cytotoxic and apoptotic activity of Triphala--an Indian herbal drug. Journal of Ethnopharmacology, 2005; 97(1): 15–20.
- 31. Aher V, Wahi AK; Immunomodulatory activity of alcohol extracts of *Terminalia chebula* Retz combretaceae. Tropical Journal of Pharmaceutical Research, 2011;10(5): 567-575.
- 32. Jagetia GC, Baliga MS, Malagi KJ, Sethukumar KM; The evaluation of the radioprotective effect of Triphala (an ayurvedic rejuvenating drug) in the mice exposed to g-radiation. Phytomedicin, 2002; 9(2): 99–108.
- 33. Singh MP, Sharma CS; Wound healing activity of *Terminalia chebula* in experimentally induced diabetic rats. International Journal of Pharm Tech Research, 2009; 1(4): 1267-1270.
- 34. Choudhary GP; Wound healing activity of ethanolic extract of *Terminalia chebula* retzius. International Journal of Pharma and Bio Sciences, 2011; 2(1): 48 52.

- 35. Mokkhasmit M, Swatdimongkol K, Satrawaha P; Study on toxicity of Thai medicinal plants. Bull Dept Med Sci, 1971; 12(2/4):36-65.
- 36. Worasuttayangkurn L; The study of toxic effects of *Terminalia chebula* Retz. In mice [dissertation]. Thailand: Mahidol University, Faculty of Graduation Studies. 2001.
- 37. Panunto W, Jaijoy K, Lerdvuthisopon N, Lertprasertsuke N, Jiruntanat N,
- Soonthornchareonnon N, Sireeratawong S; Acute and chronic toxicity studies of the water extract from dried fruits of *Terminalia chebula* Rezt. in rats. International Journal of Applied Research in Natural Products, 2011; 3 (4):36-43.
- 38. Department of Medical Sciences, Ministry of Public Health. Thai Herbal Pharmacopoeia Volume II. Nonthaburi, Thailand. 2000; 71-89.