

Research Article

Study on Growth and Reproductive Behavior in Freshwater Catfish (*Heteropneustes fossilis*)

Suchitra Kumari

Research Scholar, Jai Prakash University, Chapra, Bihar

***Corresponding author**

Suchitra Kumari

Email: suchitra_kumari@rediffmail.com

Abstract: The changes in the gonadosomatic index (GSI) of a catfish, *Heteropneustes fossilis* was studied for relation between seasonal variation in growth and reproductive cycle. The fish has only one spawning season of short duration, running from July to August as indicated by the peaks of GSI and the diameter of oocytes and testicular lobules. Both males and females mature simultaneously. The minimum GSI for female was 0.74 ± 0.12 in resting phase and maximum was observed in the spawning phase (16.49 ± 1.70). The GSI for males was minimum in resting phase (0.087 ± 0.004) and maximum in spawning phase (2.02 ± 0.181).

Keywords: gonadosomatic index, *Heteropneustes fossilis*, Rotifera, and reproductive cycle

INTRODUCTION

The environmental changes greatly influence the production of eggs varies not only among different species but also within the same species. This depends upon the length and weight of the gonads. Maturity determination by gonadosomatic ratio has proved to be a significant tool in the life of fishes. Gonads undergoing regular seasonal cyclic changes in weight, particularly in females which help to indicate the spawning season [1]. The method of studying the spawning season is to follow the seasonal changes in gonadal weight in relation to body weight which is expressed as the gonadosomatic index. The Gonadosomatic index (GSI) is one of the important parameters of the fish biology, which gives the detail idea regarding the fish reproduction and reproductive status of the species and help in ascertaining breeding period of fish [2]. The gonadosomatic index measures the cyclic changes in gonad weight in relation to total fish weight, and can be used to determine spawning periods. The objective of the present work has to determine the maturity and spawning period influenced with different seasons in the fish *Heteropneustes fossilis* and dissected to remove the gonads.

METHODS AND MATERIALS

The study was carried out from January 2010 to December 2012. Healthy mature and immature

Heteropneustes fossilis numbering 280 (140 males and 140 females) were collected from Kushinagar hatchery station, Kushinagar (Uttar Pradesh). They were brought to the laboratory and were thoroughly washed with water and blotted completely to remove excess of water and each fish was weighed on electrical balance and dissected to remove the gonads. The weight of individual fish and its gonads were recorded and GSI was calculated using the formula.

$$\text{GSI} = \frac{\text{Weight of the gonad}}{\text{Weight of fish}} \times 100$$

RESULTS AND OBSERVATIONS

On the basis of the seasonal changes in the gonads, the annual cycle of *Heteropneustes fossilis* has been divided into following five phases:

1. Resting phase (November to January)
2. Preparatory phase (February to March)
3. Pre-spawning phase (April to June)
4. Spawning phase (July to August)
5. Post-spawning phase (September to October)

In the ovary, GSI and diameter of ova in various stages of development was calculated (Table 1 and 2) while in the testes, GSI and diameter of testicular lobules was calculated (Table 3 and 4).

Table-1: Seasonal gonadosomatic indices of female *Heteropneustes fossilis*.

Phases	Months	GSI	MEAN
Resting	November	0.69±0.10	0.74±0.12
	December	0.60±0.12	
	January	0.95±0.16	
Preparatory	February	1.26±0.05	1.89±0.24
	March	2.56±0.63	P < 0.01
Pre Spawning	April	3.43±0.89	10.42±0.79
	May	8.83±1.66	P<0.01
	June	19.00±1.02	
Spawning	July	21.98±1.80	16.49±1.70
	August	11.01±1.60	P<0.01
Post Spawning	September	1.30±0.36	3.08±0.34
	October	0.87±0.09	P<0.01

Table-2: Average oocyte diameter of the female fish during its reproductive phases.

Phases	Months	Oocyte diameter in um	MEAN
Resting	November	67.82 ± 1.42	70.83 ± 2.65µm
	December	66.22 ± 1.91	
	January	82.27 ± 1.90	
Preparatory	February	99.89 ± 0.47	111.9 ± 7.09 µm
	March	124.0 ± 3.44	
Pre Spawning	April	313.60 ± 4.73	502.60 ± 44.95 µm
	May	545.00 ± 2.46	
	June	649.10 ± 2.69	
Spawning	July	665.29 ± 5.38	583.80 ± 62.12 µm
	August	506.00 ± 3.81	
Post Spawning	September	99.29 ± 3.11	94.19 ± 2.84 µm
	October	89.08 ± 2.14	

Table-3: Seasonal gonadosomatic indices of male *Heteropneustes fossilis*.

Phases	Months	Oocyte diameter in um	Mean
Resting (Control)	November	0.081±0.008	0.087± 0.004
	December	0.051±0.012	
	January	0.120±0.003	
Preparatory	February	0.130±0.007	2.85±0.058
	March	0.440±0.109	P < 0.01
Pre-spawning	April	0.506±0.046	1.71± 0.08
	May	1.505±0.09	P < 0.01
	June	3.145±0.117	
Spawning	July	3.00±0.425	2.02± 0.181
	August	1.04±0.137	P < 0.01
Post-spawning	September	0.59±0.075	0.536±0.045
	October	0.089±0.015	P < 0.01

Table-4: Average lobules diameter of the testis during different phases of the reproductive cycle in *Heteropneustes fossilis*.

Phases	Months	Testicular lobule diameter in µm	Mean
Resting (Control)	November	54.71± 3.36	51.39 ± 2.00µm
	December	49.06 ± 4.05	
	January	49.91 ± 1.58	
Preparatory	February	68.84 ± 1.20	90.82± 6.98 µm
	March	92.80 ± 2.01	
Pre-spawning	April	137.20 ± 1.50	199.40 ± 15.53 µm
	May	186.10 ± 3.44	
	June	274.83 ± 4.55	
Spawning	July	252.60± 2.28	221.50 ± 25.28 µm
	August	130.40 ± 2.05	
Post-spawning	September	86.44 ± 1.99	77.94 ± 4.08µm
	October	69.44 2.65	

In the resting phase, ovaries are very small in size. They are thin, pinkish red in colour and have inconspicuous vascular supply. GSI is 0.74 ± 0.12 . The average diameter of oocytes in this phase is $70.83 \pm 2.65 \mu\text{m}$. Oocytes are smaller in size, somewhat triangular in shape with darkly stained cytoplasm, nucleus is large and round containing 1 to 2 nucleoli. The ovaries increase in weight and size in next phase. GSI in preparatory phase is 1.89 ± 0.24 . In the preparatory phase, average diameter of oocytes increase to $111.90 \pm 7.09 \mu\text{m}$. As ovaries approach maturity during pre-spawning period, their volume and vascular supply increases significantly. GSI suddenly shoots up to 10.42 ± 0.79 . The average oocyte diameter is $502.60 \pm 44.95 \mu\text{m}$. There is reduction in the inter-follicular space because oocytes increase in size due to yolk formation. The ovaries grow considerably in size occupying large area in the posterior half of the body during spawning phase. Ovaries become very large, fill the entire peritoneal cavity and contain fully matured oocytes laden with yolk. GSI becomes 16.49 ± 1.70 . The average diameter of oocytes is $583.80 \pm 62.13 \mu\text{m}$. However, few oocytes at peri-nucleolar and yolk vesicle stage are present in the peripheral area of the ovary. In the post-spawning phase, ovaries are reduced in volume and weight and have dull colour. Vascular supply is reduced. Some unspent ova are seen. There is a sharp decline in the GSI in this phase which is 3.08 ± 0.34 .

In males testes are very small and thread like in November to January. The vascular supply is inconspicuous. GSI value is 0.087 ± 0.004 . The average diameter of seminiferous lobules in resting phase is $51.39 \pm 2.00 \mu\text{m}$. In preparatory phase, testes increase in size and vascular supply also increases. GSI value in this phase is 0.285 ± 0.058 . The average diameter of testicular lobules increases to about $90.82 \pm 6.98 \mu\text{m}$. A prominent interlobular demarcation is observed in this phase. In pre-spawning phase, testes enlarge in volume. There is rapid increase in GSI in this phase. The GSI is 1.71 ± 0.08 , diameter of testicular lobule increase to about $199.40 \pm 15.53 \mu\text{m}$. In spawning phase, testes bulge out in the abdominal cavity occupying one third of the body cavity. Vascular supply increases and testis become red in colour. GSI value in this phase is about 2.02 ± 0.18 . The average lobule is about $221.50 \pm 25.28 \mu\text{m}$ in diameter. In postspawning phase, size of the testis decreases, GSI value also falls down to about 0.53 ± 0.045 . The average lobule diameter in this phase is much reduced to about $77.94 \pm 4.08 \mu\text{m}$.

DISCUSSION

The annual reproductive cycle in the females of *Heteropneustes fossilis* has been divided into five phases such as resting, preparatory, pre-spawning, spawning and post-spawning depending upon seasonal changes in the ovary, variations in GSI, oocyte diameter and testicular lobules diameter. GSI increases gradually from preparatory phase and the increase is statistically

significant ($p < 0.01$) both in pre-spawning and spawning phases with similar condition in *Oreochromis mossambicus* [3]. In *Garra mullya* also GSI exhibits increasing trend from February onwards and highest is in July which is the spawning phase. It decreases sharply from November to January [4]. During resting phase, the ovary is predominated by the immature oocytes. These are smaller in diameter ($70.83 \pm 2.65 \mu\text{m}$) with darkly stained ooplasm and large nuclei. GSI (0.74 ± 0.12) is lowest during this period. In preparatory phase, there is a gradual increase in GSI (1.89 ± 0.24) as maturation proceeds and new sets of oogonia grow to become oocytes at different stages. It is referred to such oocytes as reserve fund eggs and called them recruitment stock eggs [5]. These yolkless oocytes play an important role in supplying the eggs to be spawned in the following spawning season. Several other workers have discussed the origin of the yolkless oocytes [6]. The growth of oocytes is generally called first growth or primary growth of oocytes. This phase of growth does not seem to bring any marked influence on ovarian weight in *Heteropneustes fossilis*. During preparatory phase, the ovaries are predominated by oocytes which are at perinuclear stage with large nuclei and many nucleoli of various sizes. The extrusion of nucleoli into the ooplasm has attracted the attention of many workers. The significance of nucleolar extrusion is reported to be for the formation of proteins [7]. In *Heteropneustes fossilis*, many nucleoli of various sizes are seen in the oocytes which are at early perinucleolar stage. The size of nuclei decreases with developing stages of the oocytes. During pre-spawning phase, female *Heteropneustes fossilis* shows rapid increase in the GSI (10.42 ± 0.79). The ovaries are enlarged and various cytological changes are observed in the oocytes indicating rapid growth and maturation. The growth during this phase is mainly due to formation of yolk vesicles and deposition of yolk. Such changes in the pre-spawning phase have been reported in the ovaries of several teleostean species. In *Heteropneustes fossilis* during this phase, oocytes proliferate and all types of oocytes are visible except the matured ones. In the spawning phase, GSI of *Heteropneustes fossilis* attains a maximum peak (16.49 ± 1.70). The ovaries during spawning phase are filled with yolk laden oocytes. Very few immature oocytes are also visible along the peripheral region of the ovary. At the end of this phase, the ovary decreases in weight not only due to ovulation or discharge of the eggs, but also due to degeneration of oocytes which is referred to as atresia. Similar condition is also reported in many other teleost species such as *Clarias batrachus* [8]. The annual cyclic changes in the males of *Heteropneustes fossilis* are noticed mainly due to maturation of testis. The size, shape, colour and length of the testes undergo variations during different reproductive phases of the life cycle. Similar changes have also been reported in other teleosts by several other workers [9]. The testes of this fish attain maximum weight between July to August corresponding to

spawning season and they show maximum GSI (2.02 ± 0.181) during this phase. It is then followed by a rapid decline. The GSI values during resting phase is very low (0.087 ± 0.004), the seminiferous lobules are small and they are mostly predominated by primary spermatogonia. There is a gradual increase in the GSI during preparatory phase (0.285 ± 0.058). Increase in the GSI is very rapid in prespawning phase (1.71 ± 0.08). In this phase, the testes are predominated by spermatocytes, spermatids and spermatozoa. GSI attains its maximum peak during the spawning phase in the month of July (2.02 ± 0.181). From post-spawning phase onwards, there is a sudden fall in the GSI which becomes 0.53 ± 0.045 . This low GSI in the post-spawning phase is due to the discharge of milt. In this phase, the wall of seminiferous lobules ruptures and spermatozoa are released out. Belsare, has reported that the testes of *Gasterosteus aculeatus* remain mature at any time of the year but their functional maturity is attained only in the breeding season (April-May) [10]. The variation in view may be because of different environmental factors which play an important role in gonadal maturation and development. The maturation of testes in *Gasterosteus aculeatus* is because of high temperature. From April to July maximum gonadal activity is reported for *Heteropneustes fossilis* [11]. The present investigation concluded that the GSI of catfish *Heteropneustes fossilis* is maximum during spawning season whereas decreases during post-spawning season.

REFERENCES

1. Dadzie SF, Abou-Seedo, Al-Shallal T; Reproductive biology of the silver Pomfret *Pampus argentens* (Euphrasen) in Kuwait water, Journal of Applied Ichthyology, 2000; 16: 247-253.
2. Shankar DS, Kulkarni RS; Somatic condition of the fish, *Notopterus notopterus* (pallas) during phases of the reproductive cycle. Journal of Environmental Biology, 2005; 26(1):49-53
3. Pathan JGK, Baile VV; Changes in protein in relation to the reproductive cycle in the male Tilapia, *Oreochromis mossambicus*. National Journal of Life Science, 2005; 1: 14-18.
4. Khan EA, Mehrotra PH; Variations of liver protein and RNA in relation to egg maturation in a hill stream teleost *Garra mullaya* (Sykes). Journal of Reproductive Biology and Comparative Endocrinology, 1999; 3(1): 47-52.
5. Hickling CF; The seasonal cycle in the Cornish pichard, *Sardina pilchardus*. (Waibum). Journal of Marine Biological Association U.K. , 1945; 26: 115-138.
6. Swarup H; The reproductive cycle and development of the gonads in *Gasterosteus aculeatus* (L.). Proceeding of Zoological Society Bengal, 1958; 11: 47-61.
7. Khanna S; An introduction to fisheries, Central book depot, Allahbad. 1956.
8. Lehri GK; The annual cycle in the ovary of the cat fish, *Clarias batrachus*, L. Acta.Anatomica, 1968; 69: 105-124.
9. Ruby SM, McMillan DB; Cyclic changes in the testis of the Brook Stickleback *Eucalia inconstans* (Kirtland). Journal of Morphology, 1970; 131: 447-466.
10. Belsare DK; Seasonal changes in the ovary of *Ophicephalus punctuators* (Bloch) Indian, Journal of Fisheries, 1962; 9: 149-157.
11. Ghosh A, Kar AB; Seasonal changes in gonads of the common Indian cat fish *Heteropneustes fossilis* (Bloch), Proceeding of Zoological Society. Bengal, 1952; 5: 29-49.