

Research Article

In vitro Anti fungal effect of *Allium sativum* and *Thymus vulgaris* on two pathogenic fungi

Saman Seyedgholizadeh^{*1}, Mohammad Hosein Sadeghi Zali², Saber Yousefi³, Amin Hashempour¹, Nazereh Hashempour¹

¹Young Researcher and Elites club, Islamic Azad University, Urmia Branch, Urmia Iran

²Department of Bacteriology, Veterinary Faculty, Islamic Azad University, Urmia Branch, Urmia, Iran

³Department of Microbiology, Virology and Immunology, Urmia University of Medical Sciences, Urmia, Iran

*Corresponding author

Saman Seyedgholizadeh

Email: saman.seyedgholizade@gmail.com

Abstract: Infectious diseases accounts for high proportion of health problems in the developing countries including India. Microorganisms have developed resistance to many antibiotics and as a result, immense clinical problem in the treatment of infectious diseases has been created. Susceptibility tests were performed by the disc diffuse on method of Bauer et al. In this study garlic has a fungicidal effect on 2 strains of fungi but thyme have a fungicidal effect on *Mucor Hymalys* and inhibit from growth of *Candida albicans*. This study showed that *Allium sativum* and *Thymus vulgaris* has antifungal effect on *Mucor Hymalys* and *Candida albicans*.

Keywords: In vitro, Anti fungal, *Allium sativum*, *Thymus vulgaris*, pathogenic fungi.

INTRODUCTION

Medicinal plants have served through ages, as a constant source of medicaments for the exposure of a variety of diseases. The history of herbal medicine is almost as old as human civilization. The plants are known to provide a rich source of botanical anthelmintics, antibacterials and insecticides[1-3].

Infectious diseases accounts for high proportion of health problems in the developing countries including India. Microorganisms have developed resistance to many antibiotics and as a result, immense clinical problem in the treatment of infectious diseases has been created [4].

There are alarming reports of market opportunistic fungal infections[5]. The infections caused by opportunistic fungi are included under new spectrum of fungal pathogens. Such fungi were earlier reported from various plants as pathogens. But now they are known to cause disease in human beings. There is an increasing awareness amongst clinicians and microbiologists pertaining to importance of infection caused by opportunistic fungi[6,7].

In the People's Republic of China, commercial *A. sativum* extracts are widely used to treat patients with systemic fungal infections [8-10].

Numerous studies have documented the antifungal [11-12] and antibacterial[13-14] effects of plant essential oils. Screening experiments with 13–52 essential oils and major active components against 5–25 microorganisms[16-17]. It also reported that thyme, clove, cinnamon, bay, oregano, garlic and lemongrass to be some of the best broad spectrum candidates for inhibition of food-borne pathogens and spoilage organisms [18].

The aim of this study is In vitro Anti fungal effect of *Allium sativum* and *Thymus vulgaris* on two pathogenic fungi.

MATERIAL AND METHODS

Collection of plant material

Thymus vulgaris were collected on March 2014 from around of Urmia city. *Allium sativum* were prepared from Rasht city on June 2014.

Culture and Maintenance of Fungi

Fungi were prepared from Iranian Research Organization for Science and Technology (IROST). After preparation this agent was sub cultured. Fungi were cultured on Saburd dextrose Agar media.

Table 1: characteristics of Fungi

S. No	Name	Type	PTCC No
1	<i>Mucor Hymalys</i>	Fungi	PTCC5292
2	<i>Candida albicans</i>	Yeast	PTCC5027

Preparation of plant essential oil

After collecting plants peeling and dry was done. After drying, essential oil with Clevenger apparatus method was done. 100gram from dry mater added to 400cc sterile water and boiled on heater for 4 hours.

Media Preparation and Its Sterilization

For the plates we prepared Saburd dextrose media. All the media prepared was then sterilized by autoclaving the media at (121°C) for 20 min.

Disc diffusion method:

Susceptibility tests were performed by the disc diffuse on method of Bauer et al [19].

RESULTS

In these tests, the effect of the essential oil on the fungi, garlic and thyme essential oils showed antifungal properties.

Table 2: Diameter of inhibition zone from garlic essential oil on ten day of the following described fungi

Fungi	Replication	First day	Third day	Tenth day
Mucor Hymalys	Rep1	Not growth	Not growth	Not growth
	Rep2	Not growth	Not growth	Not growth
	Rep3	Not growth	Not growth	Not growth
	Rep4	Not growth	Not growth	Not growth
Candida albicans	Rep1	Not growth	Not growth	Not growth
	Rep2	Not growth	Not growth	Not growth
	Rep3	Not growth	Not growth	Not growth
	Rep4	Not growth	Not growth	Not growth

Table 3: Diameter of inhibition zone from thyme essential oil on ten day of the following described fungi

Fungi	Replication	First day	Third day	Tenth day
Mucor Hymalys	Rep1	Not growth	Not growth	Not growth
	Rep2	Not growth	Not growth	Not growth
	Rep3	Not growth	Not growth	Not growth
	Rep4	Not growth	Not growth	Not growth
Candida albicans	Rep1	6 cm	6 cm	4.5 cm
	Rep2	5 cm	5 cm	4.5 cm
	Rep3	6.5 cm	6.5 cm	4.5 cm
	Rep4	5 cm	5 cm	3.5 cm

Table 4: Diameter of colony in the following described fungi as control sample on ten day

Fungi	Replication	First day	Third day	Tenth day
Mucor Hymalys	Rep1	Not growth	5 cm	Complete growth
	Rep2	Not growth	5 cm	Complete growth
	Rep3	Not growth	5 cm	Complete growth
	Rep4	Not growth	5 cm	Complete growth
Candida albicans	Rep1	Not growth	4 cm	Complete growth
	Rep2	Not growth	4.5 cm	Complete growth
	Rep3	Not growth	4.5 cm	Complete growth
	Rep4	Not growth	4 cm	Complete growth




Fig 1: Average of Diameter of inhibition zone from thyme essential oil on ten day (with cm)




Fig 2: Average of Diameter of colony in control sample on ten day (with cm)

DISCUSSION

In this study garlic has a fungicidal effect on 2 strains of fungi but thyme have a fungicidal effect on Mucor Hymalys and inhibit from growth of Candida albicans. This study showed that Allium sativum and Thymus vulgaris has antifungal effect on Mucor Hymalys and Candida albicans. Sunita Bansod and Mahendra Rai in 2008 showed that garlic has antifungal effect [7]. In 2001 Zafar Iqbal et al proved Anthelmintic Activity of Allium sativum [3]. In another study Maryam Omidbeygi et al in proved antifungal activity of thyme on *Aspergillus flavus*[18]. Despite these studies, conclude that Allium sativum and Thymus vulgaris has antifungal effect on 2 strains of fungi.

Acknowledgment: Authors are thankful to Young Researcher and Elite club (Islamic Azad university Urmia branch) for providing financial assistance in the form of a major research project. Special thanks to Dr. Ashkan Khoda bandeloo, Dr. Mahmoud pour Yousef, Mr.Reza Delshad and Mr. Amin khorsandian for helping me with my experiments.

REFERENCE

1. Satyavati GV, Raina MK, Sharma M; Medicinal Plants of India. Vol. I, pp: 201–06. Indian Council of Medical Research, New Delhi. 1976.

- Lewis WH, Elvin- Lewis MPH; Medicinal Botany Plants Affecting Man’s Health. John Wiley and Sons, New York. 1977.
- Iqbal Z, Nadeem QK, Khan MN, Akhtar MS, Waraich FN; In vitro anthelmintic activity of Allium sativum, Zingiber officinale, Curcubita mexicana and Ficus religiosa. International Journal of Agriculture and Biology, 2001; 3(4):454-457.
- Davies J; Inactivation of antibiotic and the dissemination of resistance genes. Sci, 1994; 264: 375-382.
- Singh N; Trends in the epidemiology of opportunistic fungal infections: predisposing factors and the impact of antimicrobial use practices. Clinical Infect Diseases, 2001; 33(10):1692-1696.
- Alex LA, Korch C, Selitrennikoff CP, Simon MI; COS1, a two-component histidine kinase that is involved in hyphal development in the opportunistic pathogen Candida albicans. Proc. Natl. Acad. Sci. USA, 1998; 95: 7069-7073.
- Sunita B, Rai M; Antifungal Activity of Essential Oils from Indian Medicinal Plants Against Human Pathogenic Aspergillus fumigatus and A. niger, World Journal of Medical Sciences , 2008; 3(2): 81-88.

8. Delaha E C, Garagusi VF; Inhibition of mycobacteria by garlic extract (*Allium sativum*). *Antimicrobial agents and chemotherapy*, 1985; 27(4):485-486.
9. Khan Z K, Katiya R; Potent antifungal activity of garlic (*Allium sativum*) against experimental murine disseminated cryptococcosis. *Pharmaceutical biology*, 2000; 38(2): 87-100.
10. Davis LE, Shen JK, Cai Y; Antifungal activity in human cerebrospinal fluid and plasma after intravenous administration of *Allium sativum*, *Antimicrob. Agents Chemother*, 1990; 34(4):651.
11. Aligiannis N, Kalpoutzakis E, Chinou IB, Mitakou S, Gikas E, Tsiaropoulos A; Composition and antimicrobial activity of the essential oils of *Wve taxa of Sideritis* from Greece. *Journal of Agricultural and Food Chemistry*, 2001; 49:811–815.
12. Thompson DP; Fungitoxic activity of essential oil components on food storage fungi. *Mycologia*, 1981; 81:151–153.
13. Canillac N, Mourey A; Antibacterial activity of the essential oil of *Picea excelsa* on *Listeria*, *Staphylococcus aureus* and coliform bacteria. *Food Microbiology*, 2001; 18:261–268.
14. Lachowicz KJ, Jones GP, Briggs DR, Bienvenu FE, Wan J, Wilcock A, et al.; The synergistic preservative effects of the essential oils of sweet basil (*Ocimum basilicum* L.) against acid-tolerant food microXora. *Letters in Applied Microbiology*, 1998; 26:209–214.
15. Conner DE, Beuchat LR; Effects of essential oils from plants on growth of food spoilage yeasts. *Journal of Food Science*, 1984; 49:429– 434.
16. Deans SG, Ritchie G; Antibacterial properties of plant essential oils. *International Journal of Food Microbiology*, 1987; 5:165–180.
17. Dorman HJD, Deans SG; Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *Journal of Applied Microbiology*, 2000; 88:308–316.
18. Omidbeygi M , Barzegar M, Hamidi Z, Naghdibadi H; Antifungal activity of thyme, summer savory and clove essential oils against *Aspergillus Xavus* in liquid medium and tomato paste, *Food Control*, 2007; 18:1518–1523.
19. Bauer AW, Kirby WM, Sherris JC, Turck M; Antibiotic susceptibility testing by standardized single disc method. *Am. J. Clin Pathol*, 1996; 44: 493-496.