

## Bio-Monitoring of Metal Trace Elements by Epiphytic Lichen in the Bordj Bou Arreridj Area, East of Algeria

Adjiri Fatima<sup>1</sup>, Ramdani Messaoud<sup>1\*</sup>, Lograda Takia<sup>1</sup> and Chalard Pierre<sup>2</sup>

<sup>1</sup>Laboratory of Natural Resource Valorisation, SNV Faculty, Setif 1 University, 19000 Setif, Algeria

<sup>2</sup>SIGMA Clermont, Campus des Cezeaux, CS 20265, 63178 Aubière cedex France

### Original Research Article

\*Corresponding author  
Ramdani Messaoud

#### Article History

Received: 04.02.2018

Accepted: 13.02.2018

Published: 28.02.2018

#### DOI:

10.36347/sajb.2018.v06i02.012



**Abstract:** The objective of this study is to use a bio-sensor in order to assess environmental by metal trace elements (MTE) contamination in the Bordj Bou Arreridj region, East of Algeria. The sampling of biological material from 32 locations was performed using a natural bioaccumulator represented by a lichenic species; *Xanthoria parietina*. Samples were taken and ten heavy metals (Fe, Cd, Co, Cr, Cu, Mn, Ni, Pb, Sb and Zn) were selected for analysis. The metal content recorded in various sampled sites were found greatly fluctuating and exceeded standard values with the exception of Pb at the Kseur and Ghilassa locations. The level of Fe was very high in all sampled locations with an average of  $(43184.4 \pm 16373.7 \text{ mg/Kg})$ . This important accumulation of MTE in thalli was correlated to road and rail traffic as well as to industrial activities. These results show the tolerance of *X. parietina* to metallic stresses and therefore its effectiveness as a bio-sensor in the bio-monitoring programs assessing air quality.

**Keywords:** Metal trace elements; bio-monitoring; bio-accumulation; *Xanthoria parietina*; Bordj Bou Arreridj; Algeria.

### INTRODUCTION

The level of urban air pollution is due to the intensive development of industry, road traffic, population density and urbanization, leading to the degradation of air quality [1, 2]. Bio-monitoring of air quality is the use of responses, at all levels, of biological organisms to predict and reveal environmental alterations and to follow its evolution [3]. Bio-monitoring is a very effective operational tool [2, 4-6].

It is a complementary method to the physico-chemical analyzes and highlights the impregnation of the environment by the metallic trace elements (MTE) [7].

The MTE, known as heavy metals or toxic metals, are widely reputed pollutants emitted into the atmosphere by a large number of natural and anthropogenic sources [8-10]. Epiphytic cryptogams are gaining increasing importance in environmental bio-monitoring because they can absorb and accumulate elements in their systems [11-12].

Bio-indication studies involving lichens provide much information on the qualitative status of the environment and are useful for assessing the long-term effects of bio-monitoring programs [2]. Lichens are among the most commonly used cryptogamic organisms in the bio-monitoring of MTE in air [2, 13-14]. Foliaceous lichens are highly tolerant to metal trace

elements and reflect the history of the environment [11-12, 15]. *Xanthoria parietina* is the most widely used lichen in bio-monitoring studies [2, 13, 16-19].

The aim of this study is to assess MTE environmental contamination by drawing up a map of the distribution of pollution in the 32 municipalities of the Bordj Bou Arreridj province (BBA), East of Algeria, using the lichen *Xanthoria parietina*.

### MATERIALS AND METHODS

#### Sample Collection

The foliaceous lichen *Xanthoria parietina*, widespread in the studied areas and widely used in similar studies in several countries, has been selected as a bio-accumulator [16]. Samples were collected in 34 locations spread throughout the Bordj Bou Arreridj (BBA) province (Figure 1).




Fig-1: Sampled populations of *Xanthoria parietina*

Samples were collected avoiding the usage of tools or containers which may contaminate them (we avoided steel tools or stainless steel containers whose walls contain pigments based on trace elements such as PVC). Each sampled site consists of an area of a maximum 100 m<sup>2</sup>, located near high traffic roads and secondary roads. The geographical coordinates of each sampled area were noted using a GPS (Table 1). The BBA province is characterized by a semi-arid continental climate, with hot dry summers and severe winters. Rainfall is inadequate and irregular both in time and space.

#### Samples treatment

Lichen samples were digested in the laboratory in aseptic conditions, using a mixture of HNO<sub>3</sub>/HF/H<sub>2</sub>O in Teflon containers [20-21]. A solution of our samples is prepared using the grinded solid material to which 10 ml of 40% hydrofluoric acid (HF) and 3 ml of 70% perchloric acid (HClO<sub>4</sub>) were added. Evaporation takes place on a hotplate at 160°C. After a quasi-total evaporation, 1 ml of 65% nitric acid (HNO<sub>3</sub>) and 10 ml of distilled water were added. The samples are then left for 30 min at 4°C in the refrigerator.

The dissolution of the residue is performed by placing the samples on a hotplate at 60°C for 1h. The resulting mixture is transferred to a 100 ml flask for filtration, adjusting the volume with distilled water.

**Table-1: Geographic coordinates of the sampled areas**

| Code | Locations          | Latitude North (N) | Longitude East (E) | Altitude (m) |
|------|--------------------|--------------------|--------------------|--------------|
| 1    | El Main            | 36° 21' 59.907"    | 4° 43' 56.866"     | 983          |
| 2    | Tassameurt         | 36° 15' 52.732"    | 4° 48' 11.329"     | 629          |
| 3    | Tefreg             | 36° 18' 39.333"    | 4° 42' 30.243"     | 1131         |
| 4    | Djaafra            | 36° 17' 33.561"    | 4° 39' 42.894"     | 1335         |
| 5    | Khelil             | 36° 10' 1.578"     | 5° 1' 25.946"      | 964          |
| 6    | Bordj Zemoura      | 36° 15' 59.677"    | 4° 50' 21.156"     | 913          |
| 7    | Ouled Dahmane      | 36° 13' 41.188"    | 4° 46' 50.138"     | 1250         |
| 8    | Colla              | 36° 15' 32.299"    | 4° 39' 43.326"     | 1250         |
| 9    | Teniet En Nasr     | 36° 13' 34.759"    | 4° 35' 21.930"     | 1250         |
| 10   | Ouled Sidi Brahim  | 36° 13' 52.071"    | 4° 20' 31.498"     | 580          |
| 11   | Ain-Taghrouit      | 36° 7' 31.180"     | 5° 4' 44.572"      | 907          |
| 12   | Bir Kasdali        | 36° 9' 0.853"      | 5° 1' 16.698"      | 955          |
| 13   | Sidi Embarek       | 36° 6' 13.896"     | 4° 54' 29.289"     | 1011         |
| 14   | Hasnaoua           | 36° 8' 17.610"     | 4° 48' 13.622"     | 993          |
| 15   | Medjana            | 36° 7' 55.621"     | 4° 40' 20.503"     | 1052         |
| 16   | Mansoura           | 36° 4' 16.060"     | 4° 28' 21.561"     | 807          |
| 17   | El M'hir           | 36° 6' 57.042"     | 4° 23' 2.713"      | 557          |
| 18   | Haraza             | 36° 8' 20.085"     | 4° 12' 52.610"     | 1201         |
| 19   | Ben Daoud          | 36° 3' 59.274"     | 4° 8' 42.662"      | 1050         |
| 20   | Tixter             | 36° 2' 49.300"     | 5° 5' 8.592"       | 947          |
| 21   | Ain Tesra          | 36° 2' 17.28"      | 5° 0' 75.81"       | 1039         |
| 22   | El Ansseur         | 36° 1' 47.470"     | 4° 53' 80.124"     | 980          |
| 23   | Bordj Bou Arreridj | 36° 4' 22.548"     | 4° 41' 31.452"     | 901          |
| 24   | El Achir           | 36° 3' 10.771"     | 4° 37' 54.631"     | 967          |
| 25   | Ras El Oued        | 35° 56' 58.912"    | 5° 2' 2.774"       | 1078         |
| 26   | Belimour           | 35° 58' 40.738"    | 4° 52' 25.165"     | 937          |
| 27   | El Hamadia         | 35° 58' 24.081"    | 4° 44' 29.356"     | 819          |
| 28   | Ksour              | 36° 59' 29.014"    | 4° 30' 39.281"     | 1250         |
| 29   | Ouled Brahem       | 35° 51' 53.737"    | 5° 4' 55.452"      | 1259         |
| 30   | Bordj Ghédir       | 35° 55' 8.500"     | 4° 53' 22.47"      | 1387         |
| 31   | Taglait            | 35° 50' 35.347"    | 4° 57' 10.404"     | 1434         |
| 32   | Ghilassa           | 35° 51' 35.031"    | 4° 54' 39.110"     | 1171         |

#### Analytical methods for MTE concentration measurements in lichens

The concentrations of the following elements; Fe, Cd, Co, Cu, Cr, Mn, Ni, Pb, Sb and Zn, were determined by Atomic Absorption Spectro-photometry with Flame (AASF). There are no established standards

of trace elements concentration in lichens [22]. To interpret the results of each element studied we used as a standard reference, values of the European Commission BCR information CRM-482 (Table 2) [23].

**Table-2: Certified CRM-482 values of trace elements (mg/kg) using AASF**

| MTE       | Symbols | Certified values | Uncertainty |
|-----------|---------|------------------|-------------|
| Lead      | Pb      | 40.9             | 1.40        |
| Copper    | Cu      | 7.03             | 0.19        |
| Cadmium   | Cd      | 0.56             | 0.02        |
| Zinc      | Zn      | 100.6            | 2.20        |
| Nickel    | Ni      | 2.47             | 0.07        |
| Chrome    | Cr      | 4.12             | 0.15        |
| Cobalt    | Co      | 0.32             | 0.03        |
| Iron      | Fe      | 804              | 160         |
| Manganese | Mn      | 33.0             | 0.50        |
| Antimony  | Sb      | 0.35             | 0.09        |

**Statistical Analysis**

Data were first subjected to Principal Components Analysis (PCA) to examine the relationship among the trace elements and the bio-accumulation by lichens, and also the relation between the presence of these elements and the pollution. Cluster analysis (UPGMA) was carried out on the original variables and on the Manhattan Distance Matrix to look for hierarchical associations among the

elements and the locations. Statistical analyses were carried out using STATISTICA 10 software.

**RESULTS**

The concentrations of metal elements (Fe, Cd, Co, Cr, Cu, Mn, Ni, Pb, Sb and Zn) accumulated in the *Xanthoria parietina* were determined by Atomic Absorption Spectro-photometry with Flame (Table 3).

**Table-3: MTE concentrations in *Xanthoria parietina* (mg/kg dry weight)**

| Stations                  | Pb   | Mn    | Fe      | Cu   | Sb    | Cd   | Co     | Zn     | Ni     | Cr     |
|---------------------------|------|-------|---------|------|-------|------|--------|--------|--------|--------|
| 1 El Main                 | 110  | 800   | 65900   | 78   | 158   | 3    | 638.04 | 611.37 | 99.93  | 380.95 |
| 2 Tassameurt              | 120  | 1300  | 68500   | 80   | 194   | 3    | 658.15 | 348.55 | 77.16  | 321.09 |
| 3 Tefreg                  | 96   | 800   | 46400   | 56   | 203   | 4    | 605.36 | 376.47 | 101.32 | 376.87 |
| 4 Djaafra                 | 89   | 1300  | 72800   | 63   | 187   | 3    | 692.5  | 614.25 | 93.89  | 406.8  |
| 5 Khelil                  | 133  | 500   | 36600   | 35   | 219   | 3    | 347.3  | 437.36 | 3.72   | 202.72 |
| 6 Bordj Zemoura           | 105  | 400   | 29000   | 51   | 163   | 3    | 416.84 | 323.88 | 42.3   | 274.83 |
| 7 Ouled Dahmane           | 111  | 400   | 45100   | 36   | 253   | 4    | 668.2  | 484.19 | 52.99  | 334.69 |
| 8 Colla                   | 60   | 700   | 32900   | 33   | 185   | 3    | 674.91 | 433.4  | 98.07  | 387.76 |
| 9 Teniet En Nasr          | 81   | 500   | 47800   | 55   | 145   | 2    | 673.23 | 482.75 | 95.75  | 406.8  |
| 10 Ouled Sidi Brahim      | 110  | 1300  | 71400   | 68   | 167   | 3    | 714.29 | 514.28 | 105.51 | 395.92 |
| 11 Ain Taghrouit          | 206  | 500   | 40700   | 28   | 212   | 5    | 254.29 | 916.87 | 30.68  | 127.89 |
| 12 Bir Kasdali            | 114  | 300   | 36000   | 36   | 210   | 4    | 389.19 | 372.87 | 26.96  | 201.36 |
| 13 Sidi Embarek           | 115  | 1300  | 73000   | 67   | 162   | 3    | 579.39 | 497.7  | 46.01  | 293.88 |
| 14 Hasnaoua               | 136  | 800   | 48900   | 65   | 238   | 3    | 527.44 | 408.72 | 90.17  | 359.18 |
| 15 Medjana                | 173  | 700   | 46400   | 87   | 226   | 4    | 600.34 | 429.97 | 95.28  | 390.48 |
| 16 Mansoura               | 90   | 600   | 43800   | 52   | 183   | 3    | 581.06 | 469.96 | 123.63 | 434.01 |
| 17 El M'hir               | 96   | 500   | 46000   | 67   | 178   | 3    | 811.48 | 438.8  | 100.4  | 472.11 |
| 18 Haraza                 | 93   | 400   | 50200   | 66   | 170   | 3    | 772.1  | 360.8  | 118.52 | 443.54 |
| 19 Ben Daoud              | 71   | 900   | 58000   | 33   | 186   | 3    | 814.83 | 570.66 | 133.4  | 428.57 |
| 20 Tixter                 | 169  | 400   | 32800   | 24   | 243   | 5    | 283.62 | 983.52 | 16.27  | 144.22 |
| 21 Ain Tesra              | 86   | 400   | 30100   | 43   | 164   | 3    | 434.44 | 471.04 | 7.44   | 229.93 |
| 22 El Anseur              | 99   | 1100  | 63200   | 76   | 190   | 3    | 488.06 | 539.67 | 43.69  | 292.52 |
| 23 Bordj Bou Arreridj     | 68   | 400   | 17000   | 23   | 93    | 3    | 601.17 | 549.4  | 38.58  | 303.4  |
| 24 El Achir               | 120  | 600   | 40800   | 39   | 194   | 3    | 711.77 | 390.34 | 107.83 | 378.23 |
| 25 Ras El Oued            | 93   | 400   | 29500   | 23   | 175   | 4    | 297.86 | 431.96 | 3.3    | 161.9  |
| 26 Belimour               | 66   | 500   | 23500   | 38   | 182   | 3    | 633.01 | 347.65 | 54.38  | 303.4  |
| 27 El Hamadia             | 58   | 900   | 42100   | 37   | 174   | 3    | 572.69 | 480.41 | 88.31  | 348.3  |
| 28 Ksour                  | 6    | 900   | 24200   | 27   | 226   | 3    | 591.12 | 451.23 | 95.75  | 386.39 |
| 29 Ouled Brahem           | 95   | 700   | 45100   | 51   | 190   | 3    | 380.81 | 364.77 | 4.65   | 165.99 |
| 30 Bordj Ghédir           | 91   | 500   | 37100   | 62   | 233   | 3    | 457.06 | 364.77 | 32.07  | 227.21 |
| 31 Taglait                | 67   | 400   | 30100   | 23   | 171   | 3    | 426.06 | 406.92 | 19.99  | 239.46 |
| 32 Ghilassa               | 23   | 300   | 7000    | 72   | 102   | 3    | 510.68 | 377.56 | 30.68  | 209.52 |
| <i>Average</i>            | 98,4 | 671.9 | 43184.4 | 49.8 | 186.8 | 3.3  | 556.5  | 476.6  | 65     | 313.4  |
| <i>SD</i>                 | 39,5 | 312.4 | 16373.7 | 19.3 | 35.7  | 0.6  | 152.6  | 145.9  | 40.3   | 97.3   |
| <i>CV%</i>                | 40,1 | 46,5  | 37,9    | 38,8 | 19,1  | 18,2 | 27,4   | 30,6   | 62,0   | 31,1   |
| <i>Min</i>                | 6    | 300   | 7000    | 23   | 93    | 2    | 254.3  | 323.9  | 3.3    | 127.9  |
| <i>Max</i>                | 206  | 1300  | 73000   | 87   | 253   | 5    | 814.8  | 983.52 | 133.4  | 472.1  |
| <i>Certified standard</i> | 40.9 | 33    | 804     | 7.03 | 0.35  | 0.6  | 0.32   | 100.6  | 2.47   | 4.12   |

The MTE concentrations in the 32 sampled sites are found highly variable. It is noted that the (Fe) concentrations are too high in the thalli. The high Fe rate hid the variability of the rest of the elements in the

analysis (Figure 2a), this element was removed and a second analysis was performed to reveal the variability of the MTE in *X. parietina* (Figure 2b).




Fig-2: Variability of MTE concentrations in *X. parietina* thalli

The Fe element exhibits the highest mean concentration in *X. parietina* samples with (43184.4 ± 16373.7 mg / kg), while Cd is the least abundant in the thalli (3.3 ± 0.6 mg / kg). The metals concentrations analyzed are distributed in the following order: Fe > Mn > Co > Zn > Cr > Sb > Pb > Ni > Cu > Cd.

Fe showed the greatest variation followed by Mn (671.9 ± 312.4mg/kg). Concentration levels of Co, Zn and Cr are relatively close and exceed standard values. The mean concentrations of Sb and Pb found in rural and urban communes are relatively similar (186.8 ± 35.7 mg / kg) and (98.4 ± 39.5 mg / kg), while

concentrations of Ni and Cu are the lowest in order of appearance after the Cd.

The three-dimensional spatial projection of the stations, based on the first three axes resulting from the PCA, shows the separation of populations into two more-or-less distinct groups, thus isolating three Ghilassa, Ain Taghroute and Tixter areas (Figure 3). The concentrations of MTE found in the thalli at Ain Taghrout and Tixter are high and similar, especially Cd and Zn, exceeding standard values. On the other hand, at the Ghilassa station, the thalli accumulate low concentrations of MTE, while the Pb concentration is below the standard values (40.6 mg / kg).




Fig-3: Spatial projection of stations based on the first three axes from the PCA

The use of UPGMA, a statistical method based on the Unweighted pair average and the distance from city-block (Manhattan), reflects heterogeneity in metal

accumulation in *X. parietina* thalli and confirms disaggregation of the studied sites into several clades (figure 4).




Fig-4: UPGMA based on the concentration of heavy metals

The first group is represented by the station of Ghilassa with a low MTE accumulation in the thalli, while the second group splits into two branches; The Sidi Embarek, Djaafra, Ouled Sidi Brahim, Tassameurt, El Anasseur, El Main and Ben Daoud stations with a very high Fe accumulation, exceeding standard values (804 mg / kg). The remaining sites form an amalgam representing the same groups as those coming from the PCA. Sites with high concentrations of Pb and Zn in the thalli are located close to high traffic road axes such as Ain Taghrout and Tixter and then decrease in low traffic stations.

## DISCUSSION

Assessment of levels of contamination in the BBA province for ten different metals showed Pb contamination values similar to those previously found in lichen thalli *X. parietina* in Italy [16], France [8, 24], Turkey [25], and Poland [10]. Generally, high concentrations of Pb measured in thalli near roads are closely related to vehicle emissions and fuel combustion [18, 25- 32]. In addition, in the studied areas the observed high concentrations of Pb may be related to the increase in industrial activities [8-9, 32], and the incineration of waste [24].

A low Pb accumulation in the southern mountainous sites (Kseur and Ghilassa) was observed with values between (6 and 23 mg / kg). However the concentration of Pb in northern mountainous sites was

high due to a higher air humidity level as suggested by some authors [14, 33].

The values of Zn in the thalli of *X. parietina* exceeded standard values. These high rates are due to traffic [18, 29] and industrial activities [25, 34]. The same Zn concentrations were observed in France [8-9, 24] and Turkey [25]; In contrast, low concentrations were found in Italy [16], Turkey [29] and Poland [10].

Cd concentrations are high in thalli and similar to the results observed in France [8, 9, 24], Turkey [25] and in Algeria [35]. Road traffic and anthropogenic activities [25, 36] contribute to the increase in Cd concentrations in the thalli of *X. parietina* [18, 29, 37]. The concentrations below standard values were observed in Italy [16] and Morocco [36].

The levels of Sb concentrations measured in the thalli of *X. parietina* are above standard values, whereas earlier studies in France [9, 31] and Spain [38] showed that Sb concentrations in the thalli are below standard values (0.35 mg / kg). According to the authors of these studies, high Sb levels were attributed mainly to road traffic, waste incineration and the production of batteries [9, 39]. Sb contamination spreads to rural areas where the maximum concentration is recorded in Ouled-Dahmane (253 mg / kg), and this high rate comes probably from sources outside the studied area, as reported reported by [31].

The levels of iron (Fe) in lichenic thalli are very high. The results obtained are similar to those recorded in other countries such as France [24], Turkey [25], and Poland [10]. On the other hand, the concentrations recorded in Portugal by [40] are lower than the standard values. The high rate of Fe in the studied sites has been linked to geographical factors [10, 13, 25, 37]. The other explanation for the high level of Fe in the thalli is the accumulation of dust from anthropological activities [15, 25, 32, 37, 39, 40- 42].

The results of the Cu analysis show that the levels accumulated in the thalli are high and exceed standard values. The same remarks were observed in *X. parietina*, in France [9, 24] and in Poland [10]. The highest Cu contamination was observed especially in rural communities, while the lowest contamination levels were observed in high industrial stations. The high level of Cu in rural communes is probably due to the usage of pesticides and fertilizers that can be sources of copper emission [2, 9, 16, 25]. The presence of Cu in the thalli sampled in urban sites is due to road and rail traffic, as well as to metallurgical activity [9, 10].

The concentrations of Mn recorded in the thalli of *X. parietina* are almost identical to those reported in France [9, 24] and Turkey [25]. Approximately 50% of Mn in the air is of natural origin [10, 24, 37] and to this natural source can be added an anthropic source such as the usage of fertilizers due to agricultural activities [39].

Ni levels are high and there is a gradient of concentration of this element from west to east. The same observations are cited in literature in Italy [16], Turkey [25], France [9, 24] and Poland [10]. Nickel

pollution is linked to emissions from the metallurgical industry and combustion processes [10, 18].

Chromium concentrations in the studied areas are above standard values (4.12 mg/kg) and the same results were observed in other regions such as France [9] and Italy [18, 43]. The highest accumulation of Cr was observed in western sites; which then drops in eastern sites with a minimal concentration recorded at Ain Taghrout (127.89 mg / kg). Vehicle emissions and industrial activities can be important sources of air pollution by Ni and Cr [38].

High concentrations of Co in the thalli of *X. parietina* were observed at all studied locations. Dust from quarrying and building materials can be the main reason for these high concentrations [31] however, the metal industry may be an important source of cobalt [25] as lichenic thalli may accumulate more atmospheric elements when exposed for longer periods of time [29, 33].

The classification of the elements by cluster analysis revealed two main clusters of elements with similar affinities or characteristics (figure 5). Cluster 1, included Pb, Cd, Zn, and Sb, three out of the four elements of this group (Pb, Cd and Sb) showed constant or decreased alteration loads in the present study, while Zn showed a high percentage contamination in the studied areas.

A relationship between Cd and Zn is rather common in lichens [16, 44-45]. It may be due to anthropogenic sources or metabolic interactions, since these two elements are essential for lichen metabolism [46]. The correlation between Pb and Zn indicates some affinity with atmospheric particles [31, 47].




Fig-5: Cluster dendrogram of the elements in *X. parietina*

A common feature between Cd and Zn, including the fact that they accumulate in lichen cells [48], and plant emissions are the results of indirect sources of Cd and Zn [25, 29]. According to [18], the found concentrations of Zn, Pb and Cd in sites which are far from sources of pollution are related to the usage of fertilizers and pesticides that are characteristic of the region.

Zn is generally related to the association of Pb and Cd elements. This phenomenon is ubiquitous in nature and its distribution in the atmosphere does not locate the source of the contamination [18]. More than 2% of Zn derives from vehicular traffic and of road dust; it is a component found in leaded and lead-free gasoline [10, 37, 49], confirming the significant accumulation of Zinc in thalli of lichen collected near the roads of the BBA region. 95% of Pb in plants is of atmospheric origin [10] and combustion of fuels remains the main sources of this toxic element in the environment even in mountainous areas [30].

The Sb is an element highly related to Pb in several industrial uses, because it increases the hardness of this element [9, 18]. The high coupling of Cd, Zn and Pb and their correlation with Sb also indicate their common sources in traffic activities along road axes [25]. These elements appear in vehicle components and lubricants [39], confirming that the contamination of the BBA region by these elements is of anthropogenic source.

The cluster 2 included Mn, Fe, Cu, Co, Ni and Cr, these relationships are in line with the results of other similar studies [16, 50]. All these elements substantially accumulate by particulate entrapment [32, 33, 51]. Further, the relationship of the two terrigenous macro-elements (Mn and Fe) with Ni and Cr confirmed the presence of anthropogenic influence on their distribution in lichen material. This suggested that trapped soil particulate in lichen thalli was a highly variable contribution which did not closely reflect the element concentrations in the atmosphere, but depended rather on environmental sources [15, 51].

## CONCLUSION

The results of the present survey showed that the mean concentration of all ten elements in *X. parietina* thalli, with the exception of Cd, are included between middle to high naturalness classes. Multivariate analyses allowed us to find the main correlations among the groups of element taken into account in the present study. The changes in the concentrations of the MTE, contained in the thalli, can provide useful information on the long-term assessment of air pollution and to monitor the progress of the contamination phenomenon in the time. It is important to focus on the possibility of monitoring a large study area completely lacking an instrumental network for detecting this kind of anthropogenic impact. This confirms that lichen bio-

monitoring can be useful in risk assessment for human health and it can be a powerful tool for administrators involved in environmental planning.

## ACKNOWLEDGEMENTS

The work was supported by Algerian MESRS and Chemical Laboratory of carbohydrates Heterocyclic of Clermont Ferrant, France

## REFERENCES

1. Pinho P, Bergamini A, Carvalho P, Branquinho C, Stofer S, Scheidegger C, Mâgias C. Lichen functional groups as ecological indicators of the effects of land-use in Mediterranean ecosystems. *Ecological indicators*. 2012; 15: 36-42.
2. Parzych A, Zdunczyk A, Astel A. Epiphytic lichens as bioindicators of air pollution by heavy metals in an urban area (northern Poland). *Journal of Elementology*. 2016;21(3).
3. Garrec JP, Van Haluwyn Ch. *Biosurveillance végétale de la qualité de l'air, concepts, méthodes et applications*. Editions Tec et doc, Lavoisier; Paris. 2002; 118p.
4. Rzepka MA, Cuny D. *Biosurveillance Végétale et Fongique des ETM Atmosphériques*. *Air Pur*. 2008; 75: 66-79.
5. Van Haluwyn Ch. *Quelle place et Quelles perspectives pour la biosurveillance végétale dans les études de la qualité de l'air en France*. *Air Pur*. 2009; 77: 5-10.
6. Van Haluwyn Ch, Cuny D, Garrec JP. *Définition, Concepts et Enjeux Généraux de la Biosurveillance de la qualité de l'air*. *Pollution Atmosphérique*. 2011; Numéro Spécial: 7-12.
7. Cuny D, Quarre S, Devred I, Jeanne B, Reynaert A, Rzepka MA, Cuny MA. *Evolution Spatiale et temporelle des concentrations en éléments traces métalliques de thalles de Xanthoria parietina récoltes sur la zone Industriale-Portuaire de Dunkerque*. *Air Pur*. 2009 ; 77: 43-49.
8. Cuny D, Rzepka MA, Van Haluwyn C. *Introduction à la biosurveillance végétale et Fongique de la qualité de l'air-Concept et intérêt*. *Spectra Analyse*. 2008 ; 264: 31-35.
9. Occelli F, Cuny MA, Devred I, Deram A, Quarre S, Cuny D. *Étude de l'imprégnation de l'environnement de trois bassins de vie de la région Nord-Pas-de-Calais par les éléments traces métalliques*. *Pollution Atmosphérique*. 2013;220:1-10.
10. Parzych A, Astel A, Zdunczyk A, Surowiec T. *Evaluation of urban environment pollution based on the accumulation of macro- and trace elements in epiphytic lichens*. *Journal of Environmental Science and Health*. 2016; 51(4): 297-308.
11. Paoli L, Corsini A, Bigagli V, Vannini J, Bruscoli C, Loppi S. *Long-term biological monitoring of environmental quality around a solid waste landfill assessed with lichens*. *Environmental Pollution*. 2012; 161: 70-75.

12. Ezeonyejiaku CD, Obiakor MO. Spatial Biomonitoring of Trace Element Contamination and Atmospheric Quality Assessment. *Am. J. Life. Sci. Res.* 2013; 1(2): 67-73.
13. Nimis PL, Andreussi S, Pittao E. The performance of tow lichen species as bioaccumulators of trace metals. *Sci. Total Environ.* 2001; 275: 43-51.
14. Kar S, Samal AC, Maity JP, Santra SC. Diversity of epiphytic lichens and their role in sequestration of atmospheric metals. *Int. J. Environ. Sci. Technol.* 2014; 11: 899-908.
15. Branquinho C, Catarino F. Improving the use of lichens as biomonitors of atmospheric metal pollution. D.H. BROWN. School of Biological Sciences, University of Bristol, United Kingdom, Portugal. 1997; pp. 5-16.
16. Nimis PL, Lazzarin G, Lazzarin A, Skert N. Biomonitoring of trace elements with lichens in Veneto Z NE Italy. *Sci. Total Environment.* 2000; 255: 97-111.
17. Nimis PL, Scheidegger Ch, Wolsely PA. Monitoring with lichens—Monitoring lichens. *Nato Science Series, IV.* Kluwer Academic Publishers, Netherlands, 2002; 408p.
18. Scerbo R, Ristori T, Possenti L, Lampuganini L, Barale R, Barghigiani C. Lichen (*Xanthoria parietina*) biomonitoring of trace element contamination and air quality assessment in pisa province (Tuscany, Italy). *Sci. Total Environ.* 2002; 286: 27-40.
19. Messaoud R, Takia L. The use of lichens bioindicators in the identification of air pollution. 2015.
20. Rusu AM. Sample preparation of lichens for elemental analysis. In *Monitoring with lichens—monitoring lichens 2002* (pp. 305-309). Springer, Dordrecht.
21. Hébrard-Labit C, Meffray L. Comparaison de méthodes d'analyse des éléments traces métalliques (ETM) et des hydrocarbures aromatiques polycycliques (HAP) sur les sols et les végétaux.
22. Bettinelli M, Spezia S, Bizzarri G. Trace element determination in lichens by ICP-MS. *Atomic Spectroscopy.* 1996; 17(3):133-41.
23. Quevauviller P, Herzig R, Muntau H. Certified reference material of lichen (CRM 482) for the quality control of trace element biomonitoring. *Science of the total environment.* 1996 Aug 30; 187(2):143-52.
24. Dron J, Austruy A, Agnan Y, Giraudeau P, Chamaret P. Utilisation de la biosurveillance lichénique sur la zone industrialo-portuaire de Fos-sur-Mer: retour sur trois ans de suivi à l'échelle d'un territoire intercommunal. *Pollution Atmosphérique.* 2016; 228: 1-14.
25. Demiray AD, Yolcubal I, Akyol NH, Çobanğlu G. Biomonitoring of airborne metals using the Lichen *Xanthoria parietina* in Kocaeli Province, Turkey. *Ecological Indicators.* 2012; 18: 632-643.
26. Rahali M. Cartographie de la pollution plombique dans la région d'Alger en utilisant un lichen (*Xanthoria parietina*) comme bioaccumulateur. *Pollution Atmosphérique.* 2002 ; 175: 421-432.
27. Loppi S, Frati L, Paoli L, Bigagli V, Rossetti C, Bruscoli C, Corsini A. Biodiversity of epiphytic lichens and heavy metal contents of *Flavoparmelia caperata* thalli as indicators of temporal variations of air pollution in the town of Montecatini Terme (central Italy). *Science of the Total Environment.* 2004; 326: 113-122.
28. Maher BA, Moore C, Matzka J. Spatial variation in vehicle-derived metal pollution identified by magnetic and elemental analysis of roadside tree leaves. *Atmospheric Environment.* 2008; 42: 364-373.
29. Kinalioglu K, Ozbucak TB, Kutbay HG, Huseyinova R, Bilgin A, Demirayak A. Biomonitoring of Trace Elements with Lichens in Samsun City, Turkey, *Ekoloji.* 2010; 75(19): 64-70.
30. Maatoug M, Medkour K, Ait Hammou M, Ayad N. Cartographie de la pollution atmosphérique par le plomb d'origine routière à l'aide de la transplantation d'un lichen bioaccumulateur *Xanthoria parietina* dans la ville de Tiaret (Algérie). *Pollution Atmosphérique.* 2010; 205: 93-101.
31. Agnan A, Séjalon-Delmas N, Probst A. Comparing early twentieth century and present-day atmospheric pollution in SW France: A story of lichens. *Environmental Pollution.* 2013; 172: 139-148.
32. Francová A, Chrástný V, Sillerová H, Vitková M, Kocourková J, Komárek M. Evaluating the suitability of different environmental samples for tracing atmospheric pollution in industrial areas. *Environmental Pollution.* 2017; 220(part A): 286-297.
33. Klimek B, Tarasek A, Hajduk J. Trace Element Concentrations in Lichens Collected in the Beskid Mountains, the Outer Western Carpathians. *Bull Environ Contam Toxicol.* 2015; 94: 532- 536.
34. Lin YP, Chang TK, Chung-Wen Shih CW, Tseng CH. Factorial and indicator kriging methods using a geographic information system to delineate spatial variation and pollution sources of soil heavy metals. *Environmental Geology.* 2002; 42: 900-909.
35. Sarmoum M, Djebbar R, Latreche K. Bioaccumulation de trois métaux lourds (Pb, Zn et Cd) par le lichen, *Xanthoria parietina*, dans la région Algéroise. *Revue Ecologie Environnement.* 2014 ; 10: 25-29.
36. Ouali-Alami FZ, Elabidi A, Mouhir L, Fekhaoui M, Serghini A. Utilisation des lichens comme bio-indicateurs de la pollution atmosphérique par le plomb, cadmium et zinc de la région de Rabat-Sale-Zemmour-Zaër (Maroc). *Afrique Science.* 2014; 10 (3): 89-106.

37. Aslan A, Cicek A, Yazici K, Karagoz Y, Turan M, Akkus F, Yildirim OS. The assessment of lichens as bioindicator of heavy metal pollution from motor vehicles activities. *African Journal of Agricultural Research*. 2013; 6(7): 1698-1706.
38. Achotegui-Castells A, Sardans J, Ribas A, Peñuelas J. Identifying the origin of atmospheric inputs of trace elements in the Prades Mountains (Catalonia) with bryophytes, lichens, and soil monitoring. *Environmental Monitoring and Assessment*. 2013; 185: 615-629.
39. Azimi S, Rocher V, Muller M, Moilleron R, Thevenot DR. Sources, distribution and variability of hydrocarbons and metals in atmospheric deposition in an urban area (Paris, France). *Science of the Total Environment*. 2005; 337: 223-239.
40. Branquinho C, Gaio-Oliveira G, Augusto S, Pinho P, Maguas C, Correia O. Biomonitoring spacial and temporal impact of atmospheric dust from a cement industry. *Environmental Pollution*. 2008; 151: 292-299.
41. Nieboer E, Richardson HS, Tomassini FD. Mineral uptake and release by lichens: an overview. *Bryologist*. 1978; 81(2): 226-246.
42. Bargagli R. The elemental composition of vegetation and the possible incidence of soil contamination of samples. *Science of the Total Environment*. 1995; 176 (1-3): 121-128.
43. Scerbo R, Possenti L, Lampugnani L, Ristori T, Barale R, Barghigiani C. Lichen (*Xanthoria parietina*) biomonitoring of trace element contamination and air quality assessment in Livorno Province (Tuscany, Italy). *The Science of the Total Environment*. 1999; 241: 91-106.
44. Loppi S, Pirintsos SA. Epiphytic lichens as sentinels of heavy metal pollution at forest ecosystems (central Italy). *Environ Pollut*. 2003; 121(3): 327-332.
45. Brunialti G, Frati L. Biomonitoring of nine elements by the lichen *Xanthoria parietina* in Adriatic Italy: A retrospective study over a 7-year time span. *Science of the Total Environment*. 2007; 387: 289-300.
46. Ormrod DP. Impact of trace element pollution on plants. *Air pollution and plant life*/edited by Michael Treshow. 1984.
47. Gandois L and Probst A. Localisation and mobility of trace metal in silver fir needles. *Chemosphere*. 2012; 87(2): 204-210.
48. Loppi S. Licheni come bioaccumulatori di elementi in traccia: stato della ricerca in Italia. *Biologia Ambientale*; 2006; 20(2): 69-78.
49. Jeran Z, Mrak T, Jaćimović R, Batić F, Kastelec D, Mavsar R, Simončić P. Epiphytic lichens as biomonitors of atmospheric pollution in Slovenian forests. *Environmental Pollution*. 2007; 146: 324-331.
50. Bennett JP, Wetmore CM. Changes in element contents of selected lichens over 11 years in northern Minnesota, USA. *Environ Exp Bot*. 1999; 41: 75-82.
51. Loppi S, Pirintsos SA, De Dominicis V. Soil contribution to the elemental composition of epiphytic lichens (Tuscany, central Italy). *Environ Monit Assess*. 1999; 58: 121-131.