

The Study for Determined Isolates Lactic Acid Bacteria as Commercial Starter Culture Isolated from Dadih

Anif Mukaromah Wati^{1,2*}, Mei-Jen Lin¹, Lilik Eka Radiati²

¹Department of Animal Science, National Pingtung University of Science and Technology, Pingtung County, Taiwan

²Faculty of Animal Science, Brawijaya University, Malang, East Java, Indonesia

Original Research Article

*Corresponding author

Anif Mukaromah Wati

Article History

Received: 08.02.2018

Accepted: 21.02.2018

Published: 30.03.2018

DOI:

10.36347/sajb.2018.v06i03.001



Abstract: Dadih is a traditional food from West Sumatra, Indonesia, which is made by buffalo milk that naturally fermented in bamboo tubes. Natural starter of dadih needs to be changed with artificial starter (starter culture) then dadih can be produced commercially. This study aimed to identify the capability of isolates lactic acid bacteria (LAB) from dadih to ferment milk that determined with pH value, Titratable Acidity (TA), viability of LAB, and physical characteristic during fermentation process. Different fermentation time, skim milk that added with isolates was a significant difference ($p < 0.05$) on pH value, TA, viability of LAB, and physical characteristic. During fermentation process, pH was decreased, TA and viability of LAB was increased, and made a curd formation and syneresis at the end of fermentation. The study concluded that isolates 11, 21, 25, 29, and 41 that isolated from dadih have ability as starter culture based on pH value, TA, viability of LAB, and curd formation. Then isolates can use commercially.

Keywords: pH, titratable acidity (TA), viability of LAB, curd formation, fermentation time.

INTRODUCTION

Dadih is a traditional food from West Sumatra, Indonesia, which is made by buffalo milk that naturally fermented in bamboo tubes. Indonesian dadih is still processed traditionally with natural starter from bamboo. Natural starter of dadih needs to be changed with artificial starter (starter culture) then dadih can be produced commercially.

Fermented milk product is produced by dairy industry with controlled conditions to make a good product [1]. To make starter cultures, LAB from dadih must be isolated and identified.

Dadih contains *Lactobacillus casei* subsp. *casei*, *Leuconoctoc paramasenteroides*, *Enterococcus faecalis* subsp. *liquefaciens*, and *Lactococcus lactis* subsp. *lactis* [2]. During fermentation process, isolates can identify the capability as starter culture to ferment milk with determined pH value and viability of LAB [3]. This study aimed to identify the capability of isolates LAB from dadih to ferment milk that determined with pH value, Titratable Acidity (TA), viability of LAB, and physical characteristic during fermentation process.

MATERIALS AND METHODS

Preparation of the isolates

Isolates was obtained from Dairy Laboratory, Department of Animal Science, National Pingtung University of Science and Technology. Isolates were isolated from Dadih. Isolates have characteristic rod shape, Gram stain positive, and negative motility test

and catalase test. LAB have characteristic Gram stain positive and negative motility test [4]. Isolates that used in this experiment were isolate 11, isolate 21, isolate 25, isolate 29, and isolate 41. Before using, isolates were subculture in the broth (Merck, Darmstadt, Germany) and incubated at 37°C for 20 h.

Preparation of the sample

Preparation of sample was determined from Wardani *et al.* [5] with some modification. To preparation skim milk, skim milk powder (Fonterra, Amsterdam, Netherlands) 10% (v/v) was autoclaved, added with 3% of each isolates, incubated at 37°C, and measured of pH, TA, viability of LAB, and physical characteristic of each 6 h until 30 h.

Test for TA

TA was determined with 9 g samples that diluted with 9 ml sterile water, then sample was titrated

with 0.1 N NaOH using 2 drops phenolphthalein solution (Nalacai Tesque, Inc, Kyoto, Japan) [6].

Test for pH value

pH meter (Suntex pH/ION meter SP-2500, New Taipei City, Taiwan) was used for determining pH. Before determination, pH meter was calibrated by using buffer solutions no. 4 and 7 (Mettler Toledo, Greifensee, Langacher) [7].

Test for viability of LAB

Five milliliters of samples were added to 45 ml of sterile 0.1% peptone (Merck, Darmstadt, Germany) and homogenized in a stomacher lab blender (400) (AES Laboratory, Combourg, France) for 1 min, solution was obtained to make 10^{-1} dilution. One milliliter from dilution (10^{-1}) was aseptically transferred to 9 ml sterile distilled water [7]. One milliliter of sample from tubes 10^6 - 10^8 was pipetted with micropipet to petri dishes (Alpha Plus Scientific Corp, Taoyuan Hsien, Taiwan), then 15 ml of sterilized MRS (Merck, Darmstadt, Germany) supplemented with 1.5% CaCO_3 medium (Nihon Shiyaku reagent, Taipei, Taiwan) was poured into petri dishes. The plates were incubated at 37°C for 48 h. The existence of LAB was clear zone surrounding colonies then colonies were counted by SUNTEX colony counter 570 (Suntex Instruments Co., LTD., New Taipei City, Taiwan).

STATISTICAL ANALYSIS

Statistical Analysis System (version 9.3, SAS) was used to perform data analysis and if there was a significantly different then continued by Duncan's Multiple Range Test (DMRT).

RESULTS

Capability of isolates to ferment milk was determined with measuring pH value, TA, viability of LAB, and physical characteristic during fermentation process.

pH value and TA

pH value of skim milk that adding each isolate was measured during fermentation process. There was a significant difference ($p < 0.05$) during fermentation process on pH value and TA. Isolate 21 was the strongest to decrease pH value from 5.25 to 4.61. In the other hand, isolate 41 was the weakest ranged from 5.63 to 4.85. Additionally, isolate 11 from 5.18 to 4.68, isolate 25 was decreased from 5.63 to 4.65, and isolate 29 from 5.38 to 4.89. pH and TA had negative correlation. When pH was decreased, TA was increased during fermentation process. In the end of fermentation process, isolate 21 was the lowest pH while TA value was the highest 0.68.

Viability of LAB

There was a significant difference ($p < 0.05$) during fermentation process. Viability of LAB was varied during fermentation. Increasing of fermentation time was effected to increasing viability of LAB. At the end of fermentation, isolate 21 was the highest viability of LAB 8.98 (cfu/ml) while it was the highest of TA and the lowest of pH value. Moreover, isolate 41 was the lowest 4.93 (cfu/ml) while it was the lowest of TA and the highest of pH.

Physical characteristic

There was a significant difference ($p < 0.05$) on physical characteristic in skim milk during fermentation. physical characteristic was changed by isolates. All of the skim milk was still liquid in 6 h. After 12 h, skim milk that added isolates (11, 21, and 29) began to make a curd that showed in Figure-1. Curd formation was correlation with decreasing of pH and increasing viability of LAB and TA. On the other hand, skim milk in isolates 25 and 41 were still liquid because pH was still high while viability of LAB and TA were still low.

At the 18 h fermentation time, curd formation was appeared when pH value was ranged from 4.78 (isolate 21) to 4.93 (isolate 41). At the 24 h fermentation time, skim milk in isolates 11, 21, and 25 had stronger to make a curd, but isolates 29 and 41 still had same curd with at the 18 h because isolate 29 and 41 had similar pH value, TA, and viability of LAB between 18 and 24 h. After 30 h fermentation time, isolates 11, 21 and 25 began to make a syneresis because they had low pH and high TA and viability of LAB. However, isolates 29 and 41 still had medium curd formation because they had high pH and low TA and viability of LAB.

DISCUSSION

During fermentation process, isolates had ability to change pH value, TA, viability of LAB, and physical characteristic. Increasing of TA and decreasing of pH were indicated that isolates had ability as starter culture. pH value was decreased that caused by lactic acid production. pH value during incubation was reduced by proteolytic activity of isolates [3]. RNA and protein synthesis were affected by pH value [8]. Viability of LAB was increasing during incubation while increasing of TA because lactic acid was produced by LAB from lactose in dairy products [9]. Curd formation in the end of fermentation because lactic acid was converted by LAB from lactose, then precipitated a milk protein, formed curd, and other metabolites [10].

Table-1: Quality of skim milk with adding isolates during fermentation

| Isolates | Fermentation time hour (h) ¹ | pH value ¹ | TA (%) ¹ | Viability of LAB (cfu/ml) ¹ | Physical characteristic ³ |
|----------|-----------------------------------------|---------------------------|--------------------------|----------------------------------------|--------------------------------------|
| 11 | 6 | 5.18 ± 0.01 ^{ab} | 0.32 ± 0.00 ^c | 8.81 ± 0.01 ^d | - |
| | 12 | 4.97 ± 0.01 ^b | 0.41 ± 0.00 ^d | 8.87 ± 0.01 ^c | + |
| | 18 | 4.83 ± 0.00 ^c | 0.46 ± 0.00 ^c | 8.89 ± 0.00 ^b | ++ |
| | 24 | 4.78 ± 0.00 ^d | 0.50 ± 0.00 ^b | 8.92 ± 0.00 ^a | ++++ |
| | 30 | 4.68 ± 0.00 ^e | 0.63 ± 0.00 ^a | 8.93 ± 0.00 ^a | * |
| 21 | 6 | 5.25 ± 0.01 ^a | 0.27 ± 0.01 ^e | 8.72 ± 0.02 ^d | - |
| | 12 | 4.96 ± 0.00 ^b | 0.42 ± 0.00 ^d | 8.89 ± 0.00 ^c | + |
| | 18 | 4.78 ± 0.01 ^c | 0.51 ± 0.00 ^c | 8.92 ± 0.00 ^b | +++ |
| | 24 | 4.71 ± 0.01 ^d | 0.61 ± 0.00 ^b | 8.94 ± 0.00 ^b | ++++ |
| | 30 | 4.61 ± 0.01 ^e | 0.68 ± 0.00 ^a | 8.98 ± 0.00 ^a | * |
| 25 | 6 | 5.63 ± 0.01 ^a | 0.14 ± 0.00 ^c | 8.49 ± 0.02 ^d | - |
| | 12 | 5.45 ± 0.01 ^b | 0.19 ± 0.00 ^d | 8.56 ± 0.00 ^c | - |
| | 18 | 4.83 ± 0.03 ^c | 0.47 ± 0.02 ^c | 8.88 ± 0.00 ^b | ++ |
| | 24 | 4.71 ± 0.01 ^d | 0.61 ± 0.00 ^b | 8.94 ± 0.00 ^a | ++++ |
| | 30 | 4.65 ± 0.00 ^e | 0.65 ± 0.00 ^a | 8.96 ± 0.00 ^a | * |
| 29 | 6 | 5.38 ± 0.02 ^a | 0.22 ± 0.01 ^d | 8.58 ± 0.01 ^b | - |
| | 12 | 5.04 ± 0.01 ^b | 0.40 ± 0.00 ^c | 8.71 ± 0.01 ^a | + |
| | 18 | 4.89 ± 0.01 ^c | 0.43 ± 0.00 ^b | 8.72 ± 0.01 ^a | ++ |
| | 24 | 4.90 ± 0.01 ^c | 0.43 ± 0.00 ^b | 8.72 ± 0.01 ^a | ++ |
| | 30 | 4.89 ± 0.00 ^c | 0.44 ± 0.00 ^a | 8.73 ± 0.01 ^a | ++ |
| 41 | 6 | 5.63 ± 0.01 ^a | 0.15 ± 0.01 ^d | 8.49 ± 0.02 ^d | - |
| | 12 | 5.33 ± 0.01 ^b | 0.24 ± 0.00 ^c | 8.59 ± 0.00 ^c | - |
| | 18 | 4.93 ± 0.00 ^c | 0.43 ± 0.00 ^b | 8.67 ± 0.02 ^b | ++ |
| | 24 | 4.92 ± 0.00 ^c | 0.43 ± 0.00 ^b | 8.72 ± 0.02 ^{ab} | ++ |
| | 30 | 4.85 ± 0.01 ^d | 0.45 ± 0.00 ^a | 8.74 ± 0.00 ^a | ++ |

¹Data are expressed as means values ± standard deviation (n = 3).

²Mean values in same column with difference letter are statistically significant (p<0.05).

³Physical characteristic: (-): liquid, (+): little curd, (++): medium curd, (+++): strong curd, (++++): very strong curd, (*): curd and made a syneresis



Fig-1: Curd formation of skim milk

CONCLUSION

This study shows that isolates 11, 21, 25, 29, and 41 that isolated from dadih have ability as starter culture based on pH value, TA, viability of LAB, and curd formation. Then isolates can use commercially.

ACKNOWLEDGEMENT

I sincerely thank to my research advisor Dr. Mei-Jen Lin for her advice and technical support in conducting experiments. Moreover, i would thank to National Pingtung University of Science and

Technology (NPUST), Taiwan because NPUST give me opportunity to study and funding my experiments.

REFERENCES

1. Widyastuti Y, Rohmatussolihat, Febrisiantosa A. The role of lactic acid bacteria in milk fermentation. Food and Nutrition Sciences. 2014; 5:435-442.
2. Pato U. Bile and acid tolerance of lactic acid bacteria isolated from dadih and their antimutagenicity against mutagenic heated tauco.

- Asian-Australas. J. Anim. Sci. 2003; 16(11):1680-1685.
3. Yelnetty A, Purnomo H, Purwadi, Mirah A. Biochemical characteristics of lactic acid bacteria with proteolytic activity and capability as starter culture isolated from spontaneous fermented local goat milk. J. Nat. Sci. Res. 2014; 4(10):137-146.
 4. Syukur S, Hermansyah A, Fachrial E. Probiotics and strong antimicrobial of buffalo milk fermentation (dadih) from different places in West Sumatera Indonesia. Res. J. Pharm. Biol. Chem. Sci. 2016; 7(6):386-392.
 5. Wardani, SK, Cahyanto MN, Rahayu ES, Utami T. The effect of inoculum size and incubation temperature on cell growth, acid production and curd formation during milk fermentation by *Lactobacillus plantarum* Dad 13. Int. Food Res. J. 2017; 24(3):921-926.
 6. Wang YC, Yu RC, Yang HY, Chou CC. Sugar and acid contents in soymilk fermented with lactic acid bacteria alone or simultaneously with bifidobacteria. Food Microbiology. 2003; 20(3):333-338.
 7. Abdalla MOM, Elsayed ZEA, Nour AAM. Effect of source of heat and storage period on physicochemical and microbiological characteristics of goat milk. Int. J. Dairy Sci. Process. 2015; 2(4): 18-23.
 8. Panesar PS, Kennedy JF, Knill CJ, Kosseva M. Production of L(+) Lactic Acid using *Lactobacillus casei* from Whey. Braz. Arch. Biol. Technol. 2010;53(1): 219-226.
 9. Che Othman SF, Noor Aziah AA, Ahmad R. Effects of jackfruit puree supplementation on lactic acid bacteria (*Lactobacillus acidophilus* FTDC 1295) in terms of viability and chemical compositions of dadih. Int. Food Res. J. 2012; 19(4):1687-1697.
 10. Surono IS. Traditional Indonesian dairy foods. Asia Pac. J. Clin. Nutr. 2015; 24(1):26-30.