Scholars Academic Journal of Biosciences
Abbreviated Key Title: Sch Acad J Biosci
ISSN 2347-9515 (Print) | ISSN 2321-6883 (Online)
Journal homepage: https://saspublishers.com
NLM ID:101629416

OPEN ACCESS

Oligochaetology

Earthworm Taxonomy in the Twenty-First Century: Why Species Redescription Still Matters

John Warren Reynolds[1*]


[1]Oligochaetology Laboratory, 9-1250 Weber Street East, Kitchener, ON Canada N2A 4E1, and Research Associate, New Brunswick Museum, Saint John, NB Canada E2K 1E5

DOI: https://doi.org/10.36347/sajb.2026.v14i11.001 | **Received:** 13.09.2026 | **Accepted:** 04.11.2026 | **Published:** 07.11.2026

***Corresponding author:** John Warren Reynolds
Oligochaetology Laboratory, 9-1250 Weber Street East, Kitchener, ON Canada N2A 4E1, and Research Associate, New Brunswick Museum, Saint John, NB Canada E2K 1E5

Abstract **Review Article**

Earthworm taxonomy remains fundamental to biodiversity science, ecology, biogeography, and conservation. Although molecular techniques have transformed systematic biology, many earthworm species continue to be known only from historical descriptions published during the late nineteenth and early twentieth centuries. These early accounts often lack sufficient anatomical detail, illustrations, information on intraspecific variation, or accessible type material, limiting their usefulness for modern taxonomic investigations. Species redescriptions play a critical role in addressing these deficiencies by refining diagnostic characters, documenting variation, clarifying species boundaries, and improving nomenclatural stability. Redescriptive studies also provide an essential foundation for molecular analyses, ensuring that DNA-based investigations are anchored to accurately identified specimens. The integration of traditional morphological approaches with modern molecular methodologies offers the most robust framework for understanding earthworm diversity and evolutionary relationships. As global biodiversity research increasingly depends upon reliable taxonomic data, species redescription remains a vital component of contemporary earthworm systematics and an indispensable tool for documenting and conserving soil biodiversity.



Graphical Abstract: Conceptual framework illustrating the continuing importance of species redescription in twenty-first-century earthworm taxonomy. Historical descriptions, often constrained by limited material and analytical techniques, are re-evaluated using contemporary morphological and molecular approaches. These redescriptions improve species diagnoses, enhance nomenclatural stability, and provide reliable taxonomic foundations for biodiversity assessment, ecological investigations, conservation initiatives, and evolutionary research.

Keywords: *Staphylococcus aureus*, Biofilm, Antibiotics Resistance, Multidrug Resistance, Burn Wound Infection.

Copyright © 2026 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution **4.0 International License (CC BY-NC 4.0)** which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

INTRODUCTION

Taxonomy forms the foundation of biological science. Accurate species identification underpins ecological studies, conservation planning, biogeographical investigations, and biodiversity assessments. Yet, despite remarkable advances in molecular biology and bioinformatics, many earthworm species remain inadequately described, poorly illustrated, or known only from historical accounts published more than a century ago (Blakemore, 2008; Csuzdi, 2012).

The continuing need for taxonomic revision is particularly evident in the family Megascolecidae and many other earthworm groups across Asia, Africa, South America, and Oceania. Numerous species were originally described from limited material, often based on a single specimen, using standards and methodologies available at the time. While these pioneering works remain invaluable, modern taxonomists frequently encounter difficulties when attempting to identify specimens solely from the original descriptions.

Recent taxonomic investigations have demonstrated that species redescriptions remain an essential component of contemporary systematic research and biodiversity documentation (Dayrat, 2005; Blakemore, 2008; Reynolds and Wetzel, 2021). Far from being repetitive exercises, they provide the critical framework needed to stabilize nomenclature and improve biological understanding.

HISTORICAL CHALLENGES

Many earthworm species were described during the late nineteenth and early twentieth centuries by authors such as Beddard (1895), Michaelsen (1900), Rosa (1891), Stephenson (1930), Gates (1972), and others. Their contributions established the foundations of modern oligochaete taxonomy. However, diagnostic characters that were considered sufficient at the time may no longer adequately distinguish among closely related taxa.

Several limitations are frequently encountered in historical descriptions:

- Incomplete anatomical information.
- Limited illustrations or absence of figures.
- Small sample sizes.
- Lack of information regarding intraspecific variation.
- Uncertain locality data.
- Absence of deposited type material or inaccessible collections.

As a result, taxonomists often face uncertainty when attempting to compare newly collected specimens with historical taxa.

Numerous earthworm taxa remain difficult to interpret because original descriptions were based on a single specimen or lacked illustrations of diagnostic reproductive structures. In some instances, type specimens have deteriorated, become inaccessible, or cannot be conclusively located within museum collections. Consequently, later authors may apply the same species name to different biological entities, resulting in taxonomic confusion that can persist for decades. Modern redescriptions provide an opportunity to reassess these taxa using larger series of specimens and standardized anatomical criteria.

Additional difficulties arise when original descriptions were based upon a single specimen or lacked illustrations of critical reproductive structures. In some cases, type specimens have deteriorated through time, are unavailable for study, or cannot be readily located within museum collections. Such circumstances may result in uncertainty regarding species identity and can lead to inconsistent application of names by subsequent authors. Modern redescriptions provide an opportunity to re-evaluate historical taxa using larger specimen series, standardized anatomical terminology, and contemporary diagnostic criteria, thereby reducing ambiguity and improving taxonomic resolution (Beddard, 1895; Michaelsen, 1900; Stephenson, 1930; Gates, 1972).

THE IMPORTANCE OF REDESCRIPTION

Species redescription serves multiple scientific functions (Gates, 1972; Blakemore, 2008).

First, it refines diagnostic characters and provides comprehensive morphological documentation using modern standards. Detailed examination of external and internal anatomy allows researchers to confirm species identities and distinguish between similar taxa.

Second, redescription can reveal previously overlooked variation within species. Such information is important for evaluating synonymies and understanding geographical variation across populations.

Third, redescription contributes directly to nomenclatural stability (ICZN, 1999; Reynolds and Wetzel, 2021). Clarified diagnoses reduce misidentifications and facilitate consistent species concepts among researchers working in different regions (ICZN, 1999).

Finally, redescriptions establish a stronger foundation for future molecular investigations. DNA-based studies depend upon accurately identified specimens. Without reliable taxonomy, molecular datasets may inadvertently incorporate misidentified material, creating additional confusion rather than resolving it.

Integrating Traditional and Modern Approaches

The ongoing debate between classical morphology and molecular taxonomy is often presented as a choice between competing methodologies. In reality, these approaches are complementary.

Morphological investigations remain indispensable because species descriptions are ultimately based upon observable diagnostic characters. Molecular techniques, including DNA barcoding and phylogenetic analyses, provide additional lines of evidence that can help resolve complex taxonomic questions (James *et al.*, 2010; Perez-Losada *et al.*, 2009).

The most robust systematic studies integrate both approaches whenever possible. Comprehensive morphological documentation combined with molecular evidence provides a powerful framework for species delimitation and evolutionary interpretation (Dayrat, 2005; Blakemore, 2008; James *et al.*, 2010).

Increasingly, systematic biologists advocate an integrative taxonomic framework in which multiple independent sources of evidence are evaluated simultaneously. Morphological characters, molecular datasets, geographical distributions, ecological information, and, where available, behavioural observations may all contribute to species delimitation. Such approaches provide a more comprehensive understanding of biodiversity than any single line of evidence alone. While molecular techniques frequently reveal previously unrecognized diversity, morphology remains essential because species names are ultimately linked to physical specimens and diagnosable characters preserved within museum collections (Dayrat, 2005; James *et al*., 2010; Blakemore, 2008).

At the same time, many regions of the world still lack access to extensive molecular facilities. For these areas, detailed morphological research remains the primary means of documenting biodiversity and describing new taxa.

Increasingly, taxonomists advocate an integrative approach in which multiple independent datasets are evaluated simultaneously. Morphological characters, molecular sequences, geographical distributions, ecological data, and behavioural observations may each contribute evidence regarding species limits. While molecular data can reveal cryptic diversity, morphology remains essential because taxonomic names are ultimately attached to observable specimens and diagnosable characters. Integrative taxonomy therefore seeks not to replace traditional methods but to strengthen them through multiple corroborating lines of evidence.

Significance for Biodiversity Research

Earthworms represent one of the most ecologically important groups of soil invertebrates. They influence nutrient cycling, soil structure, organic matter decomposition, and ecosystem productivity. Yet earthworm diversity remains incompletely known in many regions (Decaëns, 2010).

Accurate taxonomy directly influences our understanding of:

- Species distributions.
- Endemism.
- Invasive species dynamics.
- Conservation priorities.
- Ecological function.

Errors in species identification can propagate through ecological literature, leading to inaccurate distribution records and misleading conclusions regarding biodiversity patterns.

Consequently, taxonomic revision and species redescription should be viewed not as isolated exercises, but as essential components of biodiversity science.

Accurate taxonomy is also increasingly important for environmental monitoring programs, biodiversity databases, and conservation assessments. Misidentified specimens incorporated into faunal inventories, ecological datasets, or genetic repositories can propagate errors through multiple studies, potentially affecting conclusions regarding species distributions, endemism, and ecosystem function. Reliable taxonomic identifications improve the quality of biodiversity information and strengthen the scientific basis upon which management strategies and conservation decisions are made (Decaëns, 2010; Csuzdi, 2012; James *et al*., 2010).

In addition, accurate taxonomy is increasingly important for environmental monitoring programs and global biodiversity databases. Misidentified records entered into regional faunal inventories, conservation assessments, or molecular repositories may propagate errors across multiple studies. Reliable species identifications improve the quality of biodiversity datasets and strengthen the scientific foundation upon which environmental policy and conservation decisions are based.

Looking Forward

The future of earthworm taxonomy depends upon the continued integration of historical scholarship, field exploration, museum collections, and emerging technologies. Digitization of collections, high-resolution imaging, and molecular methods provide unprecedented opportunities for reassessing historical taxa and expanding our knowledge of global earthworm diversity (Reynolds and Wetzel, 2021).

Emerging technologies continue to transform taxonomic research. Digitization of museum collections, online biodiversity databases, high-resolution imaging systems, and the development of DNA reference libraries now permit broader comparative analyses than were previously possible. Nevertheless, the value of these resources depends upon accurate species identifications and expert taxonomic interpretation. Equally important is the training of future taxonomists capable of combining knowledge of classical literature with modern analytical methods. Without continued investment in taxonomic expertise, technological advances alone will be insufficient to address the substantial gaps that remain in our understanding of global earthworm diversity (Dayrat, 2005; Blakemore, 2008; Reynolds & Wetzel, 2021).

At the same time, there remains an urgent need for specialists capable of interpreting classical literature and applying rigorous taxonomic principles. Many species described decades ago await modern reassessment, and numerous regions remain poorly surveyed.

The rediscovery and redescription of historically neglected taxa frequently yield important biological insights while strengthening the taxonomic framework upon which ecological and evolutionary studies depend.

Despite significant technological advances, one of the greatest challenges facing earthworm systematics is the declining number of trained taxonomists worldwide. The development of digital collections, online databases, and global DNA reference libraries will facilitate future research, but these resources remain dependent upon expert interpretation. Training the next generation of taxonomists may prove as important as the development of new technologies themselves.

CONCLUSION

Species redescription remains a vital and relevant aspect of contemporary earthworm systematics. By clarifying diagnoses, documenting variation, and improving taxonomic stability, redescriptive studies provide the foundation necessary for biodiversity research, ecological investigations, and future molecular analyses. As the scientific community continues to explore the diversity of life beneath our feet, careful taxonomic scholarship will remain indispensable.

Declarations

Original Manuscript

This manuscript is original research, has not been previously published, and is not under consideration by any other publication.

Conflict of Interest

The author declares no conflict of interest.

Funding Statement

No external funding was received for this study.

Data Availability Statement

No new datasets were generated or analyzed during this review study.

Ethics Statement

This article does not involve human participants or animal experimentation.

Acknowledgement. The author thanks Wilma M. Reynolds, Oligochaetology Laboratory, for reviewing the manuscript, her comments and suggestions for improving the text.

REFERENCES

- Beddard, F.E. 1895. A Monograph of the Order Oligochaeta. Oxford: Claredon Press, *xii* + 769 pp.
- Blakemore, R.J. 2008. Cosmopolitan earthworms—An eco-taxonomic guide to the peregrine species of the world (4th ed.). Yokohama: VermEcology, 1 CD-ROM. OCLC: (OCoLC)183139488
- Csuzdi, C. 2012. Earthworm species, a searchable database. *Opuscula Zoologica Budapest* 43(1): 97–99.
- Dayrat, B. 2005. Towards integrative taxonomy. *Biological Journal of the Linnean Society* 85(3): 407–415. https://doi.org/10.1111/j.1095-8312.2005.00503.x
- Decaëns, T. 2010. Macroecological patterns in soil communities. *Global Ecology and Biogeography* 19: 287–302.
- Gates, G.E. 1972. Burmese Earthworms: An Introduction to the Systematics and Biology of Megadrile Oligochaetes with Special Reference to Southeast Asia. *Transactions of the American Philosophical Society* (New Series), 62(7): 1–326.
- International Commission on Zoological Nomenclature (ICZN) (Ride *et al.*, Eds.). 1999. International Code of Zoological Nomenclature. 4th Edition. London: International Trust for Zoological Nomenclature, *xxix* +306 pp.
- James, S.W., D. Porco, T. Decaëns, B. Richard, R. Rougerie and C. Erséus. 2010. DNA barcoding reveals cryptic diversity in lumbricid earthworms. *Molecular Ecology Resources* 10(5): 845–856. https://doi.org/10.1111/j.1755-0998.2010.02843.x
- Michaelsen, W. 1900. Oligochaeta. Lief. 10. *In:*

Das Tierreich. Berlin, Friedländer, *xxix* + 575 pp.

- Perez-Losada, M., M. Ricoy, J.C. Marshall and J. Domínguez. 2009. Phylogenetic assessment of the earthworm *Aporrectodea caliginosa* speciescomplex (Oligochaeta: Lumbricidae) based on mitochondrial and nuclearDNA sequences. *Molecular Phylogenetics and Evolution* 52: 293–302.
- Reynolds, J.W. and M.J. Wetzel. 2021. Nomenclatural and taxonomic issues in earthworm systematics: historical perspectives and future directions. *Megadrilogica* 25(1): 1–20.
- Rosa, D. 1891. Die exotischen Terricolen des k.k. naturhistorischen Hofmuseums. *Annalen des Naturhistorischen Museums in Wien* 6: 379–406.
- Stephenson, J. 1930. The Oligochaeta. London: Claradon Press. 978 pp.