

## Consequences of Antepartum Haemorrhage and its Perinatal Outcome

Md. Alauddin<sup>1\*</sup>, Mst. Marfia Khatun<sup>2</sup>, Sadia Sultana Mollika<sup>3</sup>, Farhana Sharmin<sup>4</sup>

<sup>1</sup>Senior Consultant (Gynaecology), 250 Bedded General Hospital, Jhenaidah, Bangladesh

<sup>2</sup>Junior Consultant (Gynaecology), 250 Bedded General Hospital, Jhenaidah, Bangladesh

<sup>3</sup>Medical Officer (Gynae), 250 Bedded General Hospital, Jhenaidah, Bangladesh

<sup>4</sup>Medical Officer (Gynae), 250 Bedded General Hospital, Jhenaidah, Bangladesh

DOI: [10.36347/sjams.2023.v11i08.025](https://doi.org/10.36347/sjams.2023.v11i08.025)

| Received: 15.07.2023 | Accepted: 19.08.2023 | Published: 28.08.2023

\*Corresponding author: Md. Alauddin

Senior Consultant (Gynaecology), 250 Bedded General Hospital, Jhenaidah, Bangladesh

### Abstract

### Original Research Article

**Background:** Antepartum haemorrhage is defined as bleeding per vaginum occurring after the fetus has reached the period of viability, its 20 weeks in developed countries and 28 weeks in developing countries. We evaluated the consequences of antepartum haemorrhage, their perinatal outcome to outline the proper management of neonate in order to improve perinatal morbidity and mortality. **Methods:** This prospective study was conducted in the Department of Obstetrics and Gynaecology, 250 Bedded General Hospital, Jhenaidah, Bangladesh from January to June 2022. APH fulfilling the inclusion criteria were studied. Data was recorded on the MS excel sheet for further analysis and processing. **Results:** Total 4595 deliveries were conducted, 150 cases ie 3.26% presented as APH. Placenta previa was most common. APH was commonly associated with multigravida in age group of 26-30 years. Maximum number of patients (40%) were in age group 26-30 years. In this study 71.3% patients were multigravida and 23.8% primigravida. Placenta previa was commonly associated with multigravidas whereas abruption was commonly seen in primigravidae. Mean gavidity was 2.24±1.17. Booking rate was 82%. Most cases of APH with placenta previa were seen in lower class and upper lower class of Modified Kuppaswamy classification. Risk factors included previous LSCS and D&C, hypertension, multiple pregnancies and malpresentations. Most patients underwent preterm LSCS. Fetal complications were due to prematurity. Overall perinatal mortality was 20.1%. **Conclusion:** Early diagnoses, timely referrals and transfusion facilities along with trained team of doctors with well-equipped ICU facility go a long way in avoiding APH related fetal complications.

**Keywords:** Antepartum Haemorrhage, Placenta Previa, Abruptio Placenta.

Copyright © 2023 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

## INTRODUCTION

Haemorrhage forms a big chunk amongst the causes of perinatal mortality and is in fact one of the gravest obstetrics emergencies whether antepartum or postpartum. Haemorrhage is a life threatening event and hence obstetrics has aptly been honoured as the business of blood.

Antepartum haemorrhage is defined as bleeding per vaginum occurring after the fetus has reached the period of viability but before the birth of baby. This implies bleeding from genital tract after 20 weeks of gestation until delivery in developed countries and 28 weeks in countries with low resource settings, lacking adequate neonatal facilities. Two to five percent of all pregnancies are complicated by antepartum haemorrhage [1, 2]. Royal College of Obstetricians and Gynaecologists (RCOG) defines blood loss by a combination of volume and signs of clinical shock to

guide management [3]. Antepartum haemorrhage can be due to placenta previa, abruptio placentae or extra placental causes, where first two accounts for more than half of the causes of APH. Placenta previa refers to the condition when the placenta is situated wholly or partially in the lower uterine segment and accounts for one third of all cases of APH. It is further classified as type I-if implantation is in lower segment but does not reach the internal os, Type II- placenta reaches the internal os but does not cover it, Type III- placenta covers the internal os but not at full dilatation, Type IV placenta covers internal os even at full dilatation of cervix. The second most common cause of APH is placental abruption which is defined as separation of the placenta-either partially or totally from its implantation site before delivery. Extra placental causes may include cervical polyps, cervical erosions, endocervical erosions, cancer of cervix, cervicitis, varicosities (vaginal, vulvar and cervical) vaginal infections, foreign bodies, genital lacerations, bloody show, degenerating

uterine myomata, vasa Previa and marginal placental separation. It may even be of undetermined origin [4]. Fetal complications are premature delivery, low birth weight, intrauterine death, congenital malformations and birth asphyxia. The latter is due to placental separation or hypotension in mother as a result of haemorrhage. Antepartum haemorrhage may frequently result in low birth weight babies. This can be an effect of preterm labour or repeated small events of haemorrhage causing chronic placental insufficiency and fetal growth retardation. The overall perinatal mortality increases to between 4-8% [5]. We therefore proposed to conduct a prospective study, to evaluate the consequences of antepartum haemorrhage, and its perinatal outcome, so as to outline the important causes and proper management of patient in order to improve perinatal morbidity and mortality and specify as to what areas required improvement in a developing country to improve the same.

## MATERIALS & METHODS

This prospective study was conducted in the Department of Obstetrics and Gynaecology, 250 Bedded General Hospital, Jhenaidah, Bangladesh from January to December 2022. 150 cases of APH were studied. All cases of antepartum haemorrhage  $\geq 28$  weeks of gestation were included in this study. Cases with APH below 28 weeks of gestation and patients suffering from any other bleeding disorders were excluded. The cases of antepartum haemorrhage with clinical findings and ultrasound reports confirming placenta previa, abruptio placenta or APH due to any other cause were admitted in the hospital. The data collected included maternal age, parity, gestational age,

presentation, booking status, education, occupation, residential address and severity of haemorrhage. The diagnoses of all cases were made on the basis of history, clinical evaluation which included general physical examination, per abdomen, per speculum, per vaginal exam and USG. Line of management i.e conservative or emergency lower segment caesarean section was planned according to the clinical condition of the patient and fetus. All facilities of neonatal intensive care unit (NICU) to deal with preterm and sick infants were available in paediatric unit of this institution. Data collected was transformed into MS excel sheet for further processing and analysis.

## RESULTS

**Table 1: Types of APH, (N=150)**

Total no of deliveries	4595
No of APH cases	150
Types of APH	
Abruptio Placentae	(35) 23.3%
Placenta Previa	(88) 58.7%
Undetermined	(27) 18.0%

Out of total 4595 deliveries a total 150 patients were admitted with APH. 150 patients fulfilled the inclusion criteria. The incidence of APH in our institute is 3.26%. Among 150 deliveries there were 4 twins and 1 triplet so the total number of babies delivered were 150. The comparison between causes of APH and its consequences was studied. Out of total cases, 58.7% were placenta previa, 23.3% abruptio placentae and 18.0% undetermined (Table-1).

**Table 2: Age distribution of APH, (N=150)**

Age (Years)	Age distribution (Years)	Abruptio placentae	Placenta previa	Undetermined
<20	4(50%)	1(15.5%)	3(37.5%)	0(0.0%)
20-25	25(45.5%)	9(16.4%)	11(20.0%)	10 (18.2%)
26-30	22(46.8%)	6(12.8%)	14(29.8%)	5(10.6%)
31-35	14(51.9%)	6(22.2%)	7(25.9%)	0 (0.00%)
>35	4(36.4%)	1(9.1%)	5(45.5%)	1(9.1%)
<b>Gravidity</b>				
1	16(41.0%)	7(17.9%)	10(25.6%)	6(15.4%)
2	25(44.6)	4(7.1%)	16(28.6%)	11(19.6%)
3	14(43.8)	5(11.6%)	11(25.6%)	2(4.7%)
$\geq 4$	8(34.8)	2(8.7%)	10(43.5%)	3(13.0%)
<b>Booking Status</b>				
Booked	56(47.1%)	11(9.2%)	37 (31.1%)	15(12.6%)
Un- booked	15(48.4%)	4(12.9%)	8 (25.8%)	4 (12.9%)

Maximum number of patients (40%) were in age group 26-30 years. In this study 71.3% patients were multigravida and 23.8% primigravida. Placenta previa was commonly associated with multigravidas whereas abruption was commonly seen in primigravidae. Mean gravidity was  $2.24 \pm 1.17$ . Booking

rate was 82%. Most cases of APH with placenta previa were seen in lower class and upper lower class of Modified Kuppaswamy classification. Abruptio was also commonly associated with lower and lower middle class (Table-2).

**Table 3: Relation of gestation to APH, (N=150)**

Gest Age at admission	No. of Cases	28-30w	31-33w	34-37w	>37w
Abruptio Placentae	(21) 52.5%	3(7.5%)	7(17.5%)	6(15%)	3(7.5%)
Placenta Previa	(28) 36.8%	10(13.2%)	12(15.8%)	21(27.6%)	5(6.6%)
Undetermined	(14) 41.2%	5(14.7%)	4(11.8%)	6(17.6%)	5(14.7%)
Gest Age at delivery					
Preterm	(54)56.8.1%	7 (7.4%)	9 (9.5%)	25 (26.3%)	0 (0.0%)
Term	(22) 40.0%	0 (0.0%)	0 (0.0%)	12 (21.8%)	21 (38.2%)

Table-3 shows that 77% of APH cases had mild bleeding at presentation out of which 58% were placenta previa. Severe bleeding was observed in 7% cases and 33% of which were due to abruption placentae. 21% cases of placenta previa were admitted

at 34-37 weeks of gestation and abruptio placenta at 31-33 weeks gestation. 26% of the APH delivered between 34-37 weeks of gestation whereas 40% cases delivered at gestation >37 weeks.

**Table 4: Risk Factors, (N=150)**

Previous history	No. of Patients	Abruptio placentae	Placenta previa	Undetermined
Abortion	26(39.4%)	6 (9.1%)	19 (28.8%)	15 (22.7%)
Abortion and D&C	17(40.5%)	2 (4.8%)	13 (31.0%)	10 (23.8%)
LSCS	15(42.9%)	4(11.4%)	16 (45.7%)	0 (0.00%)
Abortion, LSCS and D&C	5(71.4%)	1 (14.3.00%)	1 (14.3.00%)	0 (0.00%)
<b>Risk Factors</b>				
Hypertension	26 (53.1%)	16 (32.7%)	6 (12.2%)	1 (2.0%)
Multiparity	9 (56.3%)	2 (12.5%)	5 (31.3%)	0 (0.00%)
Twins	5 (41.7%)	4 (33.3%)	3 (25%)	0 (0.00%)
Malpresentations	7 (53.8%)	0 (0.00%)	4 (30.8%)	2 (15.4%)
IUGR	8 (47.1%)	3 (17.6%)	3 (17.6%)	3 (17.6%)
Polyhydramnios	7 (46.7%)	5 (33.3%)	0 (0.0%)	3 (20.0%)
Hypothyroidism	7 (50.0%)	4(28.6%)	2 (14.3%)	1 (7.1%)
Elderly	5 (50.0%)	0(0.00%)	3 (30%)	2 (20%)
Elderly	2 (50.0%)	2 (50.0%)	0 (0.00%)	0 (0.00%)

Table-4 shows that among 76% patients with previous LSCS and 64% with history of dilatation and curettage had placenta previa. Hypertension was noted in 53% of the patients with APH. Out of these 32.7%

patients had abruption. Multiparity, malpresentations, polyhydramnios and hypothyroidism were also studied as associated risk factors.

**Table 5: Mode of delivery, (N=150)**

Mode of delivery	No of Patients	Abruption	Placenta Previa	Undetermined
LSCS	96(64%) Preterm-66 Term- 30	25 (26.1%)	63 (65.6%)	8 (8.3%)
Vaginal Delivery	54(36%) Preterm-24 Term -30	10 (26%)	25 (46%)	19 (35%)

Table-5 shows that 64% of the patients underwent LSCS out of which 65.6% were placenta previa. 36% of the patients had vaginal delivery out of which 46% were placenta previa (Type 1). 59% of the

patients had preterm deliveries out of which 73% were LSCS and 41% of the term deliveries had 50% LSCS rate.

**Table 6: Fetal complications, (N=150)**

Fetal Complication	No. of Cases	Abruptio placentae	Placenta previa	Undetermined
Preterm	30(50.8%)	6(10.2%)	20(33.9%)	3(5.1%)
Asphyxia	14(48.3%)	4(13.8%)	8(27.6%)	3(10.3%)
IUD	12(52.2%)	5(21.7%)	5(21.7%)	1(4.3%)
Jaundice	8(47.1%)	4(23.5%)	3(17.6%)	2(11.8%)
Distress	4(50.0%)	2(25%)	1(12.5%)	1(12.5%)
Still Birth	3(50.0%)	3(50.0%)	0(0%)	0(0%)
Fever	3(50.0%)	2(33.3%)	1(16.7%)	0(0%)
Hypoglycemia	1(50.0%)	0(0%)	1(50.0%)	0(0%)

Table-6 shows that 50.8% neonates had preterm deliveries, out of which 33.9% were placenta

previa. 48.3% cases were associated with asphyxia out of which 27.6% were placenta previa.

**Table 7: Relation of birth weight and APH (N=150)**

Birth Weight	No. of Cases	Abruptio placentae	Placenta previa	Undetermined
<1	2(50%)	1(25%)	1(25%)	0(0%)
1-1.5	12(42.9%)	6(21.4%)	7(25%)	3(10.7%)
1.6-2	15(44.1%)	5(14.7%)	12(35.3%)	2(5.9%)
2.1-2.5	17(42.5%)	5(12.5%)	14(35%)	4(10%)
2.6-3	17(42.5%)	6(15%)	11(27.5%)	6(15%)
>3	2(50.0%)	2(50.0%)	0(0%)	0(0%)
<b>APGAR</b>				
0-3	14(53.8%)	7 (26.9%)	5 (19.2%)	0(0%)
4-7	18(42.9%)	8 (19.0%)	12(28.8%)	4 (9.5%)
>7	35(42.7%)	10 (12.2%)	25(30.5%)	12 (14.6%)

Table-7 shows that majority of neonates weighed between 2.1-2.5 Kgs. Low birth weight was due to prematurity. Mean 2.17+36 Kgs. 66% of the neonates had AP GAR score>7. Abruptio placenta had

significant relationship with low AP GAR score. 75% of NICU admissions were due to fetal distress out of which 48% cases were of placenta previa and 36% abruptio placenta.

**Table 8: Fetal outcome, (N=150)**

Outcome	Total	Abruption	Placenta Previa	Undetermined
Live	53(49.1%)	12(11.1%)	30(27.8%)	13(12.0%)
IUD	8(47.1%)	4(23.5%)	4(23.5%)	1(5.9%)
Still Birth	3(50.0%)	3(50.0%)	0(0.00%)	0(0.00%)
Expired in NICU	12(63.2%)	2(10.5%)	5(26.3%)	0(0.00%)
<b>Cause of death</b>				
Hypoglycemia, Seizures	6(85.7%)	0(0.00%)	0(0.00%)	1(14.3%)
RDS	16(88.9%)	1(5.6%)	1(5.6%)	0(0.00%)
Sepsis, Shock	30(62.5%)	7(14.6%)	8(16.7%)	3(6.3%)
Shock	24(49.0%)	14(28.6%)	10(20.4%)	1(2.0%)
Shock, Pulmonary Haemorrhage	11(39.3%)	14(50.0%)	3(10.7%)	0(0.00%)

Table-8 shows that out of the total 150 deliveries, 53 were live neonates, 12 expired in NICU. 13 were intrauterine deaths and 4 were still Births. Neonatal deaths in toto were 28 i.e 20.1%. Overall deaths of neonates illustrates the most common cause is neonatal shock seen in 46% followed by sepsis and shock seen in 32% neonates.

## DISCUSSION

Incidence of APH in our study was 3.26%. This was in accordance with the findings of Kedar K *et al.*, Bhandiwad A *et al.*, Bhide A *et al.*, and Arora A *et al.*, [6-9]. In our study 82% patients of APH were booked and 18% were unbooked. Tyagi P *et al.*, and Bhandiwad A *et al.*, reported higher incidences of unbooked cases in their studies. The inconsistency noted maybe because of the free facilities provided by the state government giving an easy access for antenatal checkup [7, 10]. Placenta previa was the main cause of APH i.e. 58.5%. Abruptio placentae and undetermined cases were 23.3% and 18.0% respectively. Our results were consistent with the studies conducted by Adekanle DA *et al.*, and Jejani a *et al.*, Mean age of women presenting with APH in our study was 26.7+4.5 years

with 40% in age group of 26-30 years and 39% between 20-25 years age. This is also consistent with studies conducted by Tyagi P *et al.*, and Adekanle DA *et al.*, in which 61% and 40% cases of APH were observed between 26-30 years of age respectively [10-12]. The mean gestational age at delivery in our study was 35.22+2.82 weeks and 60.1% of the patients in our study had preterm deliveries which was almost similar to studies conducted by Samal SK *et al.*, and Singhal S *et al.*, This higher rate of preterm deliveries could be due to early decisions and surgical intervention in view of maternal health [13, 14]. Hypertension was seen in 16% APH cases, out of which 68.8% cases had abruption. This was consistent with the studies conducted by Sarwar I *et al.*, and Bhandiwad A *et al.*, In our study 19% APH cases had abortions, 22% abortions and 12% patients had previous LSCS which was consistent with the study conducted by Samal SK *et al.*, Sekiguchi A *et al.*, also reported similar a incidence [5, 13, 15, 16] . Malpresentations were seen in 5.2% cases which was consistent with the findings of Samal SK *et al.*, [13], Jejani A *et al.*, [12] and Sheikh F *et al.*, [13, 17]. Prematurity was the most common cause of LBW which was seen in 70% of the neonates

weighing <2.5 kgs. 30% of neonates had birth weight >2.5Kgs. Mean birth weight was 2.17+36Kgs. Our findings were consistent with Samal SK *et al.*, Sharmila G *et al.*, and Sekiguchi A *et al.*, [13, 16, 18]. In our study 66% neonates had APGAR score above 7 at 5 mins of birth while 34% had scores less than 7. Our results were consistent with the findings of Adekunle DA *et al.*, Rajini P *et al.*, Singhal SR *et al.*, with 74.6%, 80.4% incidences of APGAR score >7 at 5 mins [11, 14, 19]. In our study the most common fetal complications observed were prematurity 50.8%, fetal distress 30%, asphyxia 48.3%, jaundice 47.1%, sepsis 2.1% still birth 50.0%, IUD 52.2%, early neonatal death 5.7%. This is consistent with the studies conducted by Kedar K *et al.*, Samal SK *et al.*, Taylor F *et al.*, and Bhandiwad A *et al.*, [7, 13, 20, 21] Perinatal mortality was observed in 20.1% of cases of APH in our study. Most common causes were prematurity related complications. Neonatal death rate was 8.6%, 11.5% cases were of IUD and these were most commonly seen with abruption followed by placenta previa. Results were similar to studies conducted by Singhal S *et al.*, and Maurya a *et al.*, Chufamo N *et al.*, calculated the perinatal mortality to be 36.9% while Arora R *et al.*, reported very high mortality rate of 61.5%. The lower neonatal death in our study may be due better neonatal care facilities with well-equipped NICU at our institute [9, 14, 18].

## CONCLUSION

Antepartum haemorrhage is a leading cause of perinatal morbidity and mortality. Placenta previa and abruptio placentae are the commonest type of APH with hypertension and multiparity being the major risk factors for APH. Most common complications include preterm deliveries and low birth weight due to early intervention. Antenatal patients with APH must be considered a high risk pregnancy. Early diagnoses, timely referrals and transfusion facilities can also aid in decreasing perinatal morbidity and mortality. Above all awareness of antenatal care during pregnancy, knowledge of various government schemes for antenatal patients, importance of institutional deliveries and adoption of various contraceptive methods (temporary as well as permanent) are the key factors to avoid various complications due to APH.

**Funding:** No funding sources.

**Conflict of Interest:** None declared.

## REFERENCES

1. Amitava, R. S., Saikat, S. R., Biswajit, N., Gourab, M., & Jayanta, M. (2010). Management of obstetric haemorrhage. *Med J Anaes*, 20(4), 499-508.
2. Mustafa Adelaja, L., & Olufemi Taiwo, O. (2011). Maternal and fetal outcome of obstetric emergencies in a tertiary health institution in South-Western Nigeria. *International Scholarly Research Notices*, 2011, 1-4 doi:10.5402/2011/160932.
3. RCOG. Antepertum Haemorrhage clinical Greentop Guidelines no 63. The Royal College of Obstetricians and Gynaecologists, 2011.
4. Arias, F., Daftary, S. N., & Bhine, G. A. (2008). Bleeding during pregnancy. In: Practical Guide to High-Risk Pregnancy and Delivery, 3 rded. New Delhi: Elsevier, Chapter, 13, 323-357.
5. Mabic, W. C. (1992). Placenta previa. *Clin Perinatol*, 19(2), 425-435.
6. Kedar, K., Uiquey, P., Pawar, A., & Choudhary, A. (2016). Maternal and fetal outcome in antepartum haemorrhage: a study at tertiary care hospital. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*, 5(5), 1386-1394.
7. Bhandiwad, A., & Bhandiwad, A. A. (2014). A Study of maternal and fetal outcomes in antepartum haemorrhage. *J Evidence Based Med Healthcare*, 1(6), 406-427.
8. Bhide, A., & Thilaganathan, B. (2004). Recent advances in the management of placenta previa. *Current Opinion in Obstetrics and Gynecology*, 16(6), 447-451.
9. Arora, R., Devi, U., & Majumdar, K. (2001). Perinatal morbidity and mortality in antepartum haemorrhage. *J Obstet Gynaec India*, 51(3), 102-104.
10. Tyagi, P., Yadav, N., Sinha, P., & Gupta, U. (2016). Study of antepartum haemorrhage and its maternal and perinatal outcome. *Int J Reprod Contracept Obstet Gynecol*, 5(11), 3972-3977.
11. Adekanle, D. A., Adeyemu, A., & Fadero, F. (2011). Antepartum haemorrhage and pregnancy outcome. *J Med and Med Sci*, 2(12), 1243-1247.
12. Jejani, A., & Kawthalkar, A. (2015). Study of obstetric outcome in antepartum haemorrhage. *Panacea Journal of Medical Science*, 5(3), 153-157.
13. Singhal, S., Nymphaea., & Nanda, S. (2007). Maternal and perinatal outcome in Antepartum Haemorrhage: A study at a Tertiary Care Referral Institute. *Internet J Gynaecol obstet*, 9(2), 1-4.
14. Samal, S. K., Rathod, S., Rani, R., & Ghose, S. (2017). Maternal and perinatal outcomes in cases of antepartum haemorrhage: a 3-year observational study in a tertiary care hospital. *Int J Reprod Contracept Obstet Gynecol*, 6(3), 1025-1029.
15. Sarwar, I., & Islam, A. (2006). Abruptio placentae and its complications at Ayub Teaching Hospital Abbottabad. *Journal of Ayub Medical College Abbottabad*, 18(1), 27-31.
16. Sekiguchi, A., Nakai, A., Kawabata, I., Hayashi, M., & Takeshita, T. (2013). Type and location of placenta previa affect preterm delivery risk related to antepartum hemorrhage. *International journal of medical sciences*, 10(12), 1683-1688.
17. Sheikh, F., Khokhar, S. A., Sirichand, P., & Shaikh, R. B. (2010). A study of antepartum haemorrhage: Maternal and Perinatal outcomes. *Medical Channel*, 16(2), 268-271.

18. Singhal, S., Nymphaea, N. S., & Nanda, S. (2008). Maternal and perinatal outcome in antepartum hemorrhage: A study at a tertiary care referral institute. *Internet J Gynecol Obstet*, 9(2), 5580.
19. Rajini, P., Devi, A. V., Singh, A., & Grishma, P. (2016). Maternal and Perinatal Outcome in Cases of Antepartum Hemorrhage in Gandhi Hospital. *Journal of Dental and Medical Sciences*, 15(6), 9-11.
20. Singhal, S. R., Chaudhary, S., Nanda, S., & Dahiya, P. (2016). Characteristics of patients of ante-partum hemorrhage at a tertiary care centre. *Indian Journal of Obstetrics and Gynecology Research*, 3(2), 86-89.
21. Tikkanen, M., Nuutila, M., Hiilesmaa, V., Paavonen, J., & Ylikorkala, O. (2006). Clinical presentation and risk factors of placental abruption. *Acta obstetrica et gynecologica Scandinavica*, 85(6), 700-705.