

To Study the Effect of Intravenous Clonidine on Attenuation of Hemodynamic Stress Response of Laryngoscopy and Intubation in Surgeries under General Anaesthesia

Ritu Masar¹, Anju Gautam^{2*}, Rajan Godwin³, Sachin Gajbhiye⁴

¹Senior Resident, Department of Anaesthesiology, M.G. Medical College, Indore, M.P., India

²Assistant Professor, Department of Anaesthesiology, G.R. Medical College, Gwalior, M.P., India

³Associate Professor, Department of Anaesthesiology, N.S.C.B. Medical College, Jabalpur, M.P., India

⁴Consultant Anaesthesiologist, N.S.C.B. Medical College, Jabalpur, M.P., India

Original Research Article

*Corresponding author

Anju Gautam

Article History

Received: 06.12.2018

Accepted: 15.12.2018

Published: 30.12.2018

DOI:

10.36347/sjams.2018.v06i12.012



Abstract: Endotracheal intubation is always associated with certain hazards. Few common complications are trauma during laryngoscopy, airway obstruction by kinking or malposition of tube, trauma to vocal cords leading to oedema and postoperative hoarseness and ulceration are not uncommon. Besides these, there are certain cardiovascular disturbances, which often go unnoticed due to lack of proper monitoring cardiac disturbances may be transient and may not lead to deleterious effects in normal healthy individuals but can lead to serious consequences in patient with cardiovascular diseases. Various methods are used to attenuate this hemodynamic response. In present study, we have used i.v. clonidine just before induction of anaesthesia to attenuate this reflex response. This randomized control study was conducted on 60 patients of ASA grade 1 and 2 after taking ethics committee approval. Patients were divided into two groups. In Group 1 (n=30), patients received i.v. clonidine just before induction of anaesthesia and Group 2 (n=30), patients received Placebo. Diastolic, systolic and mean blood pressure and heart rate were measured throughout the procedure and recorded for five minutes just after the intubation. i.v. clonidine given pre-emptively helps attenuating adverse hemodynamic response occurring due to laryngoscopy ($p < 0.05$) in comparison to control group. i.v. clonidine given pre-emptively reduces adverse hemodynamic response occurring due to laryngoscopy.

Keywords: Clonidine, Hemodynamic response, Laryngoscopy, General Anaesthesia.

INTRODUCTION

Endotracheal intubation is considered to be gold standard during surgeries and also life saving procedure during emergency condition. With the introduction of new equipment and neuromuscular blockers in the anaesthetic practice coupled with ever-growing technical knowledge and skills of anesthesiologists, endotracheal intubation has become a safe and common Procedure in modern day Anaesthesia.

Endotracheal intubation is not free of complications. Some are minor and some are major like trauma during laryngoscopy, airway obstruction by kinking or malposition of tube, trauma to vocal cords leading to oedema and postoperative hoarseness and ulceration are not uncommon. Besides these, there are certain cardiovascular disturbances, which often go unnoticed due to lack of proper monitoring cardiac disturbances may be transient and may not lead to

deleterious effects in normal healthy individuals but can lead to serious consequences in patient with cardiovascular diseases.

Cardiovascular complications namely hypertension, arrhythmias and tachycardia arising during laryngoscopy and transient disturbances in cardiac action are caused by reflex excitation of vagus nerves or sympathetic system during laryngoscopy and intubation are of special concern [1-3]. Reflex circulatory responses is more marked when intubation is done under light plane of anaesthesia [2, 4].

Several pharmacological methods are used to attenuate the blood pressure and heart rate elevation and response to laryngoscopy and intubation. These included topical anaesthesia of the oropharynx [5], intravenous lignocaine [6, 7], adrenergic blocking drugs, vasodilating drugs etc. The response can be diminished or modified locally, centrally or peripherally

and attempts have been made to accomplish this using all approaches with varying success. Regional topical anaesthesia used to block afferent impulses.

But most of the method required laryngoscopy itself produces similar response. Intravenous lignocaine and deeper inhalational techniques are used to modify the response at the central nervous system.

Other pharmacological approaches are fentanyl [8], beta blocker [9], Alfentanyl [10], calcium channel Blocker like nifedipine [11], MgSo₄ [12], Diltriazem [13], nitroglycerine [14], Buprenorphine [15], Esmolol etc. but these approaches are not proved to be entirely satisfactory because reflex action is not completely blocked.

Clonidine, an alpha-2 adrenergic agonist, initially introduced as centrally acting antihypertensive agent has been shown to reduce sympathetic out flow in response to stress. It has been in use for treatment of mild and moderate hypertension. This drug act as the reaction site in the medulla oblongata and pre-synaptically at peripheral nerve terminals to cause a reduction in the activity of sympathetic nervous system.

MATERIALS AND METHODS

The present study was conducted in NSCB Medical College and Hospital, Jabalpur. The study was carried out on 60 adult patients of either sex, in the age group 18-65 years. All the patients were normotensive and devoid of any cardiovascular disturbances, belonging to ASA grade I and II. The patients were picked up from the routine operative list scheduled to undergo various types of operating procedures like general surgical, gynaecological and orthopaedic.

The patients who required administration of general anaesthesia for the particular surgery were included in the study. Any patients who gave history of receiving antihypertensive, antiarrhythmic and steroids were excluded from the study. Patients with systolic BP of 140 mmHg or less and diastolic BP of 90 mmHg or less were considered as normotensive and only such patients were included in the study. Patients with any abnormally in the ECG were not included in the study.

All the patients were examined the day prior to the surgery. Informed consent was obtained and the patients were briefed about the drug regime and the method of administration so that full cooperation of the patients was achieved. Thorough general, physical and systemic examination was carried out to rule out any systemic disorders. All routine and special investigations of blood and urine were carried out. Chest skiagrams, electrocardiogram, ultrasonography etc. were done as per indication.

On the morning of operation, the patients were shifted to the operation theatre. Systolic blood pressure, diastolic blood pressure, and heart rate were recorded.

Patients were divided into 2 groups of 30 patients each. Depending upon the drugs employed to attenuate the cardiovascular responses during laryngoscopy and intubation, patients were randomly allotted to either group. The groups were designated as I and II.

Group I (n=30) the patients of this group received only normal saline and served as control group.

Group II (n=30) the Patients of this group received IV clonidine 3 µg/kg before induction.

ANAESTHESIA TECHNIQUE

In all the groups, the pre-medication with parasympatholytics were avoided as they could cause tachycardia which would invalidate the results. No other pre-medicant such as sedative or narcotic was given to any patient.

All the patients were preoxygenated with 100% oxygen for three minutes. Technique of anaesthesia was standardized for all the patients in the study. This consisted of induction with I.V. thiopentone sodium (2.5) 5mg/kg BW followed by I.V. succinylcholine 1.5 mg/kg BW to facilitate endotracheal intubation.

Gentle ventilation with 100% oxygen was done till cessation of fasciculation thereafter gentle laryngoscopy was performed and endotracheal intubation was done with cuffed endotracheal tube of adequate size. The cuff was immediately inflated. The endotracheal tube was then connected to the Boyles machine through Bain's circuit. Nitrous oxide: oxygen (60:40), halothane and non depolarising muscle relaxant was used for maintenance of anaesthesia. No kind of stimulation like urinary Catheterization, Ryle tube insertion cleaning draping, insertion, of additional IV line, IM injections will be allowed during the study period.

To study the effect of clonidine on cardiovascular changes during laryngoscopy and intubation, pulse rate and blood pressure were recorded at fixed time intervals. The observation were made as under:

- Just before induction
- Just before laryngoscopy
- Immediately after intubation
- One minute after intubation
- Two minutes after intubation
- Four minutes after intubation
- Five minutes after intubation

The parameter like SBP, DBP, and PR were recorded every minute after ET till up to 5 min. Patients were continuously monitored for arterial oxygen saturation. Any period of desaturation, gross fluctuation of blood pressure on either side and any abnormality in the pulse rhythm was noted and recorded.

All the relevant data was recorded in proforma prepared for the study and results thus obtained were subjected to statistical analysis by paired t-test and Z test where ever applicable and degree of significance was obtained.

RESULTS

Data obtained from the patients involved in the study were analyzed. The mean age, weight, sex, duration of anaesthesia and surgery were comparable in two groups as shown in table-1.

Preoperative heart rate, systolic, diastolic and mean blood pressures were comparable in both the groups.

Table-1: showing demographic variables of two groups

Demographic data	Group 1	Group 2
Age	37.06±10.63	37.90±10.20
Weight (Kg)	58.36±12.15	57.60±9.56
Sex (Female)	76.7%	86.7%
Duration Of Anaesthesia (Min)	97.16±17.05	98.00±20.82

Table-2: Showing pulse rate at various time interval in study groups

GROUP		PR_POP	PR_BI	PR_BL	PR_AI	PR1	PR2	PR3	PR4	PR5
I	Mean	86.9	91.2	99.4	112.6	113.3	110.1	106.6	104.9	101.7
	SD	±10.2	±9.4	±10.5	±10.9	±7.4	±7.2	±8.0	±9.0	±8.4
	N	30	30	30	30	30	30	30	30	30
t value			1.709	4.698	9.424	11.480	10.193	8.354	7.282	6.146
P			P>0.05	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001
II	Mean	82.6	76.2	82.0	88.5	86.3	83.6	80.9	78.3	75.5
	SD	±8.4	±7.3	±6.8	±6.2	±5.9	±6.0	±6.1	±5.9	±5.8
	N	30	30	30	30	30	30	30	30	30
t value			3.144	0.302	3.097	1.981	0.530	0.911	2.303	3.776
P			0.005	P>0.05	0.005	0.05	P>0.05	P>0.05	0.05	0.001

Table-3: Showing systolic arterial pressure at various time interval in study groups

GROUP		SAP_POP	SAP_BI	SAP_BL	SAP_AI	SAP1	SAP2	SAP3	SAP4	SAP5
1	Mean	125.3	125.7	126.5	163.7	154.3	143.0	137.7	133.5	128.1
	SD	±8.7	±11.3	±12.7	±16.1	±14.4	±12.2	±12.1	±10.5	±9.9
	N	30	30	30	30	30	30	30	30	30
t value			0.153	0.427	11.498	9.387	6.453	4.546	3.289	1.135
P			P>0.05	P>0.05	0.0001	0.0001	0.0001	0.0001	0.001	9999
2	Mean	126.3	117.1	110.6	119.2	116.9	114.5	112.1	109.9	107.1
	SD	±10.7	±8.5	±7.5	±7.2	±6.8	±6.1	±5.8	±5.6	±5.3
	N	30	30	30	30	30	30	30	30	30
t value			3.712	6.599	3.028	4.082	5.263	6.390	7.425	8.821
P			0.001	0.0001	0.005	0.0001	0.0001	0.0001	0.0001	0.0001

Table-4: Showing diastolic arterial pressure at various time interval in study groups

GROUP		DAP_POP	DAP_BI	DAP_BL	DAP_AI	DAP1	DAP2	DAP3	DAP4	DAP5
1	Mean	81.9	82.1	83.5	104.9	97.2	92.3	88.3	85.6	82.9
	SD	±5.6	±6.1	±7.0	±7.3	±6.8	±6.3	±5.9	±5.6	±5.5
	N	30	30	30	30	30	30	30	30	30
t value			0.176	0.975	13.675	9.547	6.795	4.344	2.594	0.702
P			P>0.05	P>0.05	0.0001	0.0001	0.0001	0.0001	0.01	P>0.05
2	Mean	80.1	74.7	71.7	77.7	75.7	73.9	71.7	70.1	68.1
	SD	±7.0	±6.3	±5.4	±5.7	±5.2	±4.5	±4.1	±3.5	±3.5
	N	30	30	30	30	30	30	30	30	30
t value			3.171	5.210	1.456	2.769	4.118	5.682	7.055	8.382
P			0.005	0.0001	P>0.05	0.01	0.0001	0.0001	0.0001	0.0001

DISCUSSION

The present study entitled "To Study The Effect of Intravenous Clonidine on Attenuation of Hemodynamic Stress Response of Laryngoscopy And Intubation In Surgeries Under General Anaesthesia" was conducted on 60 patients of ASA grade I and II of either sex of age group 18 - 50 years scheduled for surgeries under general anesthesia who were randomly divided into two groups according to the drugs received as shown below:

Group I (n=30) the patients of this group received only normal saline and served as control group.

Group II (n=30) the patients of this group received IV clonidine 3 μ g/kg before induction.

Endotracheal intubation is always associated with certain hazards. Few common complications are trauma during laryngoscopy, airway obstruction by kinking or malposition of tube, trauma to vocal cords leading to oedema and postoperative hoarseness and ulceration are not uncommon. Besides these, there are certain cardiovascular disturbances, which often go unnoticed due to lack of proper monitoring cardiac disturbances may be transient and may not lead to deleterious effects in normal healthy individuals but can lead to serious consequences in patient with cardiovascular diseases.

Later workers took pharmacological method to attenuate the blood pressure and heart rate elevation and response to laryngoscopy and intubation. These included topical anaesthesia of the oropharynx, intravenous lignocaine, adrenergic blocking drugs, vasodilating drugs etc. The response can be diminished or modified locally, centrally or peripherally and attempts have been made to accomplish this using all approaches with varying success. Regional topical anaesthesia used to block afferent impulses.

But most of the method required laryngoscopy itself produces similar response. Intravenous lignocaine and deeper inhalational techniques are used to modify the response at the central nervous system.

Other pharmacological approaches are fentanyl, beta blocker, Alfentanyl, calcium channel Blocker like nifedipine, MgSo₄, Diltriazem, nitroglycerine, Buprenorphine, Esmolol etc.

These pharmacological approaches are not proved to be entirely satisfactory because reflex action is not completely blocked.

Clonidine, an alpha-2 adrenergic agonist, initially introduced as centrally acting antihypertensive agent has been shown to reduce sympathetic out flow in response to stress. It has been in use for treatment of mild and moderate hypertension. This drug act as the

reaction site in the medulla oblongata and pre-synaptically at peripheral nerve terminals to cause a reduction in the activity of sympathetic nervous system.

All the demographic data like age, sex, weight of the patients, duration and type of the surgery were comparable in both the groups i.e. statistically insignificant ($p > 0.05$) as shown in table-1.

Pulse rate was significantly lower and maintained in clonidine group as compared to the control group as shown in table-2 and results were statistically significant ($p < 0.05$).

Blood BC and Flacke WE [16] studied the reduction in Halothane Anaesthetic requirement by clonidine, an alpha-adrenergic agonist and shown that clonidine causes marked increase in halothane anaesthetic requirement in the dogs. The effect is mediated through an alpha-adrenergic mechanism.

Batra *et al.*, [17], conducted a double blind randomized trial to study the effect of clonidine on pulse rate and blood pressure response to laryngoscopy and tracheal intubation. The increase was significantly lower in the clonidine treated group immediately after intubation ($P < 0.001$). The authors concluded that the rise in heart rate blood pressure associated with laryngoscopy and intubation during a routine induction sequence can be altered by the use of oral clonidine.

Wright *et al.*, [18], evaluated the efficacy of oral clonidine in a dose of 0.3mg as a routine premedicant compared with placebo. In the clonidine treated patients, heart rate was reduced than the control group and then persisted through induction and maintenance of anaesthesia.

Nishikawa *et al.*, [19], evaluated the effects of oral clonidine premedication on the haemodynamic changes associated with laryngoscopy and tracheal intubation. There was also a significant difference between the two groups in the incidence of systolic blood pressure increase above 180mmHg following laryngoscopy and tracheal intubation (0% versus 26%, $p < 0.05$). However, no significant difference was noticed between the two groups in heart rate response to laryngoscopy and tracheal intubation. The authors concluded that oral clonidine 5 mcg.kg.⁻¹ as a preanaesthetic medication could attenuate the presser responses associated with laryngoscopy and tracheal intubation.

Systolic and diastolic arterial blood pressures were significantly lower and maintained in clonidine group as compared to the control group as shown in table-3 and 4 and results were statistically significant ($p < 0.05$).

Carbine UA *et al.*, [20] studied the effect of clonidine on the pressor and heart rate response to endotracheal intubation. 30 patients were pretreated with either clonidine 1.25 μ g/Kg or clonidine 0.625 μ g/Kg or an equivalent volume of normal saline, given IV 15 mins before induction of anaesthesia. The attenuation of the pressor response to intubation of both clonidine groups was statistically significant compared to the saline group.

Dorothee M. Gaumann *et al.*, [21] This study was conducted to examine the haemodynamic and endocrine effects of clonidine, given as sole preanaesthetic medication, in neurosurgical patients. Though statistically significant, the observed inhibitory haemodynamic and endocrine effects of clonidine seem to be of minor clinical importance.

Leslie *et al.*, [22], conducted a randomized double blind control trial to investigate the influences of intravenous clonidine on thiopentone dose requirements, when used for induction of anaesthesia and associated haemodynamic effects. Significant decreases in thiopentone doses were observed in group receiving clonidine compared with control group.

Carbine UA, Allen *et al.*, [23] studied the effects of IV clonidine, fentanyl and saline for both their effect on the cardiovascular response to intubation and early postoperative pain. The increase in heart rate with intubation was significantly lower in the fentanyl and clonidine groups compared to the control group ($P < 0.05$) and these changes persisted for 90 seconds after intubation.

Howie *et al.*, [24], examined the effects of oral clonidine premedication on anaesthetic requirement, hormonal response haemodynamic and recovery in coronary artery bypass graft surgery patients. The authors concluded that oral clonidine decrease opioid use and lowers hormonal response while maintaining haemodynamics in patients undergoing coronary artery bypass graft surgeries with sufentanil anaesthesia.

Ramesh *et al.*, [25], studied the efficacy of clonidine in children with respect to sedation, intubation response and recovery. The authors concluded that clonidine 3mcg.kg-1 produces sedation comparable to diazepam 0.2mg.kg-1 and also attenuates the intubation response without increasing the incidence of complications.

CONCLUSION

This study was carried out to study the effect of pre-emptive doses of i.v. clonidine on attenuation of hemodynamic stress response of laryngoscopy and intubation in surgeries under general anaesthesia. We found that i.v. clonidine, when given just before anaesthesia induction reduces hemodynamic changes

like hypertension and tachycardia associated with laryngoscopy and intubation.

REFERENCES

1. Burstein CL, Lopinto, FJ, Newman W. Electrocardiographic studies during endotracheal intubation. Effects during usual routine technique. *Anesthesiology*, 1950; 1224-234.
2. King BD, Harris LC, Greifenstein FE. Reflex circulatory responses to direct laryngoscopy and tracheal intubation performed during general anaesthesia. *Anesthesiology*, 1951; 12:556-66.
3. Reid LC, Brace DE. Irritation of the respiratory tract and its reflex effect upon the heart. *Surg Gynecol Obstet*. 1940; 70:157.
4. Cummings MF, Russell WJ, Frewin DB. Effects of Pancuronium and Alcuronium on The Changes In Arterial Pressure And Plasma Catecholamine Concentrations During Tracheal Intubation. *BJA: British Journal of Anaesthesia*. 1983 Jul 1;55(7):619-23.
5. Stoelting RK. Blood pressure and heart rate changes during short-duration laryngoscopy for tracheal intubation: influence of viscous or intravenous lidocaine. *Anesthesia & Analgesia*. 1978 Mar 1;57(2):197-199.
6. Denlinger JK, Ellison N, Ominsky AJ. Effects of intratracheal lidocaine on circulatory responses to tracheal intubation. *Anesthesiology: The Journal of the American Society of Anesthesiologists*. 1974 Oct 1;41(4):409-12.
7. Hamill JF, Bedford RF, Weaver DC, Colohan HR. Lidocaine before endotracheal intubation: intravenous or larygotracheal? *Anaesthesiology*. 1981; 55: 578-81.
8. Helfman SM, Gold MI, DeLisser EA, Herrington CA. Which drug prevents tachycardia and hypertension associated with tracheal intubation: lidocaine, fentanyl, or esmolol?. *Anesthesia and analgesia*. 1991 Apr;72(4):482-6.
9. Miller DR, Martineau RJ, Wynands JE, Hill J. Bolus administration of esmolol for controlling the haemodynamic response to tracheal intubation: the Canadian Multicentre Trial. *Canadian journal of anaesthesia*. 1991 Oct 1;38(7):849-58.
10. Martineau RJ, Tousignant CP, Miller DR, Hull KA. Alfentanil controls the haemodynamic response during rapidsequence induction of anaesthesia. *Canadian Journal of Anaesthesia*. 1990 Oct 1;37(7):755-61.
11. Ali QE, Siddiqui MMH, Sabzposh SW, Aslam M, Chawla G. Effect of nifedipine pretreatment on the pressure response to laryngoscopy and endotracheal intubation. *Ind Journal Anaesth*, 1999; (43):3 (36).
12. Dullked VK. Efficacy of magnesium sulfate for attenuating the pressor response to laryngoscopy and intubation in comparison with that of lidocaine. *Br Journal of anaesthesia*, 1999 Sep: 499- 506.
13. Mikawa K, Ikegaki J, Maekawa N, Goto R, Kaetsu H, Obara H. The effect of diltiazem on the

- cardiovascular response to tracheal intubation. *Anaesthesia*. 1990 Apr;45(4):289-93.
14. Singh H, Vichitvejpaisal P, Gaines GY, White PF. Comparative effects of lidocaine, esmolol, and nitroglycerin in modifying the hemodynamic response to laryngoscopy and intubation. *Journal of clinical anesthesia*. 1995 Feb 1;7(1):5-8.
 15. Khan FA, Kamal RS. Effect of buprenorphine on the cardiovascular response to tracheal intubation. *Anaesthesia*. 1989 May;44(5):394-7.
 16. Parnass SM, Rothenberg DM, Kerchberger JP, Ivankovich AD. A single bolus dose of esmolol in the prevention of intubation-induced tachycardia and hypertension in an ambulatory surgery unit. *Journal of clinical anesthesia*. 1990 Jul 1;2(4):232-7.
 17. Flacke JW, Bloor BC, Flacke WE, Wong D, Dazza S, Stead SW, Laks H. Reduced narcotic requirement by clonidine with improved hemodynamic and adrenergic stability in patients undergoing coronary bypass surgery. *Anesthesiology*. 1987;67(1):11-9.
 18. Batra YK, Indu B, Puri GD. Attenuation of pulse rate and blood pressure response to laryngoscopy and tracheal intubation by clonidine. *International journal of clinical pharmacology, therapy, and toxicology*. 1988 Jul;26(7):360-363.
 19. Wright PMC, Carbine UA, Meclune S, Orr DA, Moore J. Preanaesthetic medication with clonidine. *Br journal of anesthesiology*. 1990, 65: 628-632.
 20. Nishikawa T, Taguchi M, Kimura T, Taguchi N, Sato Y, Dai M. Effects of clonidine premedication upon hemodynamic changes associated with laryngoscopy and tracheal intubation. *Masui. The Japanese journal of anesthesiology*. 1991 Jul;40(7):1083-8.
 21. Carabine UA, Moore J. Preanaesthetic Medication with Clonidine: A Dose-Response Study. *Survey of Anesthesiology*. 1992 Apr 1;36(2):76.
 22. Gaumann DM, Tassonyi E, Rivest RW, Fathi M, Reverdin AF. Cardiovascular and endocrine effects of clonidine premedication in neurosurgical patients. *Canadian journal of anaesthesia*. 1991 Oct 1;38(7):837-43.
 23. Leslie K, Mooney PH, Silbert BS. Effects of intravenous clonidine: effect of the cardiovascular response to intubation. *Anesth Analg*, 1991; 72: S 327.
 24. Howie MB, Hiestand DC, Jopling MW, Romanelli VA, Kelly WB, McSweeney TD. Effect of oral clonidine premedication on anesthetic requirement, hormonal response, hemodynamics, and recovery in coronary artery bypass graft surgery patients. *Journal of clinical anesthesia*. 1996 Jun 1;8(4):263-72.
 25. Ramesh VJ, Bhardwaj N, Batra YK. Comparative study of oral clonidine and diazepam as premedicants in children. *International journal of clinical pharmacology and therapeutics*. 1997 May; 35(5):218-21.