

“Clinical Profile and Platelet Trend of Dengue Fever <14 Years of Age”

Dr. Morsheda Khanam^{1*}, Dr. Kazi Iman², Dr. Farhana Rahat³, Dr. Mohammed Jamaluddin⁴, Prof. Nobo Krishna Ghosh⁵

¹Assistant Professor, Dr. M R Khan Shishu (Children) Hospital and ICH, Mirpur, Dhaka, Bangladesh

²Registrar, Dr. M R Khan Shishu (Children) Hospital and ICH, Mirpur, Dhaka, Bangladesh

³Assistant Professor, Dr. M R Khan Shishu (Children) Hospital and ICH, Mirpur, Dhaka, Bangladesh

⁴Assistant Professor, Gonosasthaya Samaj Vittik Medical College, Savar, Bangladesh

⁵Professor, Dr. M R Khan Shishu (Children) Hospital and ICH, Mirpur, Dhaka, Bangladesh

DOI: [10.36347/sjams.2019.v07i11.003](https://doi.org/10.36347/sjams.2019.v07i11.003)

| Received: 02.10.2019 | Accepted: 09.10.2019 | Published: 09.11.2019

*Corresponding author: Dr. Morsheda Khanam

Abstract

Original Research Article

Dengue is a mosquito-borne viral disease that has rapidly spread in all regions of world health organization (WHO) in recent years. Dengue virus is transmitted by female mosquitoes mainly of the species *Aedes aegypti* and to a lesser extent, *Aedes albopictus*. Children are the usual victim of dengue infection, there is paucity of published data regarding dengue infection in children in our country. We conducted a cross sectional descriptive in the Department of Paediatrics, Dr. M R Khan Shishu (children) Hospital and ICH, Mirpur-2, Dhaka over a period of 3 months from August 2018 to October 2018 among the children having Dengue infection and who were admitted in the selected hospital. Sixty nine (69) patients were enrolled in our study by using a purposive sampling technique. A detailed history was taken; clinical examinations and relevant investigations were done in every patient. The mean age of the patients was 4.96 ± 5.5 years with almost similar male and female ratio. Most of the patients presented with high grade continued type of fever (75.36%), followed by vomiting (73.91%), abdominal pain (71.01%), Itchy rash was the most important characteristic signs (82.61%). Flushed appearance observed in 66.67%, subconjunctival haemorrhage was (34.78%). Platelet count less than $100 \times 10^9/L$ were found in 44.93% patients. Tourniquet test was positive in 37.68% of cases. All of the patients had packed cell volume (PCV) and Raised serum alanine aminotransferase (ALT) was observed 43.47%. IgM or/with IgG antibodies for dengue virus were positive in 100% patients. Platelet recovery time was 4.12 ± 1.23 days in our observation. Majority (94.20%) of the patients completely recovered from the disease and only 5.80% died. High grade continued fever, vomiting with abdominal pain and itchy skin rash (with normal platelet count) were the presenting features.

Keywords: Dengue Fever (DF), Dengue Haemorrhagic Fever (DHF), Subconjunctival Haemorrhage.

Copyright © 2019: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

INTRODUCTION

Dengue is a mosquito-borne viral disease that has rapidly spread in all regions of world health organization (WHO) in recent years. Dengue virus is transmitted by female mosquitoes mainly of the species *Aedes aegypti* and, to a lesser extent, *Aedes albopictus*. Dengue is the most important arthropod-borne viral disease, and it is a major public health problem in subtropical and tropical regions. The virus is transmitted to humans by the bite of infected female mosquitoes of the genus *Aedes*. The global resurgence of dengue is thought to be due to failure to control the *Aedes* populations, uncontrolled urbanization, population growth, climate change, and increased airplane travel [1]. Dengue is a serious mosquito-borne viral disease which in recent years has become a major

international public health concern. It is the most serious viral haemorrhagic fever in the world with an annual incidence of 100 million cases per year [2]. Of them 250,000 to 500,000 cases are reported as dengue haemorrhagic fever (DHF) (because of the presence of haemorrhagic manifestations, thrombocytopenia and signs of plasma leakage) with an estimated death of about 12,000. The magnitude of dengue fever was largely unknown until it took a heavy toll in 2000 (5555 cases and 93 deaths were reported [3]. Nearly 90% of the dengue infections occur in children with risk of dying during a secondary attack is nearly 15-fold higher than that of adults [4]. Although children are the main group affected by dengue, little published data are available regarding dengue infections in children living in South Asia. In the context of Bangladesh, data of

dengue infection in children are even scarce. Most infections in children fewer than 14 years are asymptomatic or minimally symptomatic; a study of school children in Thailand found only 13% of those infected missed more than one day of school because of illness [5]. Classic dengue is more commonly seen among older children, adolescents, and adults. They are less likely to be asymptomatic. Dengue is abrupt in onset, typically with high fever accompanied by severe headache, incapacitating myalgia and arthralgia, nausea and vomiting, and rash. Rash, typically macular or maculopapular, often becoming confluent and sparing small islands of normal skin, has been reported in over half of infected people. Other signs and symptoms include flushed facies, sore throat, cough, cutaneous hyperaesthesia, and taste aberrations. Recovery may be prolonged and include depression. The present study was carried out in order to document the clinical manifestations of dengue infections in children in Bangladesh.

OBJECTIVE

General objective

To document clinical profile and platelet trend of dengue fever in children in Bangladesh

Specific objectives

To know more about sign and symptoms of dengue fever in Bangladesh

MATERIALS AND METHODS

This cross sectional descriptive study in the Department of Paediatrics, Dr. Mr. Khan Shishu (children) Hospital and ICH, Mirpur-2, Dhaka over a period of 3 months from August 2018 to October 2018. Among the children having Dengue infection and who were admitted in the selected hospital. Sixty nine (69) patients were enrolled in our study by using a purposive sampling technique. In every patient a detailed history was taken, clinical examinations and relevant investigations were done. After obtaining informed consent from the parents a total of 69 children were selected consecutively based on clinical features mentioned in National Guidelines for Clinical Management of Dengue Syndrome, Bangladesh 2005. Patient with any identified specific infection or febrile illness more than two weeks were excluded from the study. Demographic variables, presenting complaints and examination findings were recorded on a structured questionnaire. Tourniquet test was done in predicting feature of bleeding manifestations. NS1 test (at 4th day) was done for all subjects initially. All NS1 positive cases were also positive for IgM. So, NS1 test is very important for detecting dengue virus earlier. White blood cell count (WBC count), platelet count, PCV, ALT, IgM and IgG antibodies for dengue virus were investigated as supporting evidence for dengue infection. Chest X-ray and ultrasonography of whole abdomen were done in selected patients where clinical

findings were suggestive. The WHO classification and case definitions were used to classify disease as DF and DHF3. DHF was further divided into four grades (I, II, III and IV) as per National Guidelines⁷. The test statistics used to analyze the data were descriptive statistics, chi-square (χ^2) test.

RESULTS

Among 69 patients 30 were male and 39 were female. Age ranged from 6 months to 15 years with a mean of 4.96 ± 5.5 years. About sixty (60%) of them was more than 5 years old in male group and about 50% was more than 5 years in female group. Seventy five (75.36%) percent of children complained of fever >5 days with continued type of fever being predominant. Seventy two (72.46%) patients had vomiting and (71.01%) patients had abdominal pain then 62.31% myalgia, 15.49% headache, 14.49% arthralgia, 10.14% retro-orbital pain, 7.24% loose stool and 4.34% runny nose/cough. Among the signs, rash with itching was a salient feature (82.61%) followed by flushed appearance (66.67%), Haemorrhage(69.57%), subconjunctival haemorrhage (34.78%), Pleural effusion (30.43%),hepatomegaly(27.54%), Ascites (18.84%) Melaena (18.84%). About one-third (37.80%) of the patients had positive tourniquet test. Ninety seven (97.10%) had low WBC count. Forty five (44.93%) of patients platelet count with $> 100 \times 10^9/L$, 36.23% with $51- 100 \times 10^9/L$, 23.19% with $21-50 \times 10^9/L$ and 14.49% with $<20 \times 10^9/L$. All children exhibited a packed cell volume (PCV) of less than 45% and over 43.47% had raised serum alanine aminotransferase (ALT). Majority (94.20%) of the patients completely recovered from the disease and only 5.80% died. Platelet recovery time was 4.12 ± 1.23 days in our observation. All death cases had both IgM and IgG positive that means they were affected by dengue 2nd time.




Fig-1: Age and sex distribution of the study participants (n=69)

Mean age of the study participants was 6.5 ± 3.5 years

Table-1: Clinical characteristics of the study participants (n=69)

Characteristics	N (%)
Duration of fever (days)	
<5 days	18 (26.08)
>5 days	51(73.91)
(mean 4.96 ± 5.5 days)	
Type of fever	
Continued	52 (75.36)
Intermittent	13 (18.84)
Remittent	03(4.34)
Biphasic	01 (1.44)
Signs and Symptoms	
Myalgia	43(62.31)
Arthralgia	10(14.49)
Headache	11(15.49)
Retro-orbital pain	7(10.14)
Runny nose/cough	03 (4.34)
Loose stool	05(7.24)
Vomiting	50 (72.46)
Abdominal pain	49 (71.01)
Rash with itching	57 (82.61)
Flushed appearance	46 (66.67)
Signs of shock	09(13.04)
Haemorrhage	48 (69.57)
Gum bleeding	10(14.50)
Subconjunctival haemorrhage	24(34.78)
Petechiae	06 (.70)
Haematemesis	09(13.04)
Melaena	13(18.84)
Pleural effusion	21(30.43)
Hepatomegaly	19(27.54)
Splenomegaly	03 (4.35)
Ascites	13 (18.84)

Table-2: Investigations of the study participants (n=69)

Investigations	Frequencies (%)	
Tourniquet test		
Positive	26 (37.68)	
Negative	56 (81.16)	
Platelet count		
Low WBC (<4x10 ⁹ /l)	67 (97.10)	
> 100X10 ⁹ /L	31(44.93)	
51-100X10 ⁹ /L	25(36.23)	
21-50X10 ⁹ /L	16(23.19)	
< 20X10 ⁹ /L	10(14.49)	
PCV <45%	69(100)	
Raised ALT	30(43.47)	
ICT for dengue	Positive	Negative
NS1	69(100)	00
IgM/IgG	69(100)	00
Platelet recovery time	4.12± 1.23 days	
Outcome		
	Number	Percentage
Recovered	65	94.20
Death	4	5.80

DISCUSSION

This cross sectional study was done to document the clinical findings in dengue infection in Bangladeshi children. It seems to be one of the preliminary efforts of this kind in Bangladesh. The mean age of the patients was 4.96 ± 5.5 days with age range of 6 month to 15 years. Male female ratio was 1:1. Similar results were reported by Malavige *et al.* [7] and Ahmed *et al.* [8], Malavige GN *et al.* [8] found mean age of the patients 7.9±2.9 years and their age range was from 1 month to 12 years [8] while Ahmed *et al.* found mean age 9.0±2.8 years with a age range of 2.5-12 years [8]. A male preponderance with a male female ratio of 3:2 was observed by Ahmed *et al.* [8]. Sixty eight (68.29) percent of children complained of fever >5 days with continued type of fever being predominant (71.95%). Sixty(59.75%) of patients had abdominal pain, 64.63% vomiting, 47.56% myalgia, 18.29% headache, 15.85% arthralgia, 12.19% retro-orbital pain, 10.97% loose stool and 6.09% runny nose/cough. These findings were completely different from that of Rahman *et al.* [8]. Rahman *et al.* had reported headache as the most predominant symptom (91%) followed by myalgia/ arthralgia (85%) and vomiting (64%). Malavige *et al.* had reported runny nose in 20% of patients. Ahmed FU *et al.* found headache in 85%, myalgia in 73%, and retro-orbital pain in 27% & vomiting in 15% of children [7]. Diaz *et al.* had reported abdominal pain precede the onset of plasma leakage in approximately 6% of adults and children with DHF9. Rash with itching was a salient feature (69.51%) followed by flushed appearance (56.09%), Haemorrhage(58.53%), subconjunctival haemorrhage (29.26%), Pleural effusion (25.60%), hepatomegaly(23.17%), Ascites (15.85%) Melaena(15.85%). Sharply contrasting with these findings, Ahmed FU *et al.* found skin rash in only 12% of children. They also found bleeding manifestations as gum bleeding (16%), haematemesis (19%), epistaxis (12%), melaena (8%) and subconjunctival haemorrhage (4%). Rahman *et al.* described spontaneous bleeding in 25% of patients with DF8. A haemorrhagic tendency could be elicited by tourniquet test. In the present study, about 37.8% of the patients had positive tourniquet test. At the initial investigation, most of the cases showed low WBC (leukopenia), representing 81.70% of all cases. All of the patients had packed cell volume (PCV) <45% and raised alanine aminotransferase (ALT) in over 40% of the cases. Platelet count <100x10⁹/L were found in 62.20% children. In another study, the tourniquet test was positive in 47.5% and raised ALT in 49% of children [8]. Ray *et al.* demonstrated alanine aminotransferase (ALT) was abnormal in 50% of patients [12]. Ahmed *et al.* described 38% of the children with positive tourniquet test and 19% with low WBC [8]. In our study, twenty two (58.53%) presented with dengue fever (DF) remaining 38.56% with dengue haemorrhagic fever (DHF). Among them, 35.18% grade-I, 34.24% grade-II, 15.50% grade-III and 15.08 % grade-IV. Malavige GN *et al.* showed 17.3% children

with DHF. Of the DHF, 39.5% of patients had grade-I, 26.7% grade-II, 31.4% grade-III and 2.3% grade IV [8].

LIMITATIONS OF THE STUDY

This is a single centre study with a limited number of samples which can't reflect the scenarios of whole country.

CONCLUSION AND RECOMMENDATION

Most of the children with dengue fever presented with high grade continued fever with vomiting and abdominal pain. Flushed appearance with itchy skin rash and subconjunctival haemorrhage were striking features in our study. Rash with itching was observed as distinctive feature of dengue infection in children. Studies on patterns of paediatric dengue infection in different regions would help clinicians and health administrators to make more informed and evidence-based health planning decisions.

REFERENCES

1. Identification of Dengue Virus in Respiratory Specimens from a Patient Who Had Recently Traveled from a Region Where Dengue Virus Infection Is Endemic. *J Clin Microbiol.* 2007; 45(5): 1523-27.
2. Price DD, Wilson SR. Dengue Fever. Online.
3. 2008 [cited 2009 Jan 21]. Available from: URL:<http://emedicine.medscaDe.com/article/78>
4. World Health Organization. Dengue fever and dengue haemorrhagic fever. [Online], cited 2009
5. Jan 26. Available from: URL: <http://www.whoban.org/comunicable.dis/dengue.html>
6. Burke DS, Nisalak A, Johnson DE, Scott RM. A prospective study of dengue infections in Bangkok. *Am J Trop Med Hyg.* 1988; 38: 172-180.
7. Gibbons RV, Vaughn DW. Dengue: an escalating problem. *BMJ.* 2002; 324:1563-66.
8. Malavige GN, Ranatunga PK, Velathanthiri VGNS, Fernando S, Karunatilaka DH, Aaskov J. Patterns of disease in Sri Lankan dengue patients. *Arch Dis Child.* 2006; 91: 396-400.
9. Ahmed FU, Mahmood BC, Sharma JD, Hoque SM, Zaman R, Hasan MS. Dengue and Dengue Haemorrhagic Fever in Children During the 2000 Outbreak in Chittagong, Bangladesh. *Dengue Bulletin.* 2001; 25: 33-39.
10. Diaz A, Kouri G, Guzman MG. Description of the clinical picture of Dengue hemorrhagic fever/dengue shock syndrome (DHF/DSS) in adults. *Bull Pan Am health Organ.* 1988; 22: 133.
11. Malaria and Vector Borne Diseases Control Unit, Diseases Control Directorate, Directorate General of Health & Family Welfare and World Health Organization. National Guidelines for Clinical Management of Dengue Syndrome, Bangladesh. 2000.
12. World Health Organisation. Prevention and control of dengue and dengue hoemorrhagic fever: comprehensive guidelines. WHO regional publication. SEARO, WHO, 1999; 29.