

The Outcome of the Efficiency of the Condylar Plate for the Treatment of Intercondylar Fracture of Femur

Dr. Md. Shafiqul Islam^{1*}, Dr. Md. Zillur Rahman Siddique², Dr. Md. Mizanur Rahman¹, Dr. Md. Abdus Sobhan³, Dr. Naseem Begam⁴, Dr. A. B. M. Zakir Uddin¹, Dr. Md. Abul Hasnat¹, Dr. Abu Bakar Siddique⁵

¹Assistant Professor, Orthopaedic Surgery, Mugda Medical College, Mugda, Dhaka, Bangladesh

²Assistant Professor, M Abdur Rahman Medical College, Dinajpur 5200, Bangladesh

³Senior Consultant, Orthopaedic Surgery, Adhunik Sadar Hospital, Bangabandhu Rd, Thakurgaon 5100, Bangladesh

⁴Associate Professor, Radiology & Imaging, Shaheed Monsur Ali Medical College & Hospital, Uttara, Dhaka, Bangladesh

⁵Associate Professor, Orthopaedic Surgery, Ad-din Medical College, Dhaka, Bangladesh

DOI: [10.36347/sjams.2020.v08i01.024](https://doi.org/10.36347/sjams.2020.v08i01.024)

| Received: 20.12.2019 | Accepted: 27.12.2019 | Published: 17.01.2020

*Corresponding author: Dr. Md. Shafiqul Islam

Abstract

Original Research Article

Objective: In this study our main goal is to evaluate the outcome of the efficiency of the Condylar plate for the treatment of intercondylar fracture of femur. **Method:** This clinical trial was carried out at the National Institute of Traumatology and Orthopaedic Rehabilitation (NITOR), Dhaka over a period of two years between January 2017 to December 2018. A total of 15 patients of intercondylar fractures of femur meeting the selection criteria were consecutively included in the sample. **Result:** Muller type C₂ was predominant and muller type C₁ and C₃ each was 20%. Over half (53.3%) of the patients were operated after 1 -2 weeks of receiving the injury, 26.7% before 1 week and 20% after 2-3 weeks of injury. Two-thirds (66.7%) of the patients stayed in the hospital for 2-3 weeks and the rest > 3-4 weeks. 40% of the patients had excellent outcome (Insal score > 85), 46.7% good outcome (Insal score between 70-84), and 13.3% fair outcome (Insal score 60- 69). **Conclusion:** The outcome of intercondylar fractures of the femur treated by condylar plate demonstrated excellent to good result in majority of the cases. Sixty percent of the cases returned to routine preinjury activities without limitations and one-quarter with mild limitations, in terms of subjective evaluation as well most of the patients had good to excellent outcome.

Keywords: Condylar plate, intercondylar fracture, Femur.

Copyright @ 2020: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

INTRODUCTION

Supracondylar and intercondylar fractures of the distal femur historically have been difficult to treat. Numerous devices have been proposed and supported and practiced by various authors and centre. Regardless of the implant that has been used for the treatment of distal femoral fracture the principle of internal fixation has to be met. These include proper anatomic reduction of the distal femoral joint surface, stable fixation of the fracture providing proper correction of shortening, angulations and rotational deformities yet permit some stress to allow for rapid and complete healing of fracture, minimal soft tissue stripping and early active mobilization [1-3].

Now most of the intercondylar fractures are being treated by DCS and condylar plate. For DCS a minimum of 4cm of uncomminuted bone in the femoral condyle above the intercondylar notch is necessary for

successful fixation. Whereas multiple whole in the distal part of condylar plate allow multiple screws to be directed into comminuted fractures. However, different investigators claimed dubious results of intercondylar fracture of femur fixation by condylar plate [4, 5].

In this study our main goal is to evaluate the outcome of the efficiency of the Condylar plate for the treatment of intercondylar fracture of femur.

OBJECTIVE

General Objective

- To assess the outcome of the efficiency of the Condylar plate for the treatment of intercondylar fracture of femur.

Specific Objectives

- To detect types of injury.
- To identify operative outcome of the patients

METHODOLOGY

Study Type

- The study was a clinical trial.

Place and Period of Study

- The study was carried out at the National Institute of Traumatology and Orthopaedic Rehabilitation (NITOR), Dhaka over a period of two years between January 2017 to December 2018.

Study population

- Radiologically proven cases of intercondylar fracture of the femur admitted in NITOR were the study population.

Inclusion Criteria

- Adult patients of > 18 years irrespective of sex
- Intercondylar femoral fractures of Muller Type C₁, C₂ and C₃.
- Both closed and open fractures, Gustillo-I, II

Exclusion Criteria

- Patients below 18 years of age
- Pathological fractures
- Intercondylar femoral fractures of Muller Type B₁ and B₂
- Open fractures of Gustillo type IIIA, IIIB and IIIC

Sample Size

- A total of 15 patients of intercondylar fractures of femur meeting the above selection criteria were consecutively included in the sample.

Data Collection

- Data were collected using a structured questionnaire (research instrument) which contained all the variables of interest.

Detailed Procedure

In all cases a detailed history and clinical examination were done. Relevant investigations like complete blood count, random blood sugar and serum creatine were done routinely. Any associated illness such as hypertension, diabetes mellitus, pulmonary problem and concomitant injuries were excluded and if present were treated adequately. X-ray of the affected thigh including hip and knee joints (A/P and lateral view) were performed and selected cases were treated by condylar plate under general/spinal anesthesia.

Data Analysis

Collected data were analysed using software SPSS (Statistical Package for Social Sciences) version 11.5 for windows. Descriptive statistics were used to analyse the data. Analysed data were presented in the form of tables and charts with due interpretation.

RESULTS

In Table-1 shows age distribution of the patients where out of 15 patients, more than half (53.3%) were > 40 years, 20% 30 or < 30 years and 26.7% 30 - 40 years old. The mean age of the patients was 42.6 ± 16.1 years and the lowest and the highest ages were 20 and 70 years respectively. The following table given below in detail:

Table-1: Age distribution of the participants (n = 15)

Age (yrs)	Frequency	Percentage
<30	03	20.0
31-40	04	26.7
>40	08	53.3

Mean age = (42.6 ± 16.1) years; range = (20 - 70) years.

In Figure-1 shows gender distribution of the patients where most of the patients were male, 73%. The following figure is given below in detail:




Fig-1: Gender distribution of the patients

In Figure-2 shows distributions of the patients according to types of injury where muller type C₂ was predominant and muller type C₁ and C₃ each was 20%. The following figure is given below in detail:




Fig-2: Distributions of the patients according to types of injury

In Table-2 shows operative outcome of the patients where over half (53.3%) of the patients were operated after 1 -2 weeks of receiving the injury, 26.7% before 1 week and 20% after 2-3 weeks of injury. Two-thirds (66.7%) of the patients stayed in the hospital for 2-3 weeks and the rest > 3-4 weeks. The following table is given below in detail:

Table-2: Operative outcome of the patients

Time interval between injury and operation	n	%
< 1 week	04	26.7
1-2 weeks	08	53.3
>2-3 weeks	03	20.0
Duration of hospital stay		
2-3 weeks	10	66.7
>3-4 weeks	05	33.3

In Table-3 shows evaluation of outcome by Insal score 9 months after intervention where 40% of the patients had excellent outcome (Insal score > 85), 46.7% good outcome (Insal score between 70-84), and 13.3% fair outcome (Insal score 60-69). The following table is given below in detail:

Table-3: Outcome of patients based on Insal score (n = 15)


Insal score	Frequency	Percentage
> 85 (Excellent)	06	40.0
70 - 84 (Good)	07	46.7
60 - 69 (Fair)	02	13.3

In Table-4 shows outcome of patients based on subjective evaluation where forty percent of the patients rated the outcome as excellent, 46.7% good and 13.3% fair the following figure is given below in detail:

Table-4: Outcome of patients based on subjective evaluation (n = 15)

Subjective evaluation	Frequency	Percentage
Excellent	06	40.0
Good	07	46.7
Fair	02	13.3

In Figure-3 shows outcome of patients based on range of knee motion where 2(13.3%) patients exhibited wide range of knee motion (> 130°), 4(26.7%) patients slightly restricted knee motion (130-110°), 7(46.7%) moderately restricted knee motion (110-90°) and 2(13.3%) extremely restricted knee motion (< 90°). The following figure is given below in detail:

**Fig-3: Outcome of patients based on range of knee motion**

DISCUSSION

The treatment of distal femoral fractures may pose difficult problems. Surgery is the preferred treatment in fractures of the distal femur, which accounts for 4% to 7% of all femoral fractures. Distal femur fractures occur at approximately one-tenth the rate of proximal femur fractures and make up 6% of all femur fractures. There is abimodal distribution of fractures based on age and gender. Most high-energy distal femur fractures occur in males between 15 and 50 years, while most low-energy fractures occur high osteoporotic women >50 years. The most common high-energy mechanism of injury is a traffic accident (53%) and the most common low-energy mechanism is a fall at home (33%) [6].

One study claimed that "fractures in the distal third of the femur continue to perplex the surgeon, whether they are transverse, oblique, or comminuted, or supracondylar or intercondylar in a T, Y or V fashion their management still evokes much controversy because of the consistently poor results obtained.

Muller type C₂ was predominant and Muller type C₁ and C₃ each was 20%. One report demonstrated that 28.6% intercondylar fractures were Muller type C₁, 46.4% Muller type C₂ and 14.3% Muller type C₃ [7]. One study reported that, open methods had good functional results, while only 35.3% of fractures treated by closed methods had good functional results [8].

In the present study, evaluation of outcome by Insal score 12 months after intervention demonstrated that 40% of the patients had excellent outcome (Insal score > 85), 46.7% had good outcome (Insal score between 70-84) and 13.3% had fair outcome (Insal score <70).

In subjective evaluation as well 40% patients rated the outcome as excellent, 46.7% good and 13.3% fair bearing perfect consistency between subjective and objective evaluations. One report found that, excellent outcome in 25%, good in 39.2% and fair in 21.4% which is nearly consistent with our findings.

CONCLUSION

The outcome of intercondylar fractures of the femur treated by condylar plate demonstrated excellent to good result in majority of the cases. Sixty percent of the cases returned to routine preinjury activities without limitations and one-quarter with mild limitations, in terms of subjective evaluation as well most of the patients had good to excellent outcome.

REFERENCES

1. Ali F, Saleh M. Treatment of isolated complex distal femoral fractures by external fixation. Injury. 2000 Apr 1;31(3):139-46.

2. Bolhofner BR, Carmen B, Clifford P. The results of open reduction and internal fixation of distal femur fractures using a biologic (indirect) reduction technique. *Journal of orthopaedic trauma*. 1996 Aug 1;10(6):372-377.
3. Crist BD, Della Rocca GJ, Murtha YM. Treatment of acute distal femur fractures. *Orthopedics*. 2008 Jul 1;31(7):443-445.
4. Butler MS, Brumback RJ, Ellison TS, Poka A, Bathon GH, Burgess AR. Interlocking intramedullary nailing for ipsilateral fractures of the femoral shaft and distal part of the femur. *The Journal of bone and joint surgery. American volume*. 1991 Dec;73(10):1492-502.
5. Butt MS, Krikler SJ, Ali MS. Displaced fractures of the distal femur in elderly patients: operative versus non-operative treatment. *The Journal of bone and joint surgery. British volume*. 1996 Jan;78(1):110-114.
6. Chee SG, Lam KS, Tay BK, Balachandran N. Operative treatment of the intercondylar fracture of the femur. *Singapore medical journal*. 1988 Jun;29(3):276-80.
7. Cieřlik P, Piekarczyk P, Marczyński W. Results of retrograde intramedullary nailing for distal femoral fractures--own experience. *Ortopedia, traumatologia, rehabilitacja*. 2007;9(6):612-617.
8. Dominguez I, Moro ER, De JP, Cebrian JP, López-Durán LS. Antegrade nailing for fractures of the distal femur. *Clinical orthopaedics and related research*. 1998 May(350):74-79.