

## Health and Nutritional Status of Children (0-6 Years age) in Rural Field Practice Area of Katihar Medical College

Inam Binish<sup>1\*</sup>, Inam Adil<sup>2</sup>, Alam Fakhar<sup>3</sup><sup>1</sup>Tutor, Department of Community Medicine, Anugrah Narayan Magadh Medical College, Gaya, Bihar India<sup>2</sup>Specialist Medical Officer, Department of Dermatology, Jai Prakash Narayan Hospital, Gaya, Bihar, India<sup>3</sup>Associate Professor, Department of FMT Katihar Medical College, Katihar Bihar, IndiaDOI: [10.36347/sjams.2021.v09i12.014](https://doi.org/10.36347/sjams.2021.v09i12.014)

| Received: 27.10.2021 | Accepted: 30.11.2021 | Published: 24.12.2021

\*Corresponding author: Dr. Binish Inam

## Abstract

## Original Research Article

**Introduction:** India has a sizeable child population. The 2011 census in the first one in many decades which counted less absolute number of children in the 0-6 years age group compared to 2001 census count of 164 million children, there were 159 million children in 2011. **Materials and Methods:** A cross-sectional study was conducted in the rural field practice area of Katihar Medical College situated around 5 km away from Katihar. A total of 600 under six children were approached for conducting the present study, of which 592 were interviewed and selected for the study. **Results:** Out of 592 children 216 (36.4%) children were exclusively breast feed. Further it was observed that majority i.e. 137 (63.43%) of the children were normal in whom exclusive breast feeding was practiced. It was observed that out of the 548 fully immunized for age children a maximum i.e. 288 (52.55%) were normal. Maximum i.e. 11 (73.33%) of the unimmunized children were under nutrition. Majority of the children were less than 24 months of age, number of female children were slightly higher than male children. Majority of the children were Muslims followed by Hindus and only one child was Christian. Maximum number of children belonged to class IV followed by class III socio-economic class, according to modified Kuppaswamy classification. **Discussion:** According to occupation of the father and nutritional status. Majority of the fathers were involved in unskilled occupation or semiskilled occupation i.e. 356 (60.13%). Unemployed fathers had a higher proportion of under nutrition children 7(58.33%), followed by fathers involved in unskilled /semiskilled work 183(51.4%). Fathers involved in semi-professional/professional occupation had maximum proportion of normal children 7(77.78%). **Conclusion:** Majority of the children belonged to nuclear type of family. The literacy percentage was slightly higher in fathers than in mothers. Majority of fathers were labourers followed by clerical work or owned a shop while most of the mothers were housewives and only few of them worked as teachers or lectures. Maximum children were delivered in hospital. Majority of them had normal birth weight. Maximum children belonged to birth order one. Faulty feeding practices were also observed in this area. Primary Immunization was 92.5% in the study area with small percentage of children not being completely immunized.

**Keywords:** Nutritional Status, Children, Rural area.

Copyright © 2021 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

## INTRODUCTION

India has a sizeable child population. The 2011 census in the first one in many decades which counted less absolute number of children in the 0-6 years age group compared to 2001 census count of 164 million children, there were 159 million children in 2011. This is evident in the share of children in the population which declined from 16% in 2001 to 13.1% in 2011.

As compared to other states, Bihar not only has a higher poverty, but also a significantly higher proportion of children in the population. The children

the age group 0-6 years age account for 18% of the population in Bihar, compared with 13.1% in India, as per the 2011 census.

Early childhood (first 6 years) constitute the most crucial period of life, when the foundations are laid for cognitive, social, and emotional, physical/motor development and cumulative lifelong learning. The age group of 0-6 year's age not only forms a large group but they are also a vulnerable or high risk group, as 90% of growth of human brain and 50% of growth of human body occurs in this period.

**Citation:** Inam Binish *et al.* Health and Nutritional Status of Children (0-6 Years age) in Rural Field Practice Area of Katihar Medical College. Sch J App Med Sci, 2021 Dec 9(12): 1857-1863.

In India Diarrhoeal diseases are major health problem among children under the age of 5 years. Diarrhoea is a leading cause of childhood morbidity and mortality in rural areas and urban slums. Diarrhoea kills 1 million children every year in India and produces malnutrition in children. The median diarrhoeal incident rates in India ranges from 1.0 to 4.7 episodes per child per year. In a prospective study undertaken in rural settings of ICDS block in Rohtak District, it was observed that young children below the age of three years on an average experienced 4 to 6 episodes of Diarrhoea per year. In a study carried out in 1985 in urban and rural areas of 11 states shows that the median Diarrhoea incidents varied from 1.5 episodes per child per year in urban areas to 4.7 episodes in rural area.

Helminthes infestation in children is also very common in rural areas. Incidences of malaria, tuberculosis are also high in rural areas and urban slums.

According to DLH5 – 3 (2007 – 08) data only 41.1% of children (12 to 23 months) are fully immunised in rural area of Bihar. In Katihar District only 32.2% of children belong to the age group 12 to 23 months are fully immunised in rural area, 43.5% of children (12 to 23 months) received measles vaccines in rural areas of Katihar District of Bihar.

The District of Katihar is situated in the plains of North-Eastern part of Bihar state, surrounded by Purnea District in the North, Bhagalpur in the West, Sahebganj District in the South and West Bengal in the East.

Aim of study the prevalence of under-nutrition among the children less than 6 years of age. Determine association of socio-demographic factors with under-nutrition. Association of nutritional status with some epidemiological factors and to suggest measures for prevention and control of under-nutrition based on finding of study.

## MATERIALS AND METHODS

A cross-sectional study was conducted in the rural field practice area of Katihar Medical College situated around 5 km away from Katihar urban area in Department of Community Medicine, Katihar Medical College, Katihar. Ratanpur is small village located in Manihari Block of Katihar District, Bihar. Duration of the study was one year from December 2012 to November 2013.

A total of 600 under six children were approached for conducting the present study, of which 592 were interviewed and selected for the study.

The study participants were the children under the age of six years residing permanently in the rural

field practice area and available during the period of study.

The under six children of both sexes were included in the study. The age was considered in terms of completed years. The following exclusion criteria were used for the purposes of the study:

- A. Children who were seriously ill during the study period.
- B. The under six children who could not be contacted in three visits.

The data collection tool used for the study was an interview schedule that was developed at the institute with the assistance from the faculty members and other experts in relation to the children's anthropometric, dietary, immunisation, morbidity and other relevant parameters used in the study.

## METHOD OF DATA COLLECTION

A. Study was approved by Institutional Ethics Committee and informed verbal consent was obtained from all participants. The health workers informed and motivated the families to participate in the study along with the scope of future intervention, of necessary. All the caregivers of the participants were explained about the purpose of the study and were insured strict confidentiality, and then informed consent was taken from each of them before the total procedure. The participants were given the option not to participate in the study if they wanted. Data regarding family and personal characteristics were recorded by personal interview technique from the caregivers of the participants by the principal investigator himself.

- B. On an average 5-6 children were examined per day.
- C. General information/history was taken.
- D. Clinical examination and analysis of past and present medical records was done.

## RESULTS

The study was conducted to Health and Nutritional status of Children (0 – 6 years) under six year children in rural field practice area of Katihar Medical College (Ratanpur Village). A total of 600 under six children were approached for conducting the present study, of which 592 were interviewed and selected for the study.

According to the WHO recommended classification, the prevalence of underweight (low weight for age), stunting (low height for age), and wasting (low weight for height) was 158 (26.69%), 138 (23.33%), and 96 (16.19%) respectively. Severe degrees of underweight, stunting and wasting were observed in 21.62%, 7.94% and 6.51% respectively. None of the children observed were obese or overweight in the present study.

**Table-1: Age Distribution of Children According To Age Group and Nutritional Status**

Age in months	Under nutrition	Normal	Total
0 – 12	44(28.95%)	108 (71.05%)	152 (100%)
13 – 24	72(47.37%)	80 (52.63%)	152 (100%)
25 – 36	61(51.26%)	58 (48.74%)	119(100%)
37 – 48	54(60.67%)	35 (39.33%)	89(100%)
49 – 60	37(68.52%)	17 (31.48%)	54 (100%)
61 – 72	18(69.23%)	8 (30.77%)	26 (100%)
Total	286 (48.31%)	306 (51.68% )	592 (100%)

Maximum children i.e. 152 each were present in 0 – 12 months and 13 – 24 months age groups. Further it was observed that maximum proportion of under nourished children were in the age group of 61 – 72 months, i.e. 18 (69.23%), followed by 37 (68.52%) in 49 – 60 months and least proportion in 0 – 12 months i.e. 44 (28.95%). This difference was statistically highly significant.

Higher number of girls i.e. 320 was seen in the study. It was observed that 169(52.81%) of the girls and 137(50.37%) of the boys were normal. Among the undernourished maximum number were boys i.e. 135(49.63%). No significant difference was observed between boys and girls as far as under nutrition are concerned ( $p < 0.05$ ).

**Table-2: Distribution of Children According to Socio-Economic Class and Nutritional Status**

Socio economic class	Under nutrition	Normal	Total
Class II	24 (35.82%)	43(64.18%)	67 (100%)
Class III	88(51.16%)	84(48.84%)	172 (100%)
Class IV	163(48.66%)	172(51.34%)	335 (100%)
Class V	11(61.11%)	7(38.89%)	18 (100%)
Total	286(48.31%)	306 (51.68% )	592(100%)

Majority of the children i.e. 335 (56.59%) belonged to class IV followed by 172 (29.05%) in class III and no children were found in class I, according to modified Kuppaswamy classification. Further the table reveals that proportion of under nutrition was highest i.e. 11(61.11%) among children who belonged to class V socio-economic-status followed by class III and class IV, whereas lowest i.e. 24 (35.82%) in children belonging to class II category.

In this study 302 (51.01%) belonged to nuclear type of family followed by 233 (39.36%) from Joint family and least i.e. 57 (9.63%) belonged to three generation families. Proportion of underweight was higher among children from three generation families and nuclear families i.e. 29(50.88 %) and 153(50.66 %) respectively, compared to 104(44.64 %) children from joint families.

Study notes that the maximum numbers of the mothers of study population were housewife i.e. 486 (82.09%). It was observed that children of mothers employed in semi-professional/professional occupations i.e. teachers had high proportion of children being underweight i.e. 4 (66.67%) followed by mothers employed in unskilled/semiskilled mainly as labourers or domestic servants 49 (62.03%). This difference was not statistically significant.

Majority of the fathers were involved in unskilled occupation or semiskilled occupation like labourers or vendors i.e. 356 (60.13%). Unemployed fathers had a higher proportion of undernourished children 7(58.33%), followed by fathers involved in unskilled /semiskilled work 183(51.4%). Fathers involved in semi-professional/ professional occupation had maximum proportion of normal children 7(77.78%). This difference among the various groups was not statistically significant.

In the present study majority i.e 243 (41.04%) of the families had a family size of two. It was observed that proportion of under nutrition was maximum i.e 42 (62.69%) when family size was 4 or more and least i.e 65(43.33%) when family size was one. This difference was not statistically significant.

Study maximum i.e. 533(90.03%) children were delivered in hospital. Proportion of under nutrition was more i.e. 32(54.24%) in children delivered at home compared to those who delivered in hospital i.e. 254(47.65%). The difference was not statistically significant.

In this study 479 (80.91%) had normal birth weight. Maximum number of the children who had birth weight  $< 2.5$  kgs were undernourished i.e. 68(60.18%). This difference was statistically significant.

**Table-3: Distribution of Children According to Birth order and Nutritional Status:**

Birth order	Under nutrition	Normal	Total
1	129(49.43%)	132(50.57%)	261(100%)
2	87(43.17%)	115(56.93%)	202(100%)
3	49(55.68%)	39(44.32%)	88(100%)
> 4	21(51.22%)	20(48.78%)	41(100%)
Total	286(48.31%)	306 (51.68%)	592(100%)

It was observed that the proportion of under nutrition was least in children with birth order two, i.e. 87(43.17%) and was found to be high in children with


birth order three i.e. 49(55.68%). Though this difference was not statistically significant.

**Table-4: Distribution of Children According to Birth Spacing and Nutritional Status**

Birth spacing	Under nutrition	Normal	Total
< 3 years	103(50.74%)	100 (49.26%)	203
> 3 years	61(43.57%)	79(56.43%)	140
Total	164	179	343*

In the present study maximum proportion of under nutrition i.e. 103(50.74%) was seen in children in whom birth spacing was less than 3 years compared to 61(43.57%) in whom birth spacing was 3 or > 3 years. This difference was not statistically significant.

In the present study 361(60.97%) children had been feed with prelacteal feeds. proportion of under nutrition was higher among children who had received prelacteal feeds 200 (55.40 %). This difference was statistically highly significant.



**Graph-01: Distribution of Fewer than Six Children According to Exclusive Breast Feeding Practice and Nutritional Status.**

Graph shows Out of 592 children 216 (36.4%) children were exclusively breast feed. Further it was observed that majority i.e. 137 (63.43%) of the children were normal in whom exclusive breast feeding was practiced. Among the children in whom exclusive breast feeding was not practiced 207(55.05%) were

under nutrition. This difference was statistically highly significant.

It was observed that out of the 548 fully immunized for age children a maximum i.e. 288 (52.55%) were normal. Maximum i.e. 11 (73.33 %) of the unimmunized children were under nutrition. This difference was not statistically significant.

**Table-5: Distribution of Children According to Clinical Signs of under nutrition (N=592):**

Clinical sign		Frequency	Percent
General appearance	Thin	71	11.99%
	Lack of lusture	20	3.37%
Hair	Dyspigmented	1	0.17%
	Thin and sparse	14	2.36%
	Diffuse depigmentation	40	6.75%
Face	Moon face	1	0.17%
	Brown pigmentation	1	0.17%

	Pale conjunctiva	118	19.93%
Tongue	Pale and flabby	97	16.38%
	Geographic	5	0.84%
	Mottled enamel	1	0.17%
Teeth	Caries	83	14.02%
	Dry and scaly	60	10.13%
Skin	Follicular hyperkeratosis	1	0.17%

Study among the different clinical features of under nutrition maximum number of children i.e. 118 (19.93%) had pale conjunctiva, 83(14.02%) had caries, 71 (11.99%) had a thin general appearance and 60 (10.13%) had dry and scaly skin.

## DISCUSSION

All the children under the age of six years, residing permanently in the rural field practice area (Ratanpur village) and available during the period of the study were assessed regarding their socio-economic background, nutritional status, feeding practices, immunisation status, morbidity and other factors affecting the malnutrition. It was assessed by the following method:

In the depth interview of the participants. Clinical examination. Analysis of past & present medical record. The present study shows prevalence of under nutrition in under age six years children. In the present study prevalence of under nutrition was 48.31%. This findings were more or less similar with Sabale Rupali *et al.* [1] 51.8%, Gholamreza Sharifzadeh *et al.* (2010 [2] 47.3%, S Bisai K *et al.* [3] 50.00%, Jakhar *et al.* [4] 58.3 %, M. K. Goel *et al.* [5] 57.4%.

Few studies showed a higher prevalence than the present study namely RN Mishra *et al.* [6] 75%, Munesh Kumar Sharma, *et al.* [7] 72.5%, Shally Awasthi *et al.* [8] 67.3%, Srivastava Anurag *et al.* [9] 66.3%, and SP Mitra [10] 61.1%. According to Paramita Sengupta *et al.* [11] 29.5%, Mittal A *et al.* [12] 38.38% were undernourished. Distribution of age under six year's children according to socio economic status and nutritional status. The proportion of undernutrition was highest i.e. 11(61.11%) among children who belonged to class V socio-economic-status, whereas lowest i.e. 24 (35.82%) in children belonging to Class II category. These findings are in confirmation with Munesh Kumar Sharma *et al.* [7]. According to type of family and nutritional status. It was observed that 302 (51.01%) belonged to nuclear type of family followed by 233 (39.36%) from joint family and least i.e. 57 (9.63%) belonged to three generation. Proportion of under nutrition was higher among children from three generation families and nuclear families i.e. 29(50.88 %) and 153(50.66 %) respectively compared to 104(44.64 %) children from joint families. These findings are in confirmation with Srivastava Anurag, *et al.* [9], Mukhopadhyay DK *et al.* [13]. A cross-sectional study study done by M. K. Goel *et al.* [5] in slum area of Rothak city on 530

children, aged between 1-6 years, revealed that undernourishment was influenced by type of family. Kumkum kumara *et al.* [14] revealed that there was no significant difference between type of family and undernutrition ( $p>0.05$ ). Distribution of age under six year's children according to employment status of the mother and nutritional status. In the present study maximum numbers of the mothers of study population were housewife i.e.486 (82.09%). It was observed that children of mothers employed in semi-professional/professional occupation had high proportion of being under nutrition i.e. 4(66.67%) followed by mothers employed in un skilled /semiskilled 49(62.03%). This difference was not statistically significant.

These findings are in confirmation with Gholamreza Sharifzadeh *et al.* [15] Nguyen Ngoc Hien *et al.* [16] and Nakahara S *et al.* [17].

According to occupation of the father and nutritional status. Majority of the fathers were involved in unskilled occupation or semiskilled occupation i.e. 356 (60.13%). Unemployed fathers had a higher proportion of undernutrition children 7(58.33%), followed by fathers involved in unskilled /semiskilled work 183(51.4%). Fathers involved in semi-professional/professional occupation had maximum proportion of normal children 7(77.78%). This difference among the various groups was not statistically significant.

Paramita Sengupta, *et al.* [11] and Swami HM *et al.* [18] found statistically significant association of being under-nourished and having unskilled labourer father. According to prelacteal feeding and nutritional status. In the present study 361 (60.97%) children had been feed with prelacteal feeds. Proportion of under nutrition was higher among children who had received prelacteal feeds 200 (55.40%). This difference was statistically significant. These findings were in confirmation with Roy S *et al.* [19]. According to exclusive breast feeding practice and nutritional status. Out of 592 children majority i.e. 137 (63.43%) of the children were normal in whom exclusive breast feeding was practiced. Among the children in whom exclusive breast feeding was not practiced 207 (55.05%) were under nutrition. This difference was statistically significant.

These findings were in confirmation with Paramita Sengupta *et al.* [11], Nguyen Ngoc Hien *et al.* [20], Chakraborty S *et al.* [21], Panpanich R *et al.* [22]

and Surya Pathi *et al*. A study done in 2009 by Prema Ramachandran *et al*. [24] states that less than 50 per cent of infants were exclusively breastfed up to six months.

## CONCLUSION

This cross-sectional, community based epidemiological study was conducted in the rural field practice area of Katihar Medical College (Ratanpur village) from November 2012 to December 2013. This study was conducted among under six population aged 0-72 months. Various factors like socio-demography, general health profile, immunization status, feeding practices and anthropometry measurement according to new WHO growth standard (2006) were assessed.

In the present study, majority of the children were less than 24 months of age, number of female children were slightly higher than male children. Majority of the children were Muslims followed by Hindus and only one child was Christian. Maximum number of children belonged to class IV followed by class III socio-economic class, according to modified Kuppaswamy classification. Majority of the children belonged to nuclear type of family. The literacy percentage was slightly higher in fathers than in mothers. Majority of fathers were labourers followed by clerical work or owned a shop while most of the mothers were housewives and only few of them worked as teachers or lecturers. Maximum children were delivered in hospital. Majority of them had normal birth weight. Maximum children belonged to birth order one. Faulty feeding practices were also observed in this area. Primary Immunization was 92.5% in the study area with small percentage of children not being completely immunized. It was observed that 48.2% of the children were undernourished. Parent's literacy, socio-economic status and family size had an impact on better nutritional status of children. Faulty feeding practices, partial immunization, h/o frequent diarrhoea and other infections in the past one year, etc, where some of the epidemiological determinants for undernourishment. Pale conjunctiva carries thin general appearance and dry and scaly skin was most commonly present clinical features of under-nutrition.

### Summary

The first six years of child's life are very critical. In the first moments, month and years of life, every touch, movement and emotion in young child's life translates into an explosion of electrical and chemical activity in the brain and the patterns of a life time are established. In a short 36 months, children develop their ability to think, speak, learn reason and lay the foundation for their values and social behaviour as result.

India has a sizeable child population. Children in the age group of 0-6 years comprises of 13% of total population. As compared to other states, Bihar not only

has higher poverty, but also a significantly higher proportion of children in the population. Children in the age group of 0-6 years account for 19.6% of the population in Bihar, compared with 15.4% in India as a whole.

The World Health Organization (WHO) published the Child Growth Standard for infants and children up to the age of 5 years based on multi country study (Brazil, Ghana, India, Norway, Oman and USA) on growth of healthy breast-fed children under optimal conditions in April 2006. This is a most comprehensive standard than the previous standards but there is very few work applying this standard in India as it is a new tool to the world of anthropometry measurement.

## RECOMMENDATIONS

Proper nutritional care of fewer than six children with special attention of 49-72 months age group children. Mothers should be educated regarding importance of monitoring the growth of their children by maintaining the growth chart. The overall literacy and education level should be increased by proper implementation of literacy and adult literacy campaigns with special emphasis on female literacy. Importance of exclusive breast feeding should be inculcated in the minds of mothers through IEC campaigns. All pregnant women should be encouraged for hospital delivery. All woman coming for antenatal checkups should be educated about the disadvantages of giving prelacteal feeds and advised not to give any forms of prelacteal feeds. All lactating mothers should be educated regarding initiation of weaning at appropriate age. Proper antenatal coverage and care along with special attention towards nutritional status of mother should be given, which will reduce problem of low birth weight and thereby reduce risk of undernutrition during their childhood. As problem of undernutrition was more among higher birth order and with lesser birth interval children, importance of two child norm with adequate birth spacing should be explained through IEC campaigns.

Conflict of interest: Zero.

## REFERENCES

1. Dr. Sabale R., Dr. Kowli, S., Dr. Chowdhary P. Faulty feeding habits and malnutrition in under five children - missed opportunity of education in immunization clinics. *Indian Journal of Maternal and Child Health* 2012; 14(3).
2. Gholamreza, S., Hassan, M., & Soori, R. (2010). Prevalence of Malnutrition in under 6-year Olds in South Khorasan, *Iran, Iran J Pediatr*, 20(4); 435-41.
3. Bisai, S., Bose, K., Ghosh, A. (2008). Nutritional status of Lodha children in a village of Paschim Medinipur district, West Bengal. *Indian Journal of Public Health*, 52(4); 203-6.

4. Jakhar, P., Rajeshwari. (2011). Nutritional Status of Children in Rural Haryana: A Cross-Sectional Analysis. *Indian Journal of Maternal and Child Health*, 13(2).
5. Goel, M.K., Mishra, R., Gaur, D.R., Das, A. (2007). Nutrition Surveillance in 1-6 years Old Children in Urban Slums of a City in Northern India. *The Internet Journal of Epidemiology*, 5, 1.
6. Mishra, R.N., Mishra, C.P., Sen, P., Singh, T. B. (2001). Nutritional Status and Dietary Intake of Pre-School Children in Urban Slums of Varanasi. *Indian Journal of Community Medicine*, 26(2); 90.
7. Sharma, M.K., Kumar, D., Mittal, P.C., Goel, N.K. (2011). Status of Child Undernutrition: Some Socio-Demographic Concerns In Allahabad, Uttar Pradesh. *Indian J. Prev. Soc. Med*, 42(1).
8. Awasthi, S., Das, R., Verma, T., & Vir, S. (2003). Anemia and undernutrition among preschool children in uttar pradesh, india. *Indian Pediatrics*, 40; 985-90.
9. Srivastava, A., Kumar, B., Esam, M.S., Shrotriya, V.P., Mishra, S.P., Iram, S. (2012). Nutritional status of fewer than five children in urban slums of Bareilly Indian *Journal of Maternal and Child Health*, 14(1).
10. Mitra, S. P. (2007). A study of dietary intake and nutritional status of under five children in slums of Kolkata city. *Indian J Community Med*, 32; 92.
11. Sengupta, P., Philip, N., & Benjamin, A.I. (2010). Epidemiological correlates of under-nutrition in under -5 years children in an urban slum of ludhiana. *Health and Population: Perspectives and Issues*, 33; 1: 1-9.
12. Mittal, A., Singh, J., Ahluwalia, S.K. (2007). Effect of maternal factors on nutritional status of 1-5 year old children in urban slum population. *Indian J Community Med*, 32; 264-7.
13. Mukhopadhyay, D.K., Biswas, R., Chakraborty, M. (2009). Anthropometric failure, a new approach to measure undernutrition: an experience from a rural community of West Bengal, India. *J Indian Med Assoc*, 107(4); 211-36.
14. Kumkum, K. (2007). Differentials of Nutritional Status in School-Age Children and the Associated Factors Health and Population-Perspectives and Issues, 30(4); 268-77.
15. Gholamreza, S., Hassan, M., & Soori, R. (2010). Prevalence of Malnutrition in under 6-year Olds in South Khorasan, Iran, *Iran J Pediatr*, 20(4); 435-41.
16. Nguyen, N.H., Sin, K. (2008). Nutritional Status and the Characteristics Related to Malnutrition in Children Under Five Years of Age in Nghean, Vietnam. *J Prev Med Public Health*, 41(4):232-40.
17. Nakahara, S., Poudel, K.C., Lopchan, M. (2006). Availability of childcare support and nutritional status of children of non-working and working mothers in urban Nepal. *Am J Hum Biol*, 18(2):169-81.
18. Swami, H.M., Thakur, J.S., Bhatia, S.P. (2000). National immunization day to assess nutritional status of under-fives in Chandigarh. *Indian J Pediatr*, 67(1); 15-7.
19. Roy, S., Dasgupta, A., Pal, B. (2009). "Feeding practices of children in an urban slum of Kolkata". *Indian J Community Med*, 34; 362-3.
20. Nguyen, N.H., Sin, K. (2008). Nutritional Status and the Characteristics Related to Malnutrition in Children Under Five Years of Age in Nghean, Vietnam. *J Prev Med Public Health*, 41(4); 232-40.
21. Chakraborty, S., Gupta, S., Chaturvedi, B. (2006). A Study of Protein Energy Malnutrition (PEM) in Children (0 to 6 Year) in a Rural Population of Jhansi District (U.P.). *IJCM*, 31(4); 291-2.
22. Panpanich, R., Vipsupakorn, K., Brabin, B. (2003). Breastfeeding and its relation to child nutrition in rural Chiang Mai, Thailand. *J Med Assoc Thai*, 86(5); 415-9.
23. Pathi, S., Das, B.C., Rasanias, H.K., Bhalla, S. (2003). Feeding Practices versus Nutritional Status of Infants in a Rural ICDS Block of Orissa. *Health and Population, Perspectives and Issues*, 26(3);116-20.
24. Ramachandran, P., Gopalan, H.S. (2009). Undernutrition & risk of infections in preschool children. *Indian J Med Res* 2009 November, 130; 579-83.