

## Variants of Ascitic Fluid Bacterial Infection in Children with Chronic Liver Disease in a tertiary care hospital in Bangladesh

Md Liaquat Ali<sup>1\*</sup>, S. M. Quamrul Hassan<sup>2</sup>, Md. Abdul Alim Shaikh<sup>3</sup>, Md. Samsul Alom<sup>4</sup>, Md. Rukunuzzaman<sup>5</sup>, ASM Bazlul Karim<sup>6</sup>, Ahmed Abu Saleh<sup>7</sup>

<sup>1</sup>Associate Professor (cc), Department of Paediatrics, North Bengal Medical College, Sirajganj, Bangladesh

<sup>2</sup>Associate Professor, Department of Paediatrics, Shaheed M. Monsur Ali Medical College, Sirajganj, Bangladesh

<sup>3</sup>Associate Professor & Head, Department of Surgery, North Bengal Medical College, Sirajganj, Bangladesh

<sup>4</sup>Associate Professor & Head, Department of Cardiology, North Bengal Medical College, Sirajganj, Bangladesh

<sup>5</sup>Professor & Chairman Department of Paediatric Gastroenterology and Nutrition, BSMMU, Dhaka, Bangladesh

<sup>6</sup>Professor (Rtd), Department of Paediatric Gastroenterology and Nutrition, BSMMU, Dhaka, Bangladesh

<sup>7</sup>Professor, Department of Microbiology & Immunology, BSMMU, Dhaka, Bangladesh

DOI: [10.36347/sjams.2021.v09i07.016](https://doi.org/10.36347/sjams.2021.v09i07.016)

| Received: 18.04.2021 | Accepted: 27.05.2021 | Published: 29.07.2021

\*Corresponding author: Md Liaquat Ali

### Abstract

### Original Research Article

**Introduction:** Ascitic fluid bacterial infection is an infection of preexisting ascites in the absence of abdominal source. Ascitic fluid bacterial infection may presents in both symptomatic and asymptomatic children with chronic liver disease (CLD). Culture negative neutrocytic ascites (CNNA) is the commonest type of infection in children with CLD. **Objective:** To observe the variants of ascitic fluid bacterial infection in children with CLD. **Methods:** Thirty five consecutive children with clinical features suggestive of CLD, aged between 2-14 years of both sexes with clinically detected ascites and admitted at the Department of Paediatric Gastroenterology and Nutrition, Bangabandhu Sheikh Mujib Medical University (BSMMU) from January 2013 through June 2013 were enrolled in this cross sectional study. Children who had features of ascitic fluid infection like 1 fever, abdominal pain with tenderness were categorized as symptomatic and children who had no features of ascitic fluid infection were categorized as asymptomatic children. Demographic, clinical, haematological, biochemical and ascitic fluid study were done for both symptomatic and asymptomatic children. Variants of ascitic fluid bacterial infection was identified by ascitic fluid polymorphonuclear leucocyte count and culture report. **Results:** Among 35 children, majority 17 (48.6%) were between 6-10 years of age group with male predominance. About 12 (34.3%) were symptomatic and among the symptomatic children 7 (58.4%) were infected and 5 (41.6%) were non-infected, 16 (45.7%) had culture negative neutrocytic ascites (CNNA) variant of ascitic fluid bacterial infection as evident by ascitic fluid polymorphonuclear leucocyte count of  $\geq 250/\text{mm}^3$  and negative culture report. Mean ascitic fluid polymorphonuclear neutrophil count was  $515 \pm 177.82/\text{HPF}$  among infected children and  $85.47 \pm 70.60/\text{HPF}$  among non-infected children, which is statistically significant ( $p < 0.001$ ). Wilson's disease was the commonest 11(30) cause of the CLD in this study. **Conclusion:** Culture negative neutrocytic ascites (CNNA) variety of ascitic fluid infection was the only variety in this study and many of them were asymptomatic.

**Keywords:** Biochemical profile of ascitic fluid, Chronic liver disease, Cytological profile of ascitic fluid.

**Copyright © 2021 The Author(s):** This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

## INTRODUCTION

Patients with chronic liver disease (CLD) are particularly susceptible to infections with a higher prevalence in cirrhotics [1]. Ascites is a frequent complication of cirrhosis. Infections in children with CLD are an important cause of morbidity and mortality. Mechanism of increased susceptibility to infection is unclear [2]. Common bacterial infections in children with CLD are spontaneous bacterial peritonitis (SBP), urinary tract infection (UTI), community acquired pneumonia, dermatologic infections and bacteremia [3].

For the purpose of diagnosis and classification of ascitic fluid infection, culture of the ascitic fluid is essential [4]. Ascitic fluid bacteriological culture is negative in approximately 40 % of adult patients with clinical manifestation suggestive of spontaneous bacterial peritonitis [5]. Ascitic fluid infection is classified in to five categories (including three spontaneous categories) based on ascitic fluid culture results, polymorphonuclear leucocyte counts (PMN), and the presence or absence of a surgical source of infection [4]. Data on ascitic fluid infection (AFI) in children is

**Citation:** Liaquat Ali *et al.* Variants of Ascitic Fluid Bacterial Infection in Children with Chronic Liver Disease in a tertiary care hospital in Bangladesh. Sch J App Med Sci, 2021 July 9(7): 1209-1214.

limited. Among five variants, SBP and Culture negative neutrocytic ascites (CNNA) are common ascitic fluid infections. SBP/CNNA has been reported in 29-43% children with CLD. 11% with ascitic form of acute viral hepatitis (AVH) and 25% with acute liver failure (ALF) [6]. Prompt detection of ascitic fluid infection is helpful to use appropriate antibiotics for treatment of ascitic fluid infection. Ascitic fluid bacterial infection of cirrhotic patients may be symptomatic in 30% of cases [7]. Symptomatic ascitic fluid bacterial infection means patients who have fever, abdominal pain, abdominal tenderness either singly or in combination in patients of liver cirrhosis. Asymptomatic ascitic fluid bacterial infection means patients who have none of these symptoms and signs [8]. Spontaneous bacterial peritonitis (SBP) is defined as infected ascites in absence of recognizable secondary cause of infection [9]. Positive ascitic fluid culture for bacteria was considered essential to establish the diagnosis of Spontaneous Bacterial Peritonitis (SBP). However relying on ascitic fluid culture for diagnosis of SBP has the disadvantages of poor sensitivity and relatively long time before the results are known. To circumvent problem with culture, the ascitic fluid white blood cell (WBC) and Polymorphonuclear leucocyte (PMN) counts have become the standards for making a diagnosis of SBP [10]. The aim of this study was to determine the variants of ascitic fluid bacterial infections in children with chronic liver disease.

## METHODS

This present cross-sectional study was carried out enrolling 35 children clinically suggestive of Chronic Liver Disease (CLD) with ascites in the Department of Paediatric Gastroenterology and Nutrition during the period for January, 2013 through June, 2013. Children who had features of ascitic fluid infection like fever, abdominal pain with tenderness were categorized as symptomatic and children without features of ascitic fluid infection were categorized as asymptomatic. Clinical and laboratory data of both symptomatic and asymptomatic children were recorded. Complete blood count, serum albumin, ALT, AST, Bilirubin, serum Creatinine, Prothombin time, HBsAg (ELISA), anti-HCV (ELISA), Anti LKM1 antibody (ELISA), Liver biopsy, Ultrasonography (USG) and endoscopic findings were recorded whenever necessary for selected children. Ascitic fluid was aspirated with all aseptic precautions. Thirty milliliter of ascitic fluid was collected with an 18 gauze sterile needle attached to a 50 cc disposable syringe. Ascitic fluid culture was done by conventional culture method using Tryptic Soya Broth. Ten milliliter ascitic fluid was used for culture. Five ml of ascitic fluid was sent for total WBC count, differential count and absolute neutrophil count (processed by auto analyzer and finally rechecked manually). Five milliliter ascitic fluid was sent for estimation of total protein and albumin. Demographic, clinical, haematological, biochemical and ascitic fluid

study (cytology, Gram's & Ziehl Neelsen staining, total protein, albumin and culture) were done for both symptomatic and asymptomatic children. Variants of ascitic fluid bacterial infection were identified by ascitic fluid neutrophil count and culture report. Culture Negative Neutrocytic Ascites (CNNA) variety ascitic fluid infection was diagnosed by the presence of ascitic fluid polymorphonuclear leucocyte count of  $\geq 250/\text{mm}^3$  and negative culture report<sup>4</sup>. Presentation difference of both infected and non infected children were found on the basis of difference of presenting symptoms, signs, haematological data and ascitic fluid total WBC count, neutrophil count, ascitic fluid albumin and total protein level.

### Related operational definitions

#### Variants of ascitic fluid infection [4]

##### I. Spontaneous bacterial peritonitis (SBP):

Spontaneous bacterial peritonitis (SBP) means ascitic fluid neutrophil count  $\geq 250/\text{mm}^3$  and positive culture.

##### II. Culture negative neutrocytic ascites (CNNA):

Culture negative neutrocytic ascites (CNNA): Culture negative neutrocytic ascites (CNNA) means ascitic fluid neutrophil counts  $\geq 250/\text{mm}^3$  but culture negative.

##### III. Monomicrobial non-neutrocytic bacterascites:

Monomicrobial non-neutrocytic bacterascites means ascitic fluid neutrophil counts  $< 250/\text{mm}^3$  but culture positive.

##### IV) Polymicrobial bacterascites:

Polymicrobial bacterascites means ascitic fluid neutrophil counts  $< 250/\text{mm}^3$  but multiple organisms' positive culture.

##### V) Secondary bacterial peritonitis:

Secondary bacterial peritonitis means ascitic fluid neutrophil counts  $\geq 250/\text{mm}^3$  and multiple organisms' positive culture.

##### Infected group

Infected group means ascitic fluid neutrophil counts  $\geq 250/\text{mm}^3$  and/or culture positive (SBP or CNNA).

##### Non infected group

Non infected group means ascitic fluid neutrophil counts  $< 250/\text{mm}^3$  and/or culture positive.

## RESULTS

Characteristics of the studied children. Demography of the studied children (n=35). The demographic data of the total 35 children with chronic liver disease were included for this study.

**Age and sex distribution**

It was observed that the age range of the children were from 2 years to 14 years and mean age was  $7.39 \pm 3.0$  years. The highest 48.6% (n-17) incidence of CLD was found in the age group of 6-10 years. A male predominance was observed in the study.

Male were 54.3% (n-19) and female 45.7% (n-16). Out of 35 children, 12 (34.3%) were symptomatic i.e. they had features of ascitic fluid infection like fever, abdominal tenderness, and 23 (65.7%) asymptomatic (Table 1).

**Table-1: Age and gender distribution of the studied subjects (n=35)**

Age (in years)	Gender		Total (%)
	Male	Female	
2-5	07 (36.84%)	05 (31.25%)	12 (34.3%)
6-10	08 (42.1%)	09 (56.25%)	17 (48.6%)
11 and above	04 (21.05%)	02 (12.5%)	06 (17.1%)
<b>Total</b>	<b>19 (54.3%)</b>	<b>16 (45.7%)</b>	<b>35 (100%)</b>

**Compare the ascitic fluid infection in symptomatic and asymptomatic children**

Out of a total 12 symptomatic children, 7 (58.4%) were infected and 5 (41.6%) were non-infected. Out of a total 23 asymptomatic children, 9

(39.1%) were infected and 14 (60.9 %) were non-infected, but this difference is not statistically significant ( $p > 0.05$ ). (Table-2) So, there is no association between ascitic fluid infection and clinical symptoms.

**Table-2: Ascitic fluid bacterial infection in symptomatic and asymptomatic children (n=35)**

Group	Symptomatic (n-12) (34.3%)	Asymptomatic (n-23) (67.7%)	Chi-value	Df	p value
	N %	N %			
Infected	7 58.4	9 39.1	1.15	1	0.279
Non-infected	5 41.6	14 60.9			
<b>Total</b>	<b>12 100</b>	<b>23 100</b>			

**Laboratory findings of the studied children**

Table-3 shows range of ascitic fluid (AF) total WBC count of infected and non-infected children was 312-1200/ $\text{mm}^3$  and 44-312/ $\text{mm}^3$  respectively. Mean AF total WBC count of infected and non-infected children were  $641.62 \pm 231.99 / \text{mm}^3$  and  $146.05 \pm 115.6 / \text{mm}^3$  respectively. Range of ascitic fluid absolute neutrophil

count of infected and non infected children were 255-960/  $\text{mm}^3$  and 10-243/  $\text{mm}^3$  respectively and mean ascitic fluid absolute neutrophil count of infected and non-infected children were  $515 \pm 177.82 / \text{mm}^3$  and  $85.47 \pm 70.6 / \text{mm}^3$  respectively. Mean difference in cell count were statistically significant.

**Table-3: Cytological profile of ascitic fluid (n=35)**

Group	Neutrophil count/ $\text{mm}^3$ (Mean $\pm$ SD) (Range/ $\text{mm}^3$ )	WBC count / $\text{mm}^3$ (Mean $\pm$ SD) (Range/ $\text{mm}^3$ )
Infected group (n-16)	$515 \pm 177.82$ (255-960)	$641.62 \pm 231.99$ (312-1200)
Non infected (n-19)	$85.47 \pm 70.60$ (10-243)	$146.05 \pm 115.60$ (44-312)
<b>p value (T-value, df)</b>	<b>0.001*(9.68, 33)</b>	<b>0.001*(3.84, 33)</b>

\*Significant

Table 4 revealed that range of ascitic fluid total protein (AFTP) of infected and non- infected children was 0.7-2.8 gm/dl and 0.5-2.5 gm/dl respectively. Mean AFTP of infected  $1.21 \pm 0.63$  gm/dl and non infected  $1.07 \pm 0.64$  gm/dl, which was statistically not significant ( $p > 0.05$ ). Range of ascitic albumin of infected and non

infected children was 0.2-1.4 gm/dl and 0.2-0.8 gm/dl respectively. Mean ascitic fluid albumin of infected and non-infected children were  $0.56 \pm 0.35$  gm/dl and  $0.43 \pm 0.19$  gm/dl respectively which was also not significant ( $p > 0.05$ ).

**Table-4: Biochemical profile of ascitic fluid (n-35)**

Group	AFTP (gm/dl) (Mean ±SD)(Range)	AF Albumin (gm/dl) (Mean ±SD)(Range)
Infected group (n=16)	1.21±0.63 (0.7-2.8)	0.56±0.35 (0.2-1.4)
Non-infected (n=19)	1.07±0.64 (0.5-2.5)	0.43±0.19 (0.2-0.8)
<b>p value (T-value, df)</b>	0.51, (0.65, 33)	0.18 (1.35, 33)

**Ascitic fluid culture**

Regarding ascitic fluid culture, among 35 studied children none had culture positive ascitic fluid bacterial infection, though 16 children had Culture Negative Neutrocytic (CNNA) variant ascitic fluid bacterial infection evident by ascitic fluid neutrophil count of  $\geq 250$  cells/mm<sup>3</sup> and negative culture report. In the present study, 16 (45.7%) children had one type of ascitic fluid bacterial infection which is CNNA. Other children had no ascitic fluid bacterial infection (Table 5).

**Variants of ascitic fluid bacterial infections**

A variant of ascitic fluid infection was determined on the basis of ascitic fluid polymorphonuclear neutrophil count and ascitic fluid culture results. Table V shows the variants of ascitic fluid infection in this study. In this present study 16 (45.7%) children had ascitic fluid bacterial infection which was culture negative neutrocytic ascites (CNNA) other children had no ascitic fluid bacterial infection.

**Table-5: Different variants of ascitic fluid bacterial infection in studied children (n-35)**

Type of ascitic fluid bacterial infections	Number	%
Spontaneous Bacterial Peritonitis (SBP)	0	00
Culture-Negative Neutrocytic Ascites (CNNA)	16	45.7
Secondary Bacterial Peritonitis	0	00
Monomicrobial Non-neutrocytic (MNB) Bactericides	0	00
Polymicrobial bacterascites	0	00

**DISCUSSION**

In this small study of 35 children with chronic liver disease with ascites, we found culture negative neutrocytic ascites (CNNA) in 16 (45.7%) children. SBP /CNNA has been reported in 19.5% to 56.6 % children with CLD in another study [11] in children. In the present study out of 35 children with feature of CLD , 12 (34.3%) children were symptomatic i.e. they had features of ascitic fluid bacterial infection like fever, abdominal pain or tenderness and 23 (65.7%) asymptomatic i.e. they had no features of ascitic fluid bacterial infection like fever, abdominal pain or tenderness. Out of a total 12 symptomatic children, 7 (58.4%) were infected and 5 (41.6%) were non-infected. Out of a total 23 asymptomatic children, 9 (39.1%) were infected and 14 (60.9 %) were non-infected, but this difference is not statistically significant ( $p > 0.05$ ). In a study [12] in adult patients showed that out of a total 35 patients of cirrhosis with ascites 8 (22.8%) had CNNA, which is not also statistically significant ( $p > 0.05$ ). In a study [11] in children showed that a total of 28.6% children with liver disease related ascites have CNNA; 50 % were symptomatic. In a other study [13] in children CNNA was found in 16.67% cases. The reason of presence of features of ascitic fluid infection in non infected children may be presence of infection other than ascitic fluid infection like UTI or pneumonia etc. In a study [11] in children fever was the most common symptom in SBP/CNNA, was present in 44% cases and more frequent in SBP/CNNA patients those with no AFI. Other symptoms of AFI (abdominal pain, loose stools

and abdominal tenderness), though more common in SBP/CNNA, did not help in differentiating between patients with AFI and no- AFI.

In this study, most of the children were  $< 10$  years of age and the highest incidence of Chronic Liver Disease (CLD) with ascites was found in the age group of 6-10 years. The age range of the studied children was 2-14 years and the mean age was  $7.39 \pm 3.0$  years. In this study, male were 19 (54.3%) and female 16 (45.7%). This male predominance was also observed in another study done by Sarker [12] and Hossen [13] which were similar to the present study. Ascites polymorphonuclear cells increase with peritoneal infection or with other intra-abdominal inflammatory conditions such as diverticulitis, cholecystitis. In the present study, among the thirty five studied children, 16 children had ascitic fluid neutrophil count of  $\geq 250$  cells/mm<sup>3</sup> which was a diagnostic parameter of Culture Negative Neutrocytic Ascites (CNNA) type of ascitic fluid infection. In a study [12] in adult patient with Chronic liver disease (CLD) with ascites showed that mean ascitic fluid absolute neutrophil count was  $704.50 \pm 480.44$  /mm<sup>3</sup> among infected and  $56.89 \pm 45.26$  /mm<sup>3</sup> among non infected patients. Hossen [13] showed in another study in children that median ascitic fluid neutrophil count was 720/mm<sup>3</sup>, with a range of 360-3600/mm<sup>3</sup> among infected children and median ascitic fluid neutrophil count was 30/mm<sup>3</sup> with a range of 3-192/mm<sup>3</sup> among non-infected children which is statistically significant ( $p$  value=0.001). In the present study, mean ascitic fluid neutrophil count was  $515 \pm 177.82$  /mm<sup>3</sup> among infected

children and mean ascitic fluid neutrophil count among non-infected children was  $85.47 \pm 70.60/\text{cmm}^3$  which was statistically significant ( $p=0.001$ ). So, the findings of the present study in children were similar to findings of the other studies.

In sterile ascites, ascitic fluid white blood cell count is usually less than  $100/\text{mm}^3$  with a predominance of mononuclear cells and a low number of polymorphonuclear cells [15]. In another study [13] median ascitic fluid WBC counts among infected children was  $1200/\text{mm}^3$  with a range of  $600-10,000/\text{mm}^3$  and among non-infected children it was  $100/\text{mm}^3$  with a range of  $20-600/\text{mm}^3$  which is statistically significant ( $p<0.001$ ). In another study [12] in adult patient with CLD with ascites showed that mean ascitic fluid WBC count among infected patients was  $2560 \pm 1871.91/\text{mm}^3$  and mean ascitic fluid WBC count among non infected patients was  $181.85 \pm 105.30/\text{mm}^3$ . In the present study, mean ascitic fluid WBC count among infected children was  $641.62 \pm 231.99/\text{mm}^3$  and among non-infected children it was  $146.05 \pm 115.60/\text{mm}^3$  which was also statistically significant ( $p<0.001$ ).

Patients with ascitic fluid total protein  $< 1\text{gm/dl}$  were the most prone to develop ascitic fluid infection [16]. The opsonic activity of the ascitic fluid was proportional to the ascitic fluid protein concentration [17]. In a study done by Hossen, [13] mean ascitic fluid total protein was  $0.36 \pm 0.23\text{gm/dl}$  and  $1.28 \pm 1.13\text{gm/dl}$  among infected and non-infected children respectively ( $p=0.087$ ). Another study [12] in 35 adult patient with CLD with ascites showed that mean ascitic fluid total protein was  $1.53 \pm 0.61\text{g/dl}$  among infected and  $1.20 \pm 0.59\text{g/dl}$  among non infected patients. In this study mean ascitic fluid total protein was  $1.21 \pm 0.63\text{gm/dl}$  and  $1.07 \pm 0.64\text{gm/dl}$  among infected (CNNA) and non-infected children respectively. So, the mean value of ascitic fluid total protein of infected children of this present study is not similar with the previous study [12], may be due to variation of age of the study populations of these two studies, though the values of the both studies are not statistically significant ( $p>0.05$ ). Hossen [13] showed in another study in children that culture negative neutrocytic ascites (CNNA) is 16.67% In fact, the sensitivity of culture in detecting bacterial growth in neutrocytic ascites (i.e., ascitic fluid with a PMN count greater than or equal to  $250\text{ cells}/\text{mm}^3$  varies widely depending on the method of culture used. In published studies, the conventional method of culture has been found to detect bacterial growth in approximately 50 % of neutrocytic sample [4]. In a recent study [12] in adult patient of CLD with ascites, showed that out of a total 35 patients ascitic fluid culture in conventional method showed no growth of organism, though in his study, out of a total 35 patients 8 (22.8%) had PMN count of  $\geq 250/\text{mm}^3$ . In the present study in children, out of a total 35 children, ascitic fluid culture result was found

negative in all children, though, 16 (45.7%) children had neutrocytic ascites (PMN count  $\geq 250/\text{mm}^3$ ). So, this study was almost similar to the study done in adult patient<sup>12</sup>, Gene probes are now commercially available for the detection of bacteraemia; hopefully, they will lead to rapid (30 minute) and accurate detection of organisms in ascitic fluid [18] In other study [19] in children showed that *Streptococcus pneumoniae* was isolated in 9 of 12 children suffering from cirrhosis with ascites. Another study [20] in adult patients showed that ascitic fluid culture by conventional methods was positive in 46% cases and all the culture positive cases had Gram-negative bacilli: *Escherich coli* being the most common microorganism. Ascitic fluid culture was found negative in the present study; the reason may be that the media used was not enriched enough or due to neutrophil mediated killing of bacteria [17]. On the basis of our facility only aerobic culture was done, though anaerobes can cause ascitic fluid infection rarely (1%). Most episodes of CNNA are diagnosed using insensible cultured methods where there are insufficient numbers of bacteria to reach the threshold of detectability [21]. The conventional method of culture probably requires at least 100 organisms/ml [16] However even when optimal culture methods are used, a small percentage of patients grow no bacteria in their neutrocytic ascitic fluid [19]

## LIMITATIONS

- Time and resource were limited
- Sample size were small
- Only aerobic culture was one
- It was a tertiary care hospital based single Centre study.

## CONCLUSION

Culture negative neutrocytic ascites (CNNA) variety of ascitic fluid infection was the most common variety in this current study. Many of the infected cases are asymptomatic. Apart from fever, symptoms and signs are not always conclusive of the diagnosis and high index of suspicion is needed in all cases with ascites for early diagnosis of cases. Routine culture of the ascetic fluid is not always diagnostic of infection. Cytological parameters of the ascitic fluid will add to the diagnostic accuracy. Polymorphonuclear neutrophil (PMN) cell count was found significantly higher in this group of children. Further studies with larger sample size are necessary to know the actual fact about bacterial infection of ascitic fluid in children with chronic liver disease.

**Conflicts of Interest:** None declared

## REFERENCES

1. Song KH, Jeon JH, Park WB, Park SW, Kim HB, Oh MD, Lee HS, Kim NJ, Choe KW. (2009). Clinical outcomes of spontaneous bacterial

- peritonitis due to extended-spectrum beta-lactamase-producing *Escherichia coli* and *Klebsiella* species. A retrospective matched case - control study. *BMC Infectious Diseases*, 9:41
2. Taneja SK, Dhiman RK. (2011). Prevention and Management of bacterial infections in Cirrhosis, *Int J Hepatol*, 784540.
  3. Cheruvattath R, Balan V. (2007). Infections in patients with end stage liver disease. *J Clin Gastroenterol*, 41: 403-411.
  4. Runyon BA. (2010). Ascites and Spontaneous Bacterial Peritonitis. In: Sleisenger MH, Feldman M, Friedman LS & Brandit LJ. eds. *Sleinger and Fordman's Gastrointestinal and Liver Disease, Pathophysiology/Diagnosis/ Management*. 9<sup>th</sup> ed. Philadelphia: Saunders, p. 1935-1960.
  5. Rimola A, Garcia-Tsao G, Navasa M. (2000). Diagnosis, treatment and prophylaxis of spontaneous bacterial peritonitis, A consensus document. *J. Hepatol*. 32: 142-153.
  6. Poder U, Thapa BR, Prasad A, et al. (2002). Natural history and risk factors in fulminant hepatic failure. *Arch Dis Child*, 87:54-6
  7. Guarner C, Sorian G. (1997). Spontaneous Bacterial Peritonitis, *Hepatology*, 1: 203-217.
  8. Evans LT, KIM WR, Oterucha JJ, Kamnath S. (2003). Spontaneous Bacterial Peritonitis in asymptomatic out patients with Cirrhotic Ascites, *Hepatology*, 37: 897-901.
  9. Syed VA, Ansary JA, Karki P, Regmi M, Khanal B. (2007). Spontaneous bacterial peritonitis (SBP) in cirrhotic ascites: a prospective study in tertiary care hospital, Nepal. *Kathmandu Univ Med J* 5(1):48-59.
  10. Thomas, Boyer. (2003). Diagnosis and management of cirrhotic ascites. In: Zakim D, Boyer TD. (eds). *Hepatology, A text book of liver disease* 5<sup>th</sup>ed. Philadelphia: Saunders, p.639.
  11. Srivastava A, Malik R, Bolia R, Yachha KS, Podder U (2017). Prevalence, Clinical profile, and Outcome of Ascitic Fluid Infection in Children With Liver Disease. *JPGN* 64:194-199.
  12. Sarker JA. (2008). Variants of ascitic fluid bacterial infection in patients of cirrhosis, MD Thesis, Department of Hepatology, BSMMU, Dhaka Bangladesh.
  13. Hossen K, Rukunuzzaman M, Alam R, Benzamin M, Yasmin A, Thakur SB, Sonia ZF, Mazumdar WM, Karim ASMB, et al. (2019). Study of Ascitic Fluid in Children with Chronic Liver Disease in Different Variants of Peritonitis at a Tertiary Care Hospital, Bangladesh *Sch J App Med Sci*, 4: 410.
  14. Garcia-G & Wiest R. (2004). Gut microflora in the pathogenesis of the complications of cirrhosis. *Best Practice and Research: Clinical Gastroenterology*, 18:2.335-372.
  15. Garcia-Tsao, G. (2011). *Ascites', Sherlock's Diseases of the Liver and Biliary System*, 12th ed., Willey- Blackwell, Oxford. UK: 210.
  16. Runyon BA. (1986). Low protein concentration ascitic fluid is predisposed to spontaneous bacterial peritonitis. *Gastroenterology*, 4(1): 396.
  17. Runyon BA. (1988). Patients with deficient opsonic activity are predisposed to SBP. *Hepatology*, 8: 632-635.
  18. Davis TE, Fuller DD. (1991). Direct identification of bacterial isolates in blood cultures using a DNA probe. *J Clin Microbiol*, 29(10): 2193-2196.
  19. Larcher VF, Manolaki N, Vegnente A, Mowat AP. (1985). Spontaneous bacterial peritonitis in children with chronic liver disease: clinical features and etiologic factors, *J Pediatr*, 106: 907-912.
  20. Runyon BA, Hofes JC. (1984). Culture negative neutrocytic ascites; a variant of spontaneous bacterial peritonitis, *Hepatology*, 4: 1209-1211.
  21. Runyon BA. (1990). Monomicrobial non-neutrocytic bacterascites: a variant of spontaneous bacterial peritonitis, *Hepatology*, 12:710.