

Research Article

Global Solutions to elastic string with weak damping by Renormalization Group Method Based on the Classical Theory of Envelope

Wenfeng Sun^{1*}, Lili Bai²

¹College of Petroleum Engineering, Northeast Petroleum University, Daqing 163319, China.

²College of Electrical and information, Northeast Petroleum University, Daqing 163319, China.

***Corresponding author**

Wenfeng Sun

Email: sunwen565656@163.com

Abstract: By the renormalization group method based on the classical theory of envelope, the global approximate solution to damped Fisher equation is obtained.

Keywords: renormalization group; envelope theory; global approximate solutions; damped Fisher equation.

INTRODUCTION

Renormalization group (RG) [1] was initially proposed by Gell-Mann and Low to handle the divergences in the perturbation theory arising in the quantum field theory [2-6], and then many scientists found that the RG method was also powerful in other fields. Recently, Goldenfeld and his colleagues initially applied the RG to nonlinear partial differential equations, such as Barenblatt's equation [7], modified porous medium equation [8], and turbulent-energy-balance equation [9].

Their method is simple and has a lot of applications including singular and reductive perturbation problems in a unified way, and they anticipated that RG method would be useful to perturbed problems. After that, many scientists' work proved it is true [13-15]. Though Goldenfel's presentation of the method is quiet heuristic, heavily relying on the RG prescription in QFT and statistical physics, namely the mathematical foundation of their method has not been established. In fact, why RG equations even in QFT improve naive perturbation has not been elucidated. In paper [10-11], Teiji Kunihiro formulated the renormalization group method based on the theory of envelop and give a mathematical foundation to this method, then he pointed out that the RG equation can be regarded as the envelop equation, and he also proved why the RG equation can give globally improved solutions to ODEs and PDEs. His method is a more direct way to perturbed problems comparing with traditional RG method.

In this paper, we apply Kunihiro's RG method based on the theory of envelope to obtain the global approximate solutions correct up to $O(\varepsilon^2)$ to the elastic string with weak damping [12]

$$u_{xx} = u_{tt} + \varepsilon u_t \quad (1)$$

and we obtain the global solutions to Eq.(1).

Application to Damped Fisher Equation

In this section, we use the Kunihiro's RG method to solve the elastic string with weak damping (1). According to the standard procedure to deal with perturbation problems, we expanding u as $u = u_0 + \varepsilon u_1 + \dots$ gives

$$u_{0xx} = u_{0tt} \quad (2)$$

and

$$u_{1xx} = u_{1tt} + u_0 \quad (3)$$

and so on. The 0-th order solution is given by

$$u_0(x,t) = Ae^{i\lambda(x+t)} + c.c. \quad (4)$$

where c.c. denotes the complex conjugate, λ is the eigenvalue of Eq.(2), and A is a complex constant. If $\lambda \neq 0$, we take formula (4) into Eq.(3), we can get

$$u_{1xx} = u_{1tt} + Ae^{i\lambda(x+t)} + c.c. \tag{5}$$

whose solution is given by

$$u_1 = -\frac{A}{2}(t-t_0)e^{i\lambda(x+t)} \tag{6}$$

Then we have a family of surfaces $S_{t_0}(x, t)$ which is represented by

$$S_{t_0}(x, t) = u(x, t) = Ae^{i\lambda(x+t)} - \frac{\varepsilon A}{2}(t-t_0)e^{i\lambda(x+t)} \tag{7}$$

parametrized by t_0 . We can see it clear that u diverges when t approaches to infinite, so we use the Kunihiro's RG method to eliminate the secular term and get the global solution. Now let us obtain the envelope E of the family of the surfaces of (7). Assuming that A is dependent on t_0 , that is

$$A = A(t_0) \tag{8}$$

According to the standard procedure to obtain the envelope, we solve the equation

$$\frac{\partial u}{\partial t_0} = 0 \tag{9}$$

with the point of tangency at $t_0 = t$, and then we have

$$\frac{dA}{dt_0} + \frac{\varepsilon A}{2} = 0 \tag{10}$$

whose solution is given by

$$A = \bar{A}e^{-\frac{\varepsilon t}{2}} \tag{11}$$

where \bar{A} is a complex constant. Therefore the envelope is represented by

$$u = \bar{A}e^{-\frac{\varepsilon-2i\lambda t}{2}+i\lambda t} + c.c. \tag{12}$$

It is clear to see that we have eliminated the secular term with the RG method and a global approximate solution is obtained.

CONCLUSION

In this paper, we apply the Kunihiro's RG method based on the theory of envelope to the elastic string with weak damping, the result shows that the RG method is simple to eliminate the secular term and give the global approximate solution.

REFERENCE

1. Gell-Mann M, Low FE; Quantum Electrodynamics at Small Distances. Phys. Rev, 1953; 95: 1300.
2. Goldenfeld N, Martin O, Oono Y; Intermediate asymptotics and renormalization group theory. Journal of Scientific Computing, 1989; 4(4): 355-372.
3. Goldenfeld, N. (1992). Lectures on phase transitions and the renormalization group.
4. Chen LY, Goldenfeld N, Oono Y, Paquette G; Selection, stability and renormalization. Physica A: Statistical Mechanics and its Applications, 1994; 204(1): 111-133.
5. Paquette GC, Chen LY, Goldenfeld N, Oono Y; Structural stability and renormalization group for propagating fronts. Physical review letters, 1994; 72(1): 76.
6. Chen LY, Goldenfeld N, Oono Y; Renormalization group theory for global asymptotic analysis. Physical review letters, 1994; 73(10): 1311.
7. Goldenfeld N, Martin O, Oono Y, Liu F; Anomalous dimensions and the renormalization group in a nonlinear diffusion process. Physical review letters, 1990; 64(12): 1361.
8. Chen LY, Goldenfeld N, Oono Y; Renormalization-group theory for the modified porous-medium equation. Physical Review A, 1991; 44(10): 6544.
9. Chen LY, Goldenfeld N; Renormalization-group theory for the propagation of a turbulent burst. Physical Review A, 1992; 45(8): 5572.
10. Teiji K, A Geometrical formulation of the renormalization group method for global analysis Prog. Theor. Phys, 1995; 94: 835.

11. Teiji K, *Jpn. J. Ind. Appl. Math.* 1995;14: 51.
12. Holmes MH; *Introduction to Perturbation Methods*, Springer-Verlag, 1999.
13. Zhu Y, Elim HI, Foo YL, Yu T, Liu Y, Ji W, Sow CH; Multiwalled carbon nanotubes beaded with ZnO nanoparticles for ultrafast nonlinear optical switching. *Advanced Materials*, 2006; 18(5): 587-592.
14. Tu T, Cheng G; Renormalization group theory for perturbed evolution equations. *Physical Review E*, 2002; 66(4): 046625.
15. Bender M, Orszag AA; *Advanced Mathematical Methods for Scientists and Engineers* McGraw-Hill, 1978; 554.