

## Agnesis of Left Hepatic Lobe: A Rare Congenital Anomaly

S. Kirami<sup>1\*</sup>, K. Zitouni<sup>1</sup>, B. Boutakioute<sup>1</sup>, M. Ouali Idrissi<sup>1</sup>, N. Cherif Idrissi El Ganouni<sup>1</sup>

<sup>1</sup>Department of Radiology Arrazi, Mohammed VI university hospital, Cadi Ayad University, Marrakech

DOI: [10.36347/sjmcr.2022.v10i05.009](https://doi.org/10.36347/sjmcr.2022.v10i05.009)

| Received: 05.04.2022 | Accepted: 08.05.2022 | Published: 12.05.2022

\*Corresponding author: S. Kirami

Department of Radiology Arrazi, Mohammed VI university hospital, Cadi Ayad University, Marrakech

### Abstract

### Case Report

Morphologic anomalies of the liver, as opposed to many other visceral organs, are very rare. Left lobe liver agnesis is defined as the absence of liver tissue on the left side of the liver in the absence of previous disease or surgery. It is commonly an accidental finding on imaging investigations or at the time of abdominal surgery. We herein report the case of left liver lobe agnesis diagnosed in a 70-year-old woman with a history of chronic abdominal pain. The diagnosis was confirmed by ultrasonography (US) and enhanced computed tomography (CT) investigations. The main imaging findings of this condition are reviewed briefly, with associations and the most common differential diagnoses.

**Keywords:** Agnesis, Lobe, Liver, Imaging.

**Copyright © 2022 The Author(s):** This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

## INTRODUCTION

Liver lobe agnesis is defined as the absence of liver tissue on the right or left side of the gallbladder fossa, without evidence of previous surgery, trauma or any disease leading to hepatocellular dysfunction and atrophy. It remains a rare condition. Only some cases have been reported in the literature [1].

This condition is generally asymptomatic with normal liver function parameters, and it is generally a coincidental discovery, uncovered by autopsy or imaging examination [2].

## CASE REPORT

A 70-year-old female with a medical history of gallbladder ablation, presented with chronic abdominal pain and discomfort especially in the left upper quadrant. The patient had no liver disease nor ablative surgery or abdominal trauma history. Physical examination was normal. Routine laboratory tests were normal as well as the fibroscopic investigations. As a part of diagnostic work up, an abdominal ultrasound (US) was performed revealing a benign renal right cyst. In the same US examination, the right and caudate lobes of the liver were noticed as hypertrophied and the left liver lobe was not visualized. Liver parenchymal echogenicity, biliary ducts and portal veins on the right lobe were seemingly normal (figure 1). For a more detailed study, contrast enhanced computed tomography (CT) was performed. CT images revealed enlargement

of the right hepatic lobe, especially lateral segment. The left hepatic lobe was absent as well as left hepatic artery, left portal vein, and left hepatic biliary system, and the falciform ligament was no longer individualized (figure2). This appearance pleaded for left liver lobe agnesis associated to compensatory hypertrophy of right hepatic lobe.

## DISCUSSION

Hepatic lobe agnesis is a rare developmental anomaly. Only some cases have been reported in the literature. It is commonly noted incidentally at autopsy, surgery or imaging investigations. Its knowledge is important especially for adequate planning of surgical procedures. This condition was first reported by Arnold and Ashley-Montagu in 1932. The incidence of lobar agnesis has been reported to be 0.005% in 19,000 autopsy cases. The left lobe more affected than the right one [3].

The pathogenesis of hepatic lobe agnesis is owed to defective and interrupted processes during embryonic development of the liver [3].

Clinically most patients are generally asymptomatic with no consequences in prognosis. However, it is important to know whether it is acquired (atrophy) or congenital (agnesis). Some of the affected patients may have additional anomalies and anatomical changes like partial or complete absence of the right

part of the diaphragm, bowel malrotation, choledochal cyst or altered localization of the gallbladder. In addition, a tendency to develop portal hypertension [4].

Pages *et al.* categorized developmental abnormalities of liver morphology as follows: Agenesis (absence of a lobe that is substituted by fibrous tissue); aplasia (one of the lobes is small and its structure is abnormal, with few liver trabeculae, numerous bile ducts and abnormal blood vessels); or hypoplasia (one of the lobes is small but its structure is normal) [4]. According to this classification, our case could be classified as agenesis. The radiologic findings can be detected in US, CT or MRI. The absence of liver tissue to the left side of the gallbladder fossa is the most significant finding for the diagnosis. The absence of the falciform ligament, blood vessels and biliary ducts is also strong evidence of left liver lobe agenesis. The existence of few remanant nodules next to the falciform

hepatic ligament with a signal equivalent to the hepatic parenchyma in MRI are described [5, 6].

In addition to congenital agenesis, trauma, diseases such as liver cirrhosis or previous liver segment resections must also be taken into account in the diagnosis. After excluding the differential diagnoses mentioned above, the rare diagnosis of left lobe agenesis of the liver can be given. In this situation, no further specific therapy is required. However, because of the presence of concomitant hepatic disease (NASH), annual sonographic and laboratory monitoring is recommended [6].


## CONCLUSION

Left hepatic lobe agenesis is a rare congenital anomaly. Awareness about this condition is crucial for clinicians and radiologists. The imaging characteristics should be considered when evaluating a patient with hepatic lobe agenesis.

## Figures



**Fig-1:** liver ultrasonography showing the right and caudate lobes of the liver . The left hepatic lobe was absent as well as left hepatic artery, left portal vein .Liver parenchymal echogenicity, biliary ducts and portal veins on the right lobe were apparently normal.



**Fig-2:** Axial and coronal enhanced computed tomography (CT) showing enlargement of the right hepatic lobe. The left hepatic lobe is absent as well as left hepatic artery, left portal vein, and left hepatic biliary system. The falciform ligament is not individualized.

## REFERENCES

1. Örmeci, T., Erdoğan Telli, S., Örmeci, A., & Aygün, C. (2016). A rare congenital liver anomaly: Hypoplasia of left hepatic lobe.
2. Paech, D., Loi, L., Piecha, F., Bonekamp, D., Delorme, S., Schlemmer, H. P., & Mueller, S. (2019). Fehlen des linken Leberlappens. *Der Radiologe*, 59(6), 550-554.
3. Matsushita, K., Gotoh, K., Eguchi, H., Iwagami, Y., Yamada, D., Asaoka, T., ... & Mori, M. (2017). Agenesis of the left hepatic lobe undergoing laparoscopic hepatectomy for hepatocellular carcinoma: a case report. *Surgical Case Reports*, 3(1), 1-5.
4. Wen, Y. L., Kudo, M., Chung, H., Minami, Y., Suetomi, Y., Onda, H., ... & Maekawa, K. (2001). Agenesis of the left lobe of the liver: Radiologic findings. *Journal of Medical Ultrasonics*, 28(4), 181-183.
5. Ceravolo, I., Guerrieri, D., Macciucca, M. D. V., De Cristofaro, F., & Panzironi, G. (2017). MRI rare finding: Absence of the left liver lobe. *European Journal of Radiology Open*, 4, 50-52.
6. Morgan, M., Bell, D. Agenesis of the left hepatic lobe. Reference article, Radiopaedia.org.