

Accessory Pinna with Accessory Tragus – A Rare Entity**Dr. Ruta Shanmugam¹, Dr. Vidyachal Ravindra², Dr. V. U. Shanmugam³, Dr. R. G. Mariappan⁴,****Dr. Balaji Swaminathan⁵, Dr. Prem Nivas⁶**¹HOD, Dept. of ENT, ²Post Graduate, ^{3,4}Professor, ⁵Reader, ⁶Senior Resident, RMMCH, Annamalai University, Chidambaram, Tamilnadu, India***Corresponding author**

Dr. Vidyachal Ravindra

Email: vidyachal3591@gmail.com

Abstract: The objective is to report a rare case of Accessory Pinna with Accessory Tragus. We present a case of a 13 year old boy who presented with swelling in the left cheek and left pre-auricular region since childhood. We diagnosed the case to be Accessory Pinna with Accessory Tragus. Thorough evaluation was done to rule to presence of associated anomalies. The child was being ridiculed by his peers, hence surgical correction of the accessory pinna and accessory tragus was done.**Keywords:** Accessory Pinna, Accessory Tragus, Congenital Anomaly of Ear.

INTRODUCTION

The pinna develops from the first branchial cleft at 6th week of intra-uterine life by the fusion of six hillocks. Initially the pinna begins to develop in the base of the neck but as the mandible develops it migrates to the normal anatomical position by 20th week of intra-uterine life. Improper development of the external ear leads to congenital anomalies such as Polyotia, which is defined as an accessory pinna that is large enough to resemble an additional pinna. It is a rare congenital anomaly of the external ear. Accessory tragus may be found anywhere along the line of ascent - an imaginary line from tragus to angle of mouth or along anterior border of sternocleidomastoid. Accessory Pinna and Accessory Tragus are caused by failure of proper fusion of the six Hillocks of His during embryological development. It consists of skin, subcutaneous fat and may also contain elastic cartilage. There have been less than 30 reported cases of accessory pinna. Incidence of accessory tragus is approximately 5-10/1000 live births [1]. Management consists of surgical correction only for cosmetic purpose. They are also more prone to have associated renal and cardiac anomalies.

CASE REPORT

A 13 year old boy presented to the ENT OPD with a large Accessory Pinna near his left ear along with Accessory Tragus in the left cheek midway between angle of mouth and tragus (Figure I). His accessory pinna was approximately 2.5 x 2.5x1 cm in size with cartilaginous component. His accessory tragus was approximately 1.0 cm x 0.5 cm in size. Right ear was within normal limits. Rest of ENT examination was within normal limits.

Pure Tone Audiometry, Ultrasound Abdomen, CT Temporal Bone were normal. ECHO showed mild Mitral Valve Prolapse without significant Mitral Regurgitation. The child was being ridiculed by his peers hence wanted surgical correction.

Surgical Correction - Under general anesthesia, both the Accessory Tragus and Accessory Pinna were excised (Figure II & III). Post-operative period was uneventful. Patient was followed by at 1st week, 3rd week, 1st month and at 3rd months (Figure IV).

**Fig 1: Pre-Operative Picture**



Fig 2: Dissection done and Left Accessory Tragus was excised.



Fig 3: Horizontal skin incision was made and dissection done. Cartilage was exposed and excess tissue was excised.



Fig 4: Post-operative Picture – after suture removal on the 7th post-operative day.

DISCUSSION

Accessory pinna or polyotia is an extremely rare condition. It occurs due to failure of proper fusion of the six Hillocks of His during embryological development of the ear. Recent studies suggest that it results from extraordinary migration of neural crest

cells in the branchial arch during development [3]. According to Lammer; fetal exposure to retinoic acid affects the migration of the neural crest cells. It may be single or multiple, unilateral or bilateral.

The first report of polyotia was by von Bol *et al.*; in 1918. Due to rarity of Accessory Pinna, surgical techniques have not yet been established. Yung Moon *et al.*; reported a case of polyotia in 2014 where 9 year old boy presented with a large accessory pinna of size 2.3 x 2 cm. surgical corrections was done in two procedures [3]. The patient had a cartilaginous concave bowl that resembled a conchal hollow behind the accessory pinna. Surgical correction was done in two procedures. In the first procedure, the accessory pinna was resected in a wedge shape. The inferior chondrocutaneous flap was used to construct a new tragus. In the second procedure they eliminated the concha-resembling cartilaginous bowl. Young Raw *et al.*; reported a case of polyotia where surgical correction was done using extraneous tissue and dermal substitute [3].

Accessory pinna has been reported to co-exist with syndromes such as Goldenhar Syndrome, and Treacher Collins Syndrome. Children with accessory tragus are also at a higher risk of renal anomalies. Durakbasa *et al.*; reported that 4% of the children with Accessory Tragi were found to have cardiac and renal anomalies [1]. Thus ultrasound abdomen and ECHO should be mandatorily done.

CONCLUSION

In conclusion, patients presenting with Accessory Pinna and Accessory Tragus should be evaluated thoroughly to rule out presence of any syndromes. Parents should be advised to receive genetic counseling. Surgical correction is done only for cosmetic purpose.

REFERENCES

1. Ulukaya Durakbasa Ç, Mutus M, Meric F, Sirliglu E, Ayhan Y, Dedeoglu S *et al.*; Should children with isolated preauricular tags be routinely evaluated for associated renal and cardiac malformations. *Marmara Medical Journal* 2004; 17 (3): 109-112.
2. Moon I.Y, Oh K.S; Surgical Correction of an Accessory Auricle, Polyotia. *Archives of Plastic Surgery* 2014; 41 (4): 427-429.
3. Rha E.Y, Kim D.H, Byeon J.H; Surgical Treatment of Polyotia. *Archives of Craniofacial Surgery* 2015; 16 (2): 84-87.
4. Kono T, Ro H, Murakami N, Kochi M, Kuroki T, Tosa Y *et al.*; Accessory auricles affecting the tragus and cheek occurring with cervical chondrocutaneous branchial remnants: A Case Report. *JPRAS Open* 2015; 6:20 -24.
5. Chander B, Singh Dogra S, Raina R, Sharma C, Sharma R; Chondrocutaneous Branchial Remnants

or Cartilaginous Choristomas: Terminology, Biological Behavior and Salience or Bilateral Cervical Lesions. *Turk Patoloji Derg* 2014, 30: 195-200.

6. Al Aboud D; Syndromes which may feature accessory tragus: An Overview. *Journal of Pakistan Association of Dermatologists* 2012; 22(3): 262-263.
7. Kar S; Multiple Accessory Tragi: A Case Report. *International Journal of Basic and Applied Medical Sciences* 2013; 3 (1): 1-3.
8. Kohelet D, Arbel E; A prospective search for urinary tract anomalies in infants with isolated preauricular tags. *Pediatrics* 2000; 105(5): e61-e61.